




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/45, C07K 14/115, C12N 5/10, 7/01, A61K 39/155		A1	(11) International Publication Number: WO 98/53078
(21) International Application Number: PCT/US98/10551		(43) International Publication Date: 26 November 1998 (26.11.98)	
(22) International Filing Date: 22 May 1998 (22.05.98)		(74) Agents: KING, Jeffrey, J. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).	
(30) Priority Data: 60/047,575 23 May 1997 (23.05.97) US 60/059,385 19 September 1997 (19.09.97) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE DEPARTMENT OF HEALTH AND HUMANSERVICES [US/US]; Box 071, Bethesda, MD 20892 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MURPHY, Brian, R. [US/US]; 5410 Tuscarawas Road, Bethesda, MD 20816 (US). COLLINS, Peter, L. [US/US]; 12304 Village Square #401, Rockville, MD 20852 (US). DURBIN, Anna, P. [US/US]; 19440 St. Johnsbury Lane, Germantown, MD 20874 (US). SKIADOPOULOS, Mario, H. [US/US]; 8303 Aqueduct Road, Potomac, MD 20854 (US). TAO, Tao [CN/US]; 10684 Weymouth Street #104, Bethesda, MD 20814 (US).		<p>Published</p> <p>With international search report.</p> <p>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	

(54) Title: PRODUCTION OF ATTENUATED PARAINFLUENZA VIRUS VACCINES FROM CLONED NUCLEOTIDE SEQUENCES

(57) Abstract

Isolated polynucleotide molecules provide recombinant PIV genomes and antigenomes for production of recombinant PIV vaccines. The recombinant genome or antigenome can be expressed with a nucleoprotein (N), phosphoprotein (P), and a large (L) polymerase protein to produce isolated infectious PIV particles. The recombinant PIV genome and antigenome can be modified to produce desired changes, for example to incorporate attenuating mutations from biologically derived PIV mutants or to create chimeric PIV clones, to generate attenuated, immunogenic viruses for vaccine use.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

PRODUCTION OF ATTENUATED PARAINFLUENZA VIRUS
VACCINES FROM CLONED NUCLEOTIDE SEQUENCES

10

Related Applications

The present application is a continuation-in-part application of, and claims the benefit under Title 35 of U.S. Provisional Application No. 60/047,575, filed May 23, 1997, and U.S. Provisional Application No. 60/059,385, filed September 19, 1997, each of which is incorporated herein by reference.

20

Background of the Invention

Human parainfluenza viruses (HPIV), HPIV1, HPIV2, and HPIV3 are significant causes of bronchiolitis, croup and pneumonia in infants and children. Karron et al., J. Infect. Dis. 172: 1445-50 (1995); Collins et al. "Parainfluenza Viruses", p. 1205-1243. In B. N. Fields et al., eds., Fields Virology, 3rd ed, vol. 1. Lippincott-Raven Publ., Philadelphia (1996); Murphy et al., Virus Res. 11:1-15 (1988). Infections by these viruses result in substantial morbidity in children less than 3 years of age, and are responsible for approximately 20% of hospitalizations among young infants and children for respiratory tract infections.

Despite considerable efforts to develop effective vaccine therapies against HPIV, no approved vaccine agents have yet been achieved for any HPIV strain, nor for ameliorating HPIV related illnesses. To date, only two live attenuated PIV vaccine candidates have received particular attention. One of these candidates is a bovine PIV (BPIV) strain that is antigenically related to HPIV3, and which has been shown to protect animals against HPIV3. BPIV3 is attenuated, genetically stable and immunogenic in human infants and children (Karron et al., J. Inf. Dis. 171:1107-14

(1995a); Karron et al., J. Inf. Dis. 172:1445-1450, (1995b)). A second PIV3 vaccine candidate, JS cp45 is a cold-adapted mutant of the JS wildtype (wt) strain of HPIV3 (Karron et al., (1995b), supra; Belshe et al., J. Med. Virol. 10:235-42 (1982)). This live, attenuated, cold-passaged (cp) PIV3 vaccine candidate exhibits temperature-sensitive (ts), cold-adaptation (ca), and attenuation (att) phenotypes which are stable after viral replication *in vivo*. The cp45 virus is protective against human PIV3 challenge in experimental animals and is attenuated, genetically stable, and immunogenic in seronegative human infants and children (Hall et al., Virus Res. 22:173-184 (1992); Karron et al., (1995b), supra.

HPIV3 is a member of the Paramyxovirus genus of the Paramyxovirus family, order Mononegavirales. Its genome is a single strand of negative-sense RNA 15462 nucleotides (nt) in length (Galinski et al., Virology 165: 499-510, (1988); Stokes et al., Virus Res. 25:91-103 (1992)) and encodes at least eight proteins (Collins et al., supra, (1996); Galinski, supra, (1991); Spriggs and Collins, J. Gen. Virol. 67: 2705-2719, (1986)). Three of these proteins are associated with the RNA genome to form the nucleocapsid; namely the nucleocapsid protein N, phosphoprotein P, and large polymerase subunit L. Three additional proteins are associated with the envelope, namely the matrix protein M, taught to mediate viral attachment and release, the hemagglutinin-neuraminidase protein HN, and the fusion protein F. Two other proteins, HN and F, represent the neutralizing and protective antigens of PIVs (Collins et al. In Fields Virology, 3rd ed., 1:1205-43 (1996)). Significant sequence divergence in these two protective antigens among different PIVs is the basis for the type specificity of protective immunity against these pathogens (id.).

Another protein of PIV, the C protein, is encoded by an overlapping open reading frame (ORF) of the P protein mRNA (Spriggs and Collins, 1986), and the D protein is generated by RNA editing of the P cistron (Galinski et al. Virology 186:543-50 (1992)). The P mRNA also contains an internal ORF which has the potential to encode a cystein-rich domain called

V. The V ORF is also found in other paramyxoviruses and typically is accessed by RNA editing, but this is not the case with PIV. Presently, it is not known whether the PIV V ORF is expressed.

5 The viral genome of PIV also contains extragenic leader and trailer regions, possessing promoters required for viral replication and transcription. Thus, the PIV genetic map is represented as 3' leader-N-P/C/D-M-F-HN-L-trailer. Transcription initiates at the 3' end and proceeds by a
10 sequential stop-start mechanism that is guided by short conserved motifs found at the gene boundaries. The upstream end of each gene contains a gene-start (GS) signal, which directs initiation of its respective mRNA. The downstream terminus of each gene contains a gene-end (GE) motif which
15 directs polyadenylation and termination.

 Identification of attenuating mutations in *cp45* and other PIV3 vaccine candidates is of interest for a variety of reasons. In particular, it will be useful to understand the genetic basis of attenuation and to characterize the molecular
20 virology and pathogenesis of attenuated HPIV3 strains to provide clinically acceptable vaccines for use against these and other paramyxoviruses, especially HPIV1 and HPIV2 which together account for an additional 7% of pediatric hospital admissions. To achieve these and related goals, a method for
25 producing virus with a wt phenotype from cDNA is needed to determine which mutation(s) in the *cp45* virus specify the *ts*, *ca* and *att* phenotypes and which gene(s) of the BPIV3 specify the attenuation phenotype.

 The complete nucleotide sequences of the prototype
30 PIV3 strain, and of the JS wt HPIV3 and *cp45* strains have been determined (Stokes et al., supra, (1992); Stokes et al., Virus Res. 30: 43-52 (1993)). From these studies, the *cp45* strain was shown to possess at least seventeen nucleotide substitutions compared to the parental JS wt HPIV3 strain,
35 eight of which are correlated with changes to viral proteins. However, it has not been previously shown which of these identified mutations specify desired, e.g., *ts*, *ca*, and *att*, phenotypes. Recently, growth of *cp45* at nonpermissive

temperatures was reported to be complemented by coexpression of a cDNA clone of the L gene of the 47885 wt PIV3 strain (Ray et al., J. Virol. 70:580-584 (1996)), suggesting that the L gene may contain one or more mutations which contribute to the *ts* phenotype of cp45. However, the results of this study are complicated by the fact that the 47885 strain is not isogenic with the JS parent of cp45 (for example, the two viruses are 4% different at the nucleotide level, and the L proteins differ at 41 amino acid positions (Stokes et al., supra, (1992); published erratum appears in Virus Res. 27:96 (1993); Virus Res. 25:91-103. Also, this method of complementation does not provide a clear measurement of the relative contribution of the L gene mutation(s) to the overall *ts* phenotype of cp45.

Rescue and analysis of attenuating mutations in PIV3 and other RNA viruses require effective methods to manipulate cDNAs for the particular viruses of interest. Despite previous advancements identifying cDNAs for PIV, manipulation of the genomic RNA of this and other negative-sense RNA viruses has proven difficult. One major obstacle in this regard is that the naked genomic RNA of these viruses is noninfectious.

Successful methods for direct genetic manipulation of non-segmented negative strand RNA viruses have only recently begun to be developed (for reviews, see Conzelmann, J. Gen. Virol. 77:381-89 (1996); Palese et al., Proc. Natl. Acad. Sci. U.S.A. 93:11354-58, (1996)). Functional nucleocapsids have been successfully generated from the intracellular coexpression of separately transfected plasmids bearing the T7 RNA polymerase promoter and encoding either genomic or antigenomic RNA and the N, P, and L proteins. The intracellular cDNA expression is driven by T7 RNA polymerase which is produced by co-infecting with a vaccinia recombinant virus. This approach was first used to determine the transcription and replication requirements of cDNA-encoded minireplicons. Some success has been achieved in the application of these general methods to rescue infectious rabies virus, vesicular stomatitis virus (VSV), measles virus,

and Sendai virus from cDNA-encoded antigenomic RNA in the presence of the nucleocapsid N, phosphoprotein P, and large polymerase subunit L (Garcin et al., EMBO J. 14:6087-6094 (1995); Lawson et al., Proc. Natl. Acad. Sci. U.S.A.

5 92:4477-81 (1995); Radecke et al., EMBO J. 14:5773-5784 (1995); Schnell et al., EMBO J. 13:4195-203 (1994); Whelan et al., Proc. Natl. Acad. Sci. U.S.A. 92:8388-92 (1995)).

Respiratory syncytial virus (RSV) has also been recovered from cDNA encoded antigenome but this required the transfection of
10 an additional plasmid encoding the M2 ORF 1 transcription elongation factor (Collins et al., 1995).

Rescue of infectious PIV virus and other Mononegavirales members is complicated by virtue of their non-segmented negative-strand RNA genome. The genomic
15 ribonucleoprotein complexes (RNPs) of segmented genome viruses, such as influenza, are generally small in size and loosely structured, and can be assembled *in vitro* from RNA and required viral proteins. However, PIV and other Mononegavirales members feature much larger and more tightly
20 structured RNPs, which tend to be refractory to functional association *in vitro*. Furthermore, the coding potential of HPIV3 P mRNA is complicated by cotranscriptional "RNA editing" (Galinski et al., Virology 186: 543-50 (1992)). The resultant shifts in reading frame can access internal ORFs which are
25 expressed as chimeras fused to the N-terminal part of P. In addition, HPIV3 appears to differ from most other paramyxoviruses which express a chimeric V protein, as noted above. The corresponding set of proteins from HPIV3 editing has not yet been identified, and the internal V ORF of HPIV3
30 is separated from the editing site by numerous translational stop codons (Galinski et al. (1992, supra). Yet another complicating factor is that editing by BPIV3 and HPIV3 produces a novel chimeric protein D, in which the upstream half of P is fused to a domain encoded by a second internal
35 ORF (Pelet et al., EMBO J. 10: 443-448 (1991); Galinski et al., supra, (1992)). The D protein does not have a counterpart in other paramyxoviruses.

In view of the foregoing, an urgent need exists in the art for tools and methods to engineer safe and effective vaccines to alleviate the serious health problems attributable to PIV, particularly illnesses among infants and children attributable to HPIV3. Quite surprisingly, the present invention satisfies these and other related needs.

Summary Of The Invention

The present invention provides novel tools and methods for introducing defined, predetermined structural and phenotypic changes into infectious PIV. In one embodiment of the invention, an isolated polynucleotide molecule is provided which comprises an operably linked transcriptional promoter, a polynucleotide sequence encoding a PIV genome or antigenome, and a transcriptional terminator.

The PIV genome or antigenome can be a human or nonhuman PIV sequence, or a recombinantly modified version thereof. In one embodiment, the polynucleotide sequence encodes a chimeric genome or antigenome comprising a human PIV sequence recombinantly joined with a nonhuman PIV sequence, such as a gene or gene segment from bovine PIV (BPIV). In additional examples, the polynucleotide encodes a chimera of sequences from a nonhuman PIV and at least one other PIV of human or nonhuman origin.

In other embodiments, the invention provides an isolated infectious PIV particle comprising a recombinant PIV (rPIV) genome or antigenome. The isolated infectious PIV particle can be a viral or subviral particle. As used herein, subviral particle refers to any infectious PIV particle which lacks a structural element, eg., a gene segment, gene, protein, or protein functional domain, which is present in a complete virus (eg., an assembled virion including a complete genome or antigenome, nucleocapsid and envelope). Thus, one example of a subviral particle of the invention is an infectious nucleocapsid containing a genome or antigenome, and the products of N, P, and L genes. Other subviral particles are produced by partial or complete deletions or substitutions

of non-essential genes and/or their products (eg., F, HN, M, or C), among other non-essential structural elements.

The isolated infectious PIV particle is preferably a human PIV, more preferably human PIV3 (HPIV3). The invention
5 also provides isolated, infectious particles from bovine or murine PIV (BPIV or MPIV), as well as particles comprising chimeric sequences from two or more different PIV genomes, for example particles incorporating polynucleotide sequences from HPIV3 and HPIV1, from HPIV3 and HPIV2 sequences, or comprised
10 of HPIV3 and BPIV sequences.

In related aspects of the invention, isolated, infectious PIV particles are provided which incorporate nucleotide sequences from HPIV3 joined to at least one sequence from a heterologous PIV, such as HPIV1, HPIV2, BPIV
15 or MPIV. For example, entire genes of HPIV3 may be replaced by counterpart genes from other forms of PIV, such as the HN and/or F glycoprotein genes of PIV1 or PIV2. Alternatively, a selected gene segment, for example a cytoplasmic tail, transmembrane domain or ectodomain of HN or F of HPIV1 or
20 HPIV2, can be substituted for a corresponding gene segment in a counterpart HPIV3 gene to yield constructs encoding chimeric proteins, e.g. fusion proteins having a cytoplasmic tail and/or transmembrane domain of PIV3 fused to an ectodomain of PIV1 or PIV2. Alternatively, genes or gene segments from one
25 PIV can be added (i.e., without substitution) within a heterologous PIV background to create novel immunogenic properties within the resultant clone.

Other modifications can be produced by introducing into a PIV genome or antigenome a nucleotide insertion,
30 rearrangement, deletion or substitution selected to encode a desired phenotypic alteration, such as one that results in attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, improved growth in vitro. or a change in an immunogenic epitope of PIV. In one
35 aspect of the invention, mutations occurring in biologically derived, attenuated PIV are identified and introduced individually or in combination into a full-length PIV clone. Typically these mutations are single amino acid changes

displayed by biologically derived mutant viruses over a wild-type PIV, for example changes exhibited by PIV mutants having *ts*, *ca* or *att* phenotypes. These changes from biologically derived mutant PIV are incorporated into a recombinant PIV clone to specify desired characteristics in the resultant virus. Exemplary mutations include amino acid changes which specify an attenuated phenotype in the HPIV3 strain JS cp45. Among these exemplary mutations are mutations occurring within the PIV polymerase gene L specifying *ts*, *ca* or *att* phenotypes, for example amino acid substitutions occurring at Tyr₉₄₂, Leu₉₉₂, and/or Thr₁₅₅₈ of the JS wild type PIV strain. In more detailed aspects, attenuated PIV recombinants are described wherein Tyr₉₄₂ is replaced by His, Leu₉₉₂ is replaced by Phe, and/or Thr₁₅₅₈ is replaced by Ile.

Also provided within the invention are recombinant PIV having multiple, phenotype-specifying mutations introduced in selected combinations into the genome or antigenome of an infectious clone to yield desired characteristics including attenuation, temperature sensitivity, cold-adaptation, small plaque size, host range restriction, etc. For example, PIV clones are provided which incorporate at least two separate mutations adopted from a biologically derived PIV mutant, e.g., two *ts* mutations from HPIV3 JS cp45. Multiply attenuated viruses are thus obtained by selecting mutations from a "menu" of identified lesions and introducing these mutations in various combinations to calibrate a vaccine virus to selected levels of attenuation, immunogenicity and stability.

In additional embodiments, the invention provides for supplementation of one or more mutations adopted from biologically derived PIV, e.g., *ts*, *ca* or *att* mutations, with additional types of mutations involving the same or different genes. Target genes for mutation in this context include the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, fusion protein F and the C, D and V ORF products. In preferred aspects, attenuating mutations adopted from biologically derived PIV are incorporated within a chimeric

PIV recombinant, e.g., a PIV recombinant having nucleotide sequences from both HPIV3 and HPIV1, or from both HPIV and BPIV viruses.

In other embodiments, the invention provides methods for producing an infectious PIV particle, e.g., a viral or subviral particle, from one or more isolated polynucleotide molecules encoding a PIV genome or antigenome (see also copending United States provisional patent application No. 60/047575, filed May 23 1997, incorporated herein by reference in its entirety). To produce an infectious PIV particle according to these methods, an expression vector comprising an isolated polynucleotide molecule encoding a PIV genome or antigenome is coexpressed in a cell or cell-free system with an expression vector comprising one or more isolated polynucleotide molecules encoding N, P, and L proteins of a PIV, whereby an infectious PIV particle is produced.

The PIV genome or antigenome and the N, P, and L proteins may be coexpressed by a single expression vector, or by separate expression vectors. In alternate embodiments, the N, P, and L proteins are each encoded on separate expression vectors.

Within the aforementioned methods, the polynucleotide molecule encoding the PIV genome or antigenome may correspond to a genomic or antigenomic sequence of human, bovine or murine PIV. Alternatively, the PIV encoding polynucleotide may be a chimera of a human PIV genomic or antigenomic sequence and at least one non-human PIV genomic or antigenomic sequence. In additional methods for producing infectious PIV, the polynucleotide encoding the PIV genome or antigenome is a chimera of two or more human PIV genomes, for example a polynucleotide containing sequences from HPIV3 joined to sequences from one or more related forms of human PIV, such as human PIV1 or human PIV2. Individual genes of human PIV3 may be substituted by counterpart genes from heterologous PIV, for example the HN and F glycoprotein genes of PIV1 or PIV2, to yield a modified genome or antigenome encoding a chimeric PIV. Alternatively, a selected, heterologous gene segment, such as a cytoplasmic tail,

transmembrane domain or ectodomain of HN or F of HPIV1 or HPIV2, can be substituted for a counterpart gene segment in a different PIV type or different gene, e.g., HN or F of HPIV3, to yield constructs encoding chimeric proteins, e.g. fusion proteins having a cytoplasmic tail and/or transmembrane domain of PIV3 fused to an ectodomain of PIV1 or PIV2.

In yet additional methods for producing infectious PIV, the PIV genome or antigenome is modified to yield a chimera of a human PIV genomic or antigenomic sequence and at least one non-human PIV sequence, for example a polynucleotide containing sequences from both human and bovine PIV.

In other methods for producing infectious PIV, the PIV genome or antigenome is modified by a nucleotide insertion, rearrangement, deletion or substitution selected to encode a desirable phenotypic alteration, such as one that results in attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV. Alternatively, the polynucleotide molecule encoding the PIV genome or antigenome can be modified to encode non-PIV molecules, e.g., a cytokine, a T-helper epitope, a restriction site marker, or a protein of a different microbial pathogen (e.g., virus, bacterium or fungus) capable of eliciting a protective immune response in the intended host. In one embodiment, the PIV genome or antigenome is modified to encode protein from a human RSV or from measles virus.

In other embodiments of the invention a cell or cell-free expression system (e.g., a cell-free lysate) is provided which incorporates an expression vector comprising an isolated polynucleotide molecule encoding a PIV genome or antigenome, and an expression vector comprising one or more isolated polynucleotide molecules encoding N, P, and L proteins of a PIV. Upon expression, the genome or antigenome and N, P, and L proteins combine to produce an infectious PIV particle, such as a viral or subviral particle. The isolated polynucleotide molecules encoding the PIV genome or antigenome and the one or more isolated polynucleotide molecules encoding N, P, and L proteins of PIV can be expressed by a single

vector, or the genome and one or more of the N, P, and L proteins can be incorporated into two or more separate vectors.

5

Brief Description of the Drawings

Figure 1 illustrates construction of the plasmid p218(131), encoding a complete copy of HPIV3 genomic RNA. The diagram of p218(A*A'CC'D'E) (top left) illustrates the location of the T7 terminator and delta ribozyme adjacent to nucleotide 1 of the HPIV3 genome. The top right diagram of p(E*FF'GHIJKL*) illustrates the locations of the 7 sequencing markers (asterisk), see Fig. 2 below, and the T7 promoter adjacent to nucleotide 15462 of HPIV3. The 15 overlapping clones used to construct the full length genomic cDNA are indicated by letters on the outside of the two plasmids. The unique restriction sites *Ngo*MI and *Xho*I were used to clone p218(131) which encodes the complete negative-sense genomic RNA of HPIV3 illustrated at the bottom. The location of each sequencing marker in p218(131) differentiating recombinant JS from JS wt is indicated by an asterisk.

20

Figure 2 depicts sequence markers in cDNA encoded genomic RNA of the HN (1) and L(2) genes of HPIV3. Sequences are negative sense, mutations are boxed.

1) A non-coding nt substitution at 7903 removes a *Hga* I site from JS wt. Nucleotide substitutions in JS cDNA at 7913 and 7915 create a *Sca* I site and change amino acid 370 from proline to threonine, ablating the antibody epitope recognized by mAb 170/7 and 423/6. Also, the HN gene was found to contain an additional point mutation at nt 7593 which had not been recognized previously. This results in a threonine to isoleucine change at amino acid position 263 in the HN protein.

30

2) The three incidental non-coding mutations in the L gene of JS cDNA that occurred during plasmid construction or assembly are boxed.

35

Figure 3 includes a diagram (not to scale) of the PIV3-CAT cDNA (top, drawn as a circular plasmid) and the

encoded minigenome RNA (bottom, drawn as a single strand of negative-sense RNA, 3' to 5').

Plasmid: The T7 promoter is shown as a black arrow pointing in the direction of transcription. Positions are shown for the delta ribozyme, the T7 transcriptional terminator, the inserted vaccinia virus transcriptional terminators (T₇CT and T₇AT), restriction enzyme sites, and the coding sequence for the various domains of the encoded minigenome.

Minigenome RNA: The 5' (right hand) end of the minigenome is defined by the promoter for T7 RNA polymerase, which contributes an extension of two nonviral G residues. These are not included in length calculations. The 3' end is defined by the ribozyme and is free of any heterologous nucleotides. PIV3-specific regions are shown as open boxes. The CAT ORF is shaded and the negative-sense complements of the translational initiation (UAC) and termination (AUU) codons are shown. Nucleotide positions are indicated according to the shortest, 898-nucleotide minigenome. Nucleotide lengths of specific regions are given in parentheses. The sequence of positions 90 to 124 (3' to 5', negative-sense) is shown at the bottom; PIV3-specific sequence is underlined, the complement of a vaccinia virus transcriptional terminator is overlined, the complement of the XbaI site is italicized, and the complement of the translational start codon for the CAT ORF is in bold. The site where 0 to 6 G residues were inserted, resulting in PIV3-CAT 0 to +6, is indicated. Abbreviations: GS, transcriptional gene-start motif; GE, transcriptional gene-end termination/polyadenylation motif; NT, nontranslated gene sequence.

Figure 4 depicts expression of CAT enzyme by minigenomes PIV3-CAT 0 to +6. Cells were transfected with plasmids encoding the indicated PIV3-CAT minigenome and the N, P and L proteins, and were infected simultaneously with vaccinia virus recombinant vTF7-3 expressing T7 RNA polymerase. Lysates were prepared 48 h post-infection, and samples representing 3000 cells were analyzed for CAT

activity. Activities are expressed relative to the +2 minigenome, which is a multiple of six, as 100. Samples in which the L plasmid was omitted did not have CAT activity detectable under these conditions.

5 Figures 5A and 5B depict synthesis of
mini-antigenomic RNA by PIV3-CAT minigenomes 0 to +6. Cells
were transfected and infected as described for Fig. 4, except
for controls in which N plasmid (lane 1) or L plasmid (lane 2)
was omitted. The PIV3-CAT minigenome plasmid which was
10 transfected in each case is indicated. Cell lysates were
prepared at 48 h post-transfection: one aliquot was treated
with micrococcal nuclease (treated) and the second was
mock-treated (untreated), followed by processing for RNA
purification. The RNAs were analyzed by Northern blot
15 hybridization with a negative-sense CAT riboprobe. As a
marker and control for nuclease treatment, the RSV-CAT C2
minigenome (Grosfeld et al., J. Virol. 69:5677-5686 (1995),
incorporated herein by reference in its entirety) was
complemented by the RSV N, P and L proteins and was processed
20 in parallel (Lane 10). Autoradiograms of the hybridized blots
are shown in Fig. 5A. Fig. 5B shows phosphorimager analysis
in which the amount of hybridization relative to the +2
minigenome was calculated separately for the treated and
untreated samples.

25 Figures 6A and 6B depict synthesis of polyadenylated
mRNA by PIV3-CAT minigenomes 0 to +6. Using aliquots of cells
from the same experiment as shown in Figs. 5A and 5B, total
intracellular RNA was harvested at 48 h post-transfection and
fractionated by oligo(dT) chromatography into bound and
30 unbound fractions, which were analyzed by Northern blot
hybridization with a negative-sense CAT riboprobe. Fig. 6A
shows autoradiograms of the hybridized blots. Fig. 6B shows
the phosphorimager analysis in which the amount of
hybridization was normalized relative to the bound RNA
35 fraction of the +2 minigenome, this was calculated separately
for the bound and unbound fractions.

Figures 7A and 7B depict accumulation of
intracellular minigenome in response to plasmid encoding

PIV3-CAT minigenomes 0 to +6. This is a continuation of the experiment shown in Figs. 5A and 5B. Thus, cells which had been transfected with plasmids encoding a minigenome and the N, P and L proteins and infected with vTF7-3 were harvested 48 h post-transfection and cell lysates were prepared. One aliquot of lysate was treated with micrococcal nuclease (treated) and the other was mock-treated (untreated), followed by processing for RNA purification. The RNAs were analyzed by Northern blot hybridization with a positive-sense CAT riboprobe, whereas in Fig. 5A the riboprobe was negative-sense. Fig. 7A shows autoradiograms of the hybridized blots. Fig. 7B shows phosphorimager analysis in which the amount of hybridization relative to the +2 minigenome was calculated separately for the treated and untreated samples.

Figures 8-10 illustrate construction of p3/7(131)2G. p3/7(131)2G encodes a complete positive-sense HPIV3 antigenome and contains two G residues between the T7 promoter and nucleotide 1 of the antigenome. The construction involved modification separately of the left and right halves of cDNA 218(131) encoding the negative-sense genome. Using the same strategy, a second cDNA, p3/7(131) was constructed which is identical to p3/7(131)2G except that the two G residues were omitted.

Figure 8 depicts construction of p(Left+2G), performed by replacing the ribozyme and T7 transcription terminator of p218(A*A'CC'D*E) with a T7 promoter including two G residues. p218 (A*A'CC'D*E) was used as the template in a PCR that amplified the left-hand 1224 nucleotides of the HPIV3 genome (black rectangle labeled "PIV3seq.") using a left hand PCR primer that introduced the T7 promoter. This PCR fragment was cloned into the MluI-PmlI window of p218(A*A'CC'D*E), resulting in p(Left+2G).

Figure 9 depicts construction of p(Right+), performed by replacing the T7 promoter of p(E*FF'GHIHKL*) with a ribozyme and T7 terminator placed adjacent to PIV3 nucleotide 15642 in the positive-sense (antigenome) strand. p(E*FF'GHIHKL*) was used as a template in a PCR that amplified

the right-hand 649 nucleotides of the HPIV3 genome (black rectangle labeled "PIV3 seq.") using a mutagenic oligonucleotide that added part of the delta ribozyme (including a naturally-occurring RsrII site). This PCR product was cloned into the RsrII-BssHII window of p3/7 to yield pPIV3-3/7, thus reconstructing a complete ribozyme flanked by the T7 terminator. The SwaI-NgoMI fragment of pPIV3-3/7 was cloned into the SwaI-NgoMI windows of p(E*FF'GHIHKL*), resulting in p(Right+). The locations of the seven sequence markers are indicated with asterisks.

Figure 10 depicts construction of p3/7(131)2G. The NgoMI-XhoI fragment of p(Right+) was cloned into the NgoMI-XhoI window of p(Left+2G), resulting in p3/7(131)2G. The positions of HPIV3 genes are indicated (not to scale). The locations of sequence markers are indicated with asterisks.

Figure 11 shows sequence confirmation of a negative sense recombinant PIV3. A 1379 bp fragment (nucleotides 7334-8713) spanning the mutations at 7903, 7913, and 7915 was generated by RT-PCR of RNA from infected cells and then analyzed by cycle-sequencing. The mutations differentiating recombinant PIV from JS wt are indicated by arrows. The complete sequence from nt 7897 to 7922 is shown in the margin next to each gel, with the three nucleotide differences indicated in bold.

Figure 12 provides a map of plasmid pTM(L)942/992/1558, which contains the PIV3 L cDNA with amino acid substitutions at positions 942, 992, and 1558 in the L protein sequence. The relative position of each of coding change is indicated, together with the aa difference and the naturally-occurring restriction site which was deliberately ablated as a marker. Restriction sites used for cloning (*Sph*I, *Pvu*AI, *Bam*HI and *Nhe*I) are indicated. The arrow shows the direction of the L protein coding sequence and is numbered according to amino acid position.

Figure 13 is a schematic representation of recombinant PIV3 viruses bearing representative mutations within the invention.

Figure 14 illustrates construction of cDNA encoding the chimeric PIV3-PIV1 antigenome in which the PIV3 HN and F ORFs were replaced by those of PIV1. First (starting from the bottom left) the PIV3 F and HN genes were subcloned (as *BspE*-I

5 I-*Xho* I and *EcoR* I-*Spe* I fragments) from the full-length PIV3 cDNA clone p3/7(131)2G (Panel A) to generate pLit.PIV3.F3a and pLit.PIV3.HN4 (Panel B), respectively. PCR mutagenesis (Panel C) was performed to delete the complete coding regions of the PIV3 F and HN genes and to introduce new restriction sites

10 (boxed). PIV1 F and HN cDNAs were prepared from infected-cell RNA by RT-PCR and cloned into pLITMUS28 (Panel D). These clones, pLit.1Fwt and pLit.1HNwt, were used as templates for modification of the PIV1 F and HN genes at their start and stop codons to introduce new restriction sites compatible with

15 those introduced in the PIV3 deletions (Panel E). Chimeric F and HN genes were constructed by importing the PIV1 coding regions into the PIV3 deletions to generate pLit.PIV3-1.Fhc and pLit.PIV3-1.HNhc (Panel F). The chimeric F and HN were assembled together to generate pSE.PIV3-1.hc (Panel G). The F

20 and HN chimeric segment was introduced into full-length PIV3 clone p3/7(131)2G by replacing its *BspE* I-*Sph* I fragment to generate pFLC.2G+.hc (Panel H). The small boxes between genes represent the gene end, intergenic, and gene start regions, and the lines represent the non-coding regions. Shaded

25 portions are from PIV1, open boxes are from PIV3. The black arrows depict the T7 promoter, while black boxes depict the hepatitis delta virus (HDV) ribozyme.

Figure 15A is a diagram of the chimeric HN and F genes of the chimeric rPIV3-1 (middle) in parallel with that

30 of rPIV3/JS (top; alternatively referred to herein as rJS and JS wt rPIV3) and PIV1/Wash64 (bottom). The four junction regions containing the sequence transitions from PIV3 to PIV1 are boxed and numbered I to IV. Each small box between genes represents the gene end, intergenic, and gene start region, and the lines represent the non-coding regions. RT-PCR

35 primers, specific to PIV3 M and L genes (primer pair A at top) or specific to PIV1 M and HN genes (primer pair B at bottom) used in Figure 14B, are depicted as arrows.

Figure 15B shows an evaluation of RT-PCR products generated using virion RNA (vRNA) as template and the PIV3- or PIV1-specific primer pairs described in Figure 14A. The reverse transcription step was omitted in a duplicate of each template, as indicated, to confirm that the PCR product was derived from RNA. For rPIV3-1, the PIV3-specific primer pair A gave rise to the expected 4.6 kb product, whereas primer pair B, specific to PIV-1 sequences not present in rPIV3-1, did not yield product. Positive controls using rPIV3/JS (labeled as rPIV3) or PIV1/Wash64 (labeled as PIV1) vRNA are shown in parallel.

Figure 16 represents sequences of PIV3-PIV1 junctions in the RT-PCR products of rPIV3-1 shown in Figure 14B were determined. The sequence for each of the four junction regions (Regions I-IV) is presented and aligned with the corresponding regions of rPIV3/JS (top line) and PIV1/Wash64 (bottom line), which were sequenced in parallel from RT-PCR products. Vertical bars indicate sequence identity, and the boxed regions indicate introduced mutations and restriction sites. The Gln to Glu codon change in the chimeric F gene is indicated by shaded box. Start and stop codons are underlined.

Figure 16 shows sequencing gels for region III (left) and IV (right) of RT-PCR products of rPIV3-1 compared with rPIV3 (left) or PIV1 (right). Start and stop codons are marked by a box, restriction sites are marked by arrows.

Figure 18 depicts multicycle growth of parental and chimeric PIVs in tissue culture. LLC-MK2 cell monolayers were inoculated with virus at an MOI of 0.01, and virus-infected cells were incubated at 32°C in the presence of trypsin. Tissue culture supernatants were harvested at 24 hour intervals, frozen, and analyzed in the same TCID₅₀ assay using hemadsorption to identify virus-infected cultures. Each point represents the mean titer of three separate cultures, with S.E. Indicated. The dotted horizontal line indicates the lower limit of viral detection.

Figure 19 illustrates introduction of the three L gene mutations of cp45 into pFLC.2G+.hc, the antigenomic cDNA

clone of the chimeric virus rPIV3-1. pTM(L)942/992/1558 (A) is a plasmid clone of the L gene that carries the three mutations found in cp45. The mutation at amino acid position 942 in the PIV3 L protein is a tyr. (wt) to his (cp45) substitution and a nearby Eae I site was ablated to mark this site. Similarly, the 992 mutation is a leu to phe change with a Bsr I site ablated and the 1558 mutation is a thr to ile change with an Ava II site ablated. The 2.9 kb SphI-NheI fragment present in pTM(L)942/992/1558 was introduced into pFLC.2G+hc (B), the plasmid carrying the full length cDNA clone of rPIV3-1, to give pFLC+hc.cp45L (C). For the constructs in (B) and (C), the black boxes indicate the location of the hepatitis D virus ribozyme and the T7 terminator, the shaded regions are the PIV1 HN and F ORFs, and the open boxes represent sequences derived from PIV3. pFLC.2G+.hc.cp45L was used in the transfection to yield the attenuated chimeric recombinant virus designated rPIV3-1.cp45L.

Description of the Specific Embodiments

The present invention provides compositions and methods for producing and modifying infectious PIV from isolated polynucleotide molecules, preferably cDNA. Infectious PIV particles are produced by a recombinant coexpression system that permits introduction of defined changes into infectious PIV. These modifications are useful in a wide variety of applications, including the development of live attenuated vaccine strains bearing predetermined, defined attenuating mutations.

Infectious PIV of the invention are produced by intracellular or cell-free coexpression of one or more isolated polynucleotide molecules that encode a PIV genome or antigenome RNA, together with one or more polynucleotides encoding viral proteins necessary to generate a transcribing, replicating nucleocapsid. Among the viral proteins useful for coexpression to yield infectious PIV are the major nucleocapsid protein (N) protein, nucleocapsid phosphoprotein (P), large (L) polymerase protein, fusion protein (F),

hemagglutinin-neuraminidase glycoprotein (HN), and matrix (M) protein. Also useful in this context are products of the C, D and V ORFs of PIV.

5 cDNAs encoding a PIV genome or antigenome are constructed for intracellular or *in vitro* coexpression with the necessary viral proteins to form infectious PIV. By "PIV antigenome" is meant an isolated positive-sense polynucleotide molecule which serves as a template for synthesis of progeny PIV genome. Preferably a cDNA is constructed which is a
10 positive-sense version of the PIV genome corresponding to the replicative intermediate RNA, or antigenome, so as to minimize the possibility of hybridizing with positive-sense transcripts of complementing sequences encoding proteins necessary to generate a transcribing, replicating nucleocapsid.

15 In some embodiments of the invention the genome or antigenome of a recombinant PIV (rPIV) need only contain those genes or portions thereof necessary to render the viral or subviral particles encoded thereby infectious. Further, the genes or portions thereof may be provided by more than one
20 polynucleotide molecule, i.e., a gene may be provided by complementation or the like from a separate nucleotide molecule. In other embodiments, the PIV genome or antigenome encodes all functions necessary for viral growth, replication, and infection without the participation of a helper virus or
25 viral function provided by a plasmid or helper cell line.

By "recombinant PIV" is meant a PIV or PIV-like viral or subviral particle derived directly or indirectly from a recombinant expression system or propagated from virus or subviral particles produced therefrom. The recombinant
30 expression system will employ a recombinant expression vector which comprises an operably linked transcriptional unit comprising an assembly of at least a genetic element or elements having a regulatory role in PIV gene expression, for example, a promoter, a structural or coding sequence which is
35 transcribed into PIV RNA, and appropriate transcription initiation and termination sequences.

To produce infectious PIV from a cDNA-expressed PIV genome or antigenome, the genome or antigenome is coexpressed

with those PIV N, P and L proteins necessary to (i) produce a nucleocapsid capable of RNA replication, and (ii) render progeny nucleocapsids competent for both RNA replication and transcription. Transcription by the genome nucleocapsid provides the other PIV proteins and initiates a productive infection. Alternatively, additional PIV proteins needed for a productive infection can be supplied by coexpression.

Synthesis of PIV antigenome or genome together with the above-mentioned viral proteins can also be achieved *in vitro* (cell-free), e.g., using a combined transcription-translation reaction, followed by transfection into cells. Alternatively, antigenome or genome RNA can be synthesized *in vitro* and transfected into cells expressing PIV proteins.

In certain embodiments of the invention, complementing sequences encoding proteins necessary to generate a transcribing, replicating PIV nucleocapsid are provided by one or more helper viruses. Such helper viruses can be wild type or mutant. Preferably, the helper virus can be distinguished phenotypically from the virus encoded by the PIV cDNA. For example, it is desirable to provide monoclonal antibodies which react immunologically with the helper virus but not the virus encoded by the PIV cDNA. Such antibodies can be neutralizing antibodies. In some embodiments, the antibodies can be used in affinity chromatography to separate the helper virus from the recombinant virus. To aid the procurement of such antibodies, mutations can be introduced into the PIV cDNA to provide antigenic diversity from the helper virus, such as in the HN or F glycoprotein genes.

In alternate embodiments of the invention, the N, P, L and other desired PIV proteins are encoded by one or more non-viral expression vectors, which can be the same or separate from that which encodes the genome or antigenome. Additional proteins may be included as desired, each encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome. Expression of the genome or antigenome and proteins from transfected plasmids can be achieved, for example, by each cDNA being under the control of a promoter

for T7 RNA polymerase, which in turn is supplied by infection, transfection or transduction with an expression system for the T7 RNA polymerase, e.g., a vaccinia virus MVA strain recombinant which expresses the T7 RNA polymerase (Wyatt et al., Virology, 210: 202-205 (1995), incorporated herein by reference in its entirety). The viral proteins, and/or T7 RNA polymerase, can also be provided by transformed mammalian cells or by transfection of preformed mRNA or protein.

A PIV antigenome may be constructed for use in the present invention by, e.g., assembling cloned cDNA segments, representing in aggregate the complete antigenome, by polymerase chain reaction or the like (PCR; described in, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202, and *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, San Diego (1990), each incorporated herein by reference in its entirety) of reverse-transcribed copies of PIV mRNA or genome RNA. For example, a first construct is generated which comprises cDNAs containing the left hand end of the antigenome, spanning from an appropriate promoter (e.g., T7 RNA polymerase promoter) and assembled in an appropriate expression vector, such as a plasmid, cosmid, phage, or DNA virus vector. The vector may be modified by mutagenesis and/or insertion of synthetic polylinker containing unique restriction sites designed to facilitate assembly. For ease of preparation the N, P, L and other desired PIV proteins can be assembled in one or more separate vectors. The right hand end of the antigenome plasmid may contain additional sequences as desired, such as a flanking ribozyme and tandem T7 transcriptional terminators. The ribozyme can be hammerhead type (e.g., Grosfeld et al., (1995), *supra*), which would yield a 3' end containing a single nonviral nucleotide, or can be any of the other suitable ribozymes such as that of hepatitis delta virus (Perrotta et al., Nature 350:434-436 (1991), incorporated herein by reference in its entirety) which would yield a 3' end free of non-PIV nucleotides. The left- and righthand ends are then joined via a common restriction site.

A variety of nucleotide insertions, deletions and rearrangements can be made in the PIV genome or antigenome during or after construction of the cDNA. For example, specific desired nucleotide sequences can be synthesized and
5 inserted at appropriate regions in the cDNA using convenient restriction enzyme sites. Alternatively, such techniques as site-specific mutagenesis, alanine scanning, PCR mutagenesis, or other such techniques well known in the art can be used to introduce mutations into the cDNA.

10 Alternative means to construct cDNA encoding the genome or antigenome include reverse transcription-PCR using improved PCR conditions (e.g., as described in Cheng et al., Proc. Natl. Acad. Sci. USA 91:5695-5699 (1994)), incorporated
15 herein by reference) to reduce the number of subunit cDNA components to as few as one or two pieces. In other embodiments different promoters can be used (e.g., T3, SP6) or different ribozymes (e.g., that of hepatitis delta virus. Different DNA vectors (e.g., cosmids) can be used for propagation to better accommodate the larger size genome or
20 antigenome.

Isolated polynucleotides (e.g., cDNA) encoding the genome or antigenome may be inserted into appropriate host cells by transfection, electroporation, mechanical insertion, transduction or the like, into cells which are capable of
25 supporting a productive PIV infection, e.g., HEP-2, FRhL-DBS2, LLC-MK2, MRC-5, and Vero cells. Transfection of isolated polynucleotide sequences may be introduced into cultured cells by, for example, calcium phosphate-mediated transfection (Wigler et al., Cell 14: 725 (1978); Corsaro and Pearson, Somatic Cell Genetics 7: 603 (1981); Graham and Van der Eb, Virology 52: 456 (1973)), electroporation (Neumann et al., EMBO J. 1: 841-845 (1982)), DEAE-dextran mediated transfection (Ausubel et al., (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY (1987), cationic lipid-mediated
30 transfection (Hawley-Nelson et al., Focus 15: 73-79 (1993)) or a commercially available transfection reagent, e.g., LipofectACE® (Life Technologies, Gaithersburg, MD) or the like
35

(each of the foregoing references are incorporated herein by reference in its entirety).

As noted above, in some embodiments of the invention the N, P, L and other desired PIV proteins are encoded by one or more helper viruses which is phenotypically distinguishable from that which encodes the genome or antigenome. The N, P, L and other desired PIV proteins can also be encoded by one or more expression vectors which can be the same or separate from that which encodes the genome or antigenome, and various combinations thereof. Additional proteins may be included as desired, encoded by its own vector or by a vector encoding one or more of the N, P, L and other desired PIV proteins, or the complete genome or antigenome.

The compositions and methods of the invention permit analysis and manipulation of PIV molecular biology and pathogenic mechanisms using, e.g., defined mutations to alter the function or expression of selected PIV proteins. Using these methods and compositions, one can readily distinguish mutations responsible for desired phenotypic changes from silent incidental mutations, and select phenotype-specific mutations for incorporation into a recombinant PIV genome or antigenome for vaccine production.

Modifications of PIV provided within the invention are directed toward the production of improved vaccine viruses, e.g., to enhance viral attenuation and vaccine immunogenicity, to ablate epitopes associated with undesirable immunopathology, to accommodate antigenic drift, etc. To achieve these and other objectives, the compositions and methods of the invention allow for a wide variety of modifications to be introduced into a PIV genome or antigenome for incorporation into infectious, recombinant PIV. For example, foreign genes or gene segments encoding protective antigens or epitopes may be added within a PIV clone to generate PIV virus strains capable of inducing immunity to both PIV and another virus or agent from which the protective antigen was derived. Alternatively, foreign genes may be inserted, in whole or in part, encoding modulators of the

immune system, such as cytokines, to enhance immunogenicity of a vaccine virus.

Other mutations which may be included within PIV clones of the invention include, for example, substitution of
5 heterologous genes or gene segments (e.g., a gene segment encoding a cytoplasmic tail of a glycoprotein gene) with a counterpart gene or gene segment in a PIV clone.

Alternatively, the relative order of genes within a PIV clone can be changed, the PIV genome promoter can be replaced with
10 its antigenome counterpart, or selected gene(s) rendered non-functional (e.g., by functional ablation involving introduction of a stop codon to prevent expression of the gene). Other modifications in a PIV clone can be made to facilitate manipulations, such as the insertion of unique
15 restriction sites in various non-coding or coding regions or elsewhere. In addition, nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

Thus, by providing infectious clones of PIV the invention permits a wide range of alterations to be
20 recombinantly produced within the PIV genome (or antigenome), yielding defined mutations which specify desired phenotypic changes. By "infectious clone" is meant cDNA or its product, synthetic or otherwise, RNA capable of being directly incorporated into infectious virions which can be transcribed
25 into genomic or antigenomic RNA capable of serving as a template to produce the genome of infectious viral or subviral particles. As noted above, defined mutations can be introduced by a variety of conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or
30 antigenome. The use of genomic or antigenomic cDNA subfragments to assemble a complete genome or antigenome cDNA as described herein has the advantage that each region can be manipulated separately, where small cDNA subjects provide for better ease of manipulation than large cDNA subjects, and then
35 readily assembled into a complete cDNA. Thus, the complete antigenome or genome cDNA, or a selected subfragment thereof, can be used as a template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a

single-stranded phagemid form, such as using the MUTA-gen® kit of Bio-Rad Laboratories (Richmond, CA), or a method using the double-stranded plasmid directly as a template such as the Chameleon® mutagenesis kit of Strategene (La Jolla, CA), or by the polymerase chain reaction employing either an oligonucleotide primer or a template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome cDNA. A variety of other mutagenesis techniques are known and can be routinely adapted for use in producing the mutations of interest in a PIV antigenome or genome cDNA of the invention.

Thus, in one illustrative embodiment mutations are introduced by using the MUTA-gene® phagemid in vitro mutagenesis kit available from Bio-Rad Laboratories. In brief, cDNA encoding an PIV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform CJ236 cells (Life Technologies). Phagemid preparations are prepared as recommended by the manufacturer. Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome is then amplified.

Mutations can vary from single nucleotide changes to the introduction, deletion or replacement of large cDNA segments containing one or more genes or genome segments. Genome segments can correspond to structural and/or functional domains, e.g., cytoplasmic, transmembrane or ectodomains of proteins, active sites such as sites that mediate binding or other biochemical interactions with different proteins, epitopic sites, e.g., sites that stimulate antibody binding and/or humoral or cell mediated immune responses, etc. Useful genome segments in this regard range from about 15-35 nucleotides in the case of genome segments encoding small functional domains of proteins, e.g., epitopic sites, to about 50, 75, 100, 200-500, and 500-1,500 or more nucleotides.

The ability to introduce defined mutations into infectious PIV has many applications, including the manipulation of PIV pathogenic and immunogenic mechanisms.

For example, the functions of PIV proteins, including the N, P, M, F, HN, and L proteins and C, D and V ORF products, can be manipulated by introducing mutations which ablate or reduce the level of protein expression, or which yield mutant protein. In one such exemplary modification, a sequence at the cleavage site of the F protein can be modified and the effects of this modification on growth in tissue culture and infection and pathogenesis of the resultant PIV can be routinely determined in experimental animals.

Various genome RNA structural features, such as promoters, intergenic regions, and transcription signals, can also be routinely manipulated within the methods and compositions of the invention. The effects of trans-acting proteins and cis-acting RNA sequences can be readily determined, for example, using a complete antigenome cDNA in parallel assays employing PIV minigenomes (Dimock, et al., J. Virol. 67: 2772-8 (1993), incorporated herein by reference in its entirety), whose rescue-dependent status is useful in characterizing those mutants that may be too inhibitory to be recovered in replication-independent infectious virus.

The present invention further provides tools and methods to readily distinguish between silent incidental mutations and mutations responsible for phenotype differences, for example by introducing suspect mutations, separately and in various combinations, into the genome or antigenome of infectious wild-type (i.e., for one or more phenotypic character such as temperature sensitivity, replication in a selected host, etc.) PIV. This process permits identification of mutations responsible for desired vaccine phenotypes such as attenuation, temperature sensitivity, cold-adaptation, small plaque size, host range restriction, etc. Mutations identified by these methods can then be introduced in various combinations to modify a vaccine virus to an appropriate level of attenuation, etc., as desired. Moreover, the present invention provides the ability to combine mutations from different strains of virus into a single vaccine strain.

As noted above, mutations incorporated within recombinantly altered PIV clones may be selected based on

their ability to alter expression and/or function of a selected PIV protein, yielding a desired phenotypic change, or for a variety of other purposes. Desired phenotypic changes include, e.g., changes in viral growth in culture, temperature sensitivity, plaque size, attenuation, and immunogenicity. For example, a polynucleotide sequence encoding the genome or antigenome can be modified by a nucleotide insertion, rearrangement, deletion or substitution to specify attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, alteration in gene expression, or a change in an immunogenic epitope.

In one aspect of the invention, mutations occurring in biologically derived, attenuated PIV are identified and introduced individually or in combination into a full-length PIV clone, and the phenotypes of rescued recombinant viruses containing the introduced mutations are determined. In exemplary embodiments, amino acid changes displayed by biologically derived mutant viruses over a wild-type PIV, for example changes exhibited by PIV mutants having *ts*, *ca* or *att* phenotypes, are incorporated within recombinant PIV clones. These changes from biologically derived mutant PIV specify desired characteristics in the resultant clones, e.g., an attenuation phenotype specified by a mutation adopted from the HPIV3 mutant JS cp45. These changes are preferably introduced into recombinant virus using two or three nucleotide changes compared to a corresponding wild type or biologically derived mutant sequence, which has the effect of stabilizing the mutation against genetic reversion.

The present invention also provides recombinant PIV having multiple, phenotype-specifying mutations introduced in selected combinations into the genome or antigenome of an infectious clone. This process, coupled with evaluation of phenotype, provides mutant recombinant PIV having such desired characteristics as attenuation, temperature sensitivity, cold-adaptation, small plaque size, host range restriction, etc. Thus, exemplary PIV clones are disclosed herein which incorporate one or more, and preferably at least two attenuating mutations, e.g., *ts*, *ca* or *att* mutations adopted

from a biologically derived PIV mutant, such as JS cp45. Target genes for adopting biologically derived mutations in a recombinant PIV in this context include the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, fusion protein F and the C, D and V ORF products. Also targeted are extragenic sequences, eg., sequences in the 3' leader or trailer regions of a PIV genome, and in cis-acting elements such as gene start and gene end sequences, eg., the N gene start signal. Exemplary mutations incorporated in recombinant PIV herein include one or more nucleotide substitutions specifying amino acid change(s) in the polymerase L gene, e.g., at Tyr₉₄₂, Leu₉₉₂, and/or Thr₁₅₅₈. For example, PIV recombinants are disclosed wherein Tyr₉₄₂ is replaced by His, Leu₉₉₂ is replaced by Phe, and/or Thr₁₅₅₈ is replaced by Ile. These mutations have been successfully incorporated in various exemplary PIV recombinants herein, including r942, r992, r1558, r942/992, r992/1558, r942/1558, or r942/992/1558 recombinants described in the Examples below. Other exemplary mutations adopted from a biologically derived PIV mutant include one or more mutations in the N protein, including specific mutations at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45. In more detailed aspects, these mutations are represented as Val₉₆ to Ala or Ser₃₈₉ to Ala. Also disclosed in the Examples below are recombinant PIV which encode an amino acid substitution in the C protein, eg., a mutation at a position corresponding to Ile₉₆ of JS cp45, preferably represented by a substitution of Ile₉₆ to Thr. Further exemplary mutations adopted from biologically derived PIV mutants include one or more mutations in the F protein, including mutations adopted from JS cp45 at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀ of JS cp45, preferably represented by acid substitutions Ile₄₂₀ to Val or Ala₄₅₀ to Thr. Other PIV recombinants within the invention adopt one or more amino acid substitutions in the HN protein, as exemplified hereinbelow by a recombinant PIV adopting a mutation at a position corresponding to residue Val₃₈₄ of JS cp45, preferably represented by the substitution Val₃₈₄ to Ala.

Yet additional examples within this aspect of the invention include recombinant PIV which incorporate one or more mutations in an extragenic sequence, eg., a 3' leader sequence of recombinant PIV genome or antigenome. Exemplary mutations in this context include mutations in the 3' leader occurs at one or more positions corresponding to nucleotide 23, 24, and/or 28 of JS cp45, for example a T to C change at nucleotide 23, a C to T change at nucleotide 24, or a G to T change at nucleotide 28. Yet additional extragenic mutations include one or more mutations in a N gene start sequence, as exemplified hereinbelow by a mutation in the N gene start sequence at a position corresponding to nucleotide 62 of JS cp45, preferably represented by a A to T change. The above exemplary mutations adopted from biologically derived mutant PIV are evaluated and combined into recombinant PIV in the Examples below to result, individually and/or in combination, in novel, attenuated candidate vaccine strains, as exemplified by the recombinants designated herein as rcp45, rcp45 3'NCFHN, rcp45 3'NL, rcp45 3'N, and rcp45 F. Other PIV recombinants within the invention will incorporate a plurality and up to a full complement of the mutations present in one or more of these exemplary recombinants, as well as mutations identified in other biologically derived mutant PIV strains identified and adopted in a recombinant PIV according to the teachings herein.

Mutations identified according to the methods disclosed herein are compiled into a "menu" and introduced in various combinations to calibrate a vaccine virus to a selected level of attenuation, immunogenicity and stability. In preferred embodiments, the invention provides for supplementation of one or more mutations adopted from biologically derived PIV, e.g., ts, ca or att mutations, with additional types of mutations involving the same or different genes. Target genes for mutation in this context also include the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, fusion protein F and the C, D and V ORF products. In one aspect, recombinant PIVs are provided wherein at least

one attenuating mutation occurs in the PIV polymerase gene L and involves a nucleotide substitution specifying a *ts* or *att* phenotype adopted from a biologically derived mutant PIV strain, for example JS *cp45*. Exemplary HPIV3 recombinants disclosed herein include the r942, r992, r1558, r942/992, r992/1558, r942/1558, or r942/992/1558 recombinants described in the Examples below. These exemplary PIV clones incorporate one or more nucleotide substitutions resulting in an amino acid change in the polymerase gene, e.g., at Tyr₉₄₂, Leu₉₉₂, and/or Thr₁₅₅₈. For example, PIV recombinants are disclosed wherein Tyr₉₄₂ is replaced by His, Leu₉₉₂ is replaced by Phe, and/or Thr₁₅₅₈ is replaced by Ile. Preferably, two or three mutations are incorporated in a codon specifying an attenuating mutation adding increased stability against phenotypic reversion.

In additional aspects, a variety of other genetic alterations can be produced in a recombinant PIV genome or antigenome for incorporation into infectious recombinant PIV, alone or together with one or more attenuating mutations adopted from a biologically derived mutant PIV. Heterologous genes (e.g. from different PIV strains or non-PIV sources such as another virus, e.g., RSV or measles virus) may be inserted or substituted, in whole or in part, the order of genes changed, gene overlap removed, the PIV genome promoter replaced with its antigenome counterpart, and even entire, non-essential genes deleted. In one aspect, a selected PIV gene, for example the C, D, or V ORF, is functionally deleted to yield a recombinant PIV having novel phenotypic characteristics, for example enhanced growth *in vitro* and/or attenuation *in vivo*. An infectious PIV clone of the invention can also be engineered to enhance its immunogenicity and induce a level of protection greater than that provided by natural infection, or to ablate epitopes associated with undesirable immunopathologic reactions. Enhanced immunogenicity of the vaccines produced by the present invention addresses one of the greatest obstacles to controlling PIV, namely the incomplete nature of immunity induced by natural infection. In this context, additional

gene(s) or gene segment(s) may be inserted into or proximate to the PIV genome or antigenome which may be placed under the control of a common or independent set of transcription signals. Genes of interest include those encoding cytokines (e.g., IL-2 through IL-15, especially IL-2, IL-6 and IL-12, etc.) and proteins rich in T helper cell epitopes. The additional protein can be expressed either as a separate protein or as a chimera engineered from a second copy of one of the PIV proteins, such as HN. This provides the ability to modify and improve the immune response against PIV both quantitatively and qualitatively.

Other mutations useful within the invention involve replacement of the 3' end of genome with its counterpart from antigenome, which is associated with changes in RNA replication and transcription. In addition, intergenic regions can be shortened or lengthened or changed in sequence content. In yet additional aspects, PIV useful in a vaccine formulation can be conveniently modified to accommodate antigenic drift in circulating virus. Typically the modification will be in the HN and/or F proteins. For example, a selected antigenic form of an entire HN or F gene, or the segment(s) encoding particular immunogenic regions thereof, is incorporated into a PIV genome or antigenome cDNA by replacement of a counterpart region in the infectious clone, or by adding one or more copies of the gene such that several antigenic forms are represented in the resultant clone. Progeny virus produced from the modified PIV cDNA are then used in vaccination protocols against emerging strains.

Other mutations for use in infectious PIV of the invention include mutations in cis-acting signals identified during mutational analysis of PIV minigenomes. For example, insertional and deletional analysis of leader and trailer and flanking sequences identify viral promoters and transcription signals and provide a series of mutations associated with varying degrees of reduction of RNA replication or transcription. Saturation mutagenesis (whereby each position in turn is modified to each of the nucleotide alternatives) of these cis-acting signals also identifies mutations which

reduce or increase RNA replication or transcription. Any of these mutations can be inserted into the complete antigenome or genome as described herein.

Additional modifications in PIV clones can be made to facilitate manipulations, such as the insertion of unique restriction sites in various intergenic regions or elsewhere. Nontranslated gene sequences can also be removed to increase capacity for inserting foreign sequences.

Certain substitutions, insertions, deletions or rearrangements of genes or gene segments within recombinant PIV of the invention (e.g., substitutions of a gene segment encoding a selected protein or protein region, for instance a cytoplasmic tail, transmembrane domain or ectodomain, an epitopic site or region, a binding site or region, an active site or region containing an active site, etc.) are made in structural or functional relation to an existing, "counterpart" gene or gene segment from the same or different PIV or other source. Such modifications yield novel recombinants having desired phenotypic changes compared to wild-type or parental PIV or other viral strains. For example, recombinants of this type may express a chimeric protein having a cytoplasmic tail and/or transmembrane domain of one PIV fused to an ectodomain of another PIV. Other exemplary recombinants of this type express duplicate protein regions, such as duplicate immunogenic regions.

As used herein, "counterpart" genes, gene segments, proteins or protein regions, are typically from heterologous sources (e.g., from different PIV genes, or representing the same (i.e., homologous or allelic) gene or gene segment in different PIV types or strains). Typical counterparts selected in this context share gross structural features, e.g., each counterpart may encode a comparable protein or protein structural domain, such as a cytoplasmic domain, transmembrane domain, ectodomain, binding site or region, epitopic site or region, etc. Counterpart domains and their encoding gene segments embrace an assemblage of species having a range of size and sequence variations defined by a common biological activity among the domain or gene segment variants.

For example, two selected protein domains encoded by counterpart gene segments within the invention share substantially the same qualitative activity, such as providing a membrane spanning function, a specific binding activity, an immunological recognition site, etc. More typically, a specific biological activity shared between counterparts, e.g., between selected protein segments or proteins, will be substantially similar quantitatively, i.e., they will not vary in respective quantitative activity levels by more than 30%, preferably by no more than 20%, more preferably by no more than 5-10%.

Counterpart genes and gene segments, as well as other polynucleotides disclosed herein for producing recombinant PIV within the invention, preferably share substantial sequence identity with a selected polynucleotide "reference sequence," e.g., with another selected counterpart sequence. As used herein, a "reference sequence" is a defined sequence used as a basis for sequence comparison, for example, a segment of a full-length cDNA or gene, or a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions wherein a polynucleotide sequence may be compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal

alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith & Waterman, Adv. Appl. Math. 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, J. Mol. Biol. 48:443 (1970), by the search for similarity method of Pearson & Lipman, Proc. Natl. Acad. Sci. USA 85:2444 (1988) (each of which is incorporated by reference), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI, incorporated herein by reference), or by inspection, and the best alignment (i.e., resulting in the highest percentage of sequence similarity over the comparison window) generated by the various methods is selected. The term "sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window

of comparison. The reference sequence may be a subset of a larger sequence.

In addition to these polynucleotide sequence relationships, proteins and protein regions encoded by recombinant PIV of the invention are also typically selected to have conservative relationships, i.e. to have substantial sequence identity or sequence similarity, with selected reference polypeptides. As applied to polypeptides, the term "sequence identity" means peptides share identical amino acids at corresponding positions. The term "sequence similarity" means peptides have identical or similar amino acids (i.e., conservative substitutions) at corresponding positions. The term "substantial sequence identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity or more (e.g., 99 percent sequence identity). The term "substantial similarity" means that two peptide sequences share corresponding percentages of sequence similarity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Abbreviations for the twenty naturally occurring amino acids used herein follow conventional usage

(Immunology - A Synthesis (2nd ed., E.S. Golub & D.R. Gren, eds., Sinauer Associates, Sunderland, MA, 1991), incorporated herein by reference). Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α,α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, ω -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). Moreover, amino acids may be modified by glycosylation, phosphorylation and the like.

The infectious PIV produced from cDNA-expressed genome or antigenome can be any of the PIV or PIV-like strains, e.g., human, bovine, murine, etc. To engender a protective immune response, the PIV strain may be one which is endogenous to the subject being immunized, such as human PIV being used to immunize humans. The genome or antigenome can be modified, however, to express heterologous PIV genes or gene segments, or genes or gene segments from other heterologous sources, e.g., RSV or measles virus. Thus, infectious PIV intended for administration to humans may be human PIV that has been modified to contain genes or gene segments from a bovine or murine PIV type such as for the purpose of attenuation. BPIV3 possesses host range mutations that restrict its replication in rhesus monkeys and humans (Karron et al., supra, 1995a; van Wyke Coelingh et al., 1988), each incorporated herein by reference in its entirety). Gene(s), mutations and cis-acting regulatory sequences of BPIV3 that specify desired phenotypes, e.g., host range restriction, will be identified by their substitution for corresponding sequences in rPIV of the invention, and incorporated within further modified rPIV to develop yet additional useful vaccine agents. Similarly, mutations of JS cp45 which are known to impart non-ts host-range attenuating mutations for the rhesus monkey (Hall et al., supra, (1992))

will likewise be identified and incorporated into modified rPIV vaccine agents of the invention. Alternatively, a bovine PIV may be modified to contain genes that encode, e.g., proteins or immunogenic epitopes that elicit protection against human PIV infection. For example, human PIV glycoprotein genes can be substituted for counterpart bovine glycoprotein genes, such that the bovine PIV elicits a protective immune response in humans against human PIV strains.

10 In exemplary embodiments, individual genes, gene segments, or single or multiple nucleotides of one PIV are substituted by counterpart sequence(s) from a heterologous PIV or other source. For example, heterologous gene segments, such as one encoding a cytoplasmic tail, transmembrane domain or ectodomain, an epitopic site or region, a binding site or region, an active site or region containing an active site, etc., of a selected protein from one PIV is substituted for a counterpart gene segment in another PIV to yield novel recombinants, for example recombinants expressing a chimeric protein having a cytoplasmic tail and/or transmembrane domain of one PIV fused to an ectodomain of another PIV. Preferred genome segments in this regard range from about 15-35 nucleotides in the case of gene segments encoding small functional domains of proteins, e.g., epitopic sites, to about 25 50, 75, 100, 200-500, or 500-1,500 or more nucleotides for gene segments encoding larger domains or protein regions.

In one aspect of the invention, selected domains of HN and/or F proteins of one PIV strain are substituted into a heterologous PIV clone to produce a recombinant virus capable of stimulating a cross-protective immune response against both PIV strains in an immunized host. In other aspects, modified PIV clones are provided which comprise a chimera of a human PIV genomic or antigenomic sequence and at least one non-human PIV sequence, for example a polynucleotide containing sequences from both human and bovine PIV. The replacement of a human PIV coding sequence or non-coding sequence (e.g., a promoter, gene-end, gene-start, intergenic or other cis-acting element) with a counterpart bovine or murine PIV sequence

yields recombinants having a variety of possible attenuating effects. For example, a host range effect will often arise from a heterologous PIV gene not functioning efficiently in a human cell, from incompatibility of the heterologous sequence or protein with a biologically interactive human PIV sequence or protein (e.g., a sequence or protein that ordinarily cooperates with the substituted sequence or protein for viral transcription, translation, assembly, etc.), among other useful attenuating effects. In yet another aspect of the invention, insertion of foreign genes or gene segments, and in some cases of noncoding nucleotide sequences, into the PIV genome results in a desired increase in genome length causing yet additional, desired phenotypic effects. Increased genome length is expected to result in attenuation of the resultant PIV clone, dependent in part upon the length of the insert. In addition, the expression of certain proteins from a gene inserted into recombinant PIV will result in attenuation of the virus due to the action of the protein. This has been described for IL-2 expressed in vaccinia virus (see, e.g., Flexner et al., Nature 33:-259-62 (1987)) and also would be expected for gamma interferon.

Deletions, insertions, substitutions and other mutations involving changes of whole viral genes or gene segments in recombinant PIV of the invention yield highly stable vaccine candidates, which are particularly important in the case of immunosuppressed individuals. Many of these mutations will result in attenuation of resultant vaccine strains, whereas others will specify different types of desired phenotypic changes. For example, certain viral genes are known which encode proteins that specifically interfere with host immunity (see, e.g., Kato et al., EMBO. J. 16:578-87 (1997), incorporated herein by reference). Ablation of such genes in vaccine viruses is expected to reduce virulence and pathogenesis and/or improve immunogenicity.

In preferred aspects of the invention, the modified PIV clones represent a chimera of two or more human PIV genomes, for example a clone containing polynucleotide sequences from HPIV3 joined to sequences from one or more

heterologous human PIV, such as HPIV1 and HPIV2. Thus, individual genes or gene segments of human PIV3 may be replaced or supplemented with counterpart genes or gene segments from HPIV 1 or HPIV2, or visa versa. In one example
5 described hereinbelow, the invention provides a PIV clone, rPIV3-1, into which both the HN and F glycoprotein genes of HPIV1 are substituted for their counterpart genes in an HPIV3 background, yielding a chimeric virus having immunological characteristics representative of both parental strains.

10 In additional aspects of the invention, chimeric PIV or PIV clones having other alterations of genes or gene segments, as described above, are further modified by introducing one or more attenuating mutations adopted from a biologically derived mutant PIV, e.g., HPIV3 JS cp45 to
15 achieve an attenuated, or further attenuated, chimeric mutant derivative. For example, one or more human PIV coding or non-coding polynucleotides may be substituted with a counterpart sequence from a heterologous human PIV, bovine PIV or murine PIV as described above, and this alteration may be combined
20 with one or more mutations specifying, e.g., a *ts*, *ca* or *att* phenotype adopted from a biologically derived attenuated PIV mutant, to yield an attenuated or further attenuated (i.e., compared to either the chimeric clone or biologically derived parent virus) vaccine virus. Alternatively, functional
25 deletion of a non-essential gene or gene segment, such as the C, D or V ORF, may be combined in a recombinant PIV with one or more mutations specifying a *ts*, *ca* or *att* phenotype from biologically derived PIV mutants to yield an attenuated vaccine strain. These combinatorial modifications yield
30 recombinant PIV having desired phenotypic characteristics, e.g., increased yield of virus, enhanced attenuation, and/or genetic resistance to reversion from an attenuated phenotype, due to the combined effects of the different selected mutations.

35 In one combinatorial mutation design, a modified PIV is provided which comprises a chimera of a human PIV genomic or antigenomic sequence and at least one non-human PIV sequence, for example a polynucleotide containing sequences

from both human and bovine PIV, and which also incorporates one or more mutations adopted from biologically derived PIV, e.g., one or more naturally occurring *ts*, *ca* or *att* mutations. Alternatively, the modified PIV can be a chimera of two or more human PIV genomes, for example a polynucleotide containing sequences from HPIV3 joined to sequences from one or more heterologous human PIVs, such as HPIV1 and HPIV2, which further incorporates one or more *ts*, *ca* *att* or other selected mutations from biologically derived PIV (e.g., a nucleotide substitution specifying a *ts*, *ca* or *att* phenotype adopted from a biologically derived mutant PIV strain such as JS cp45). In more detailed aspects, individual genes or gene segments of human PIV3 are replaced or supplemented with counterpart genes or gene segments from HPIV1 or HPIV2, or visa versa, in a clone that is attenuated or further attenuated by, e.g., a nucleotide change encoding an amino acid substitution conferring a *ts* mutation in the large polymerase L gene. For example, the invention provides PIV clones having the HN and/or F glycoprotein genes of HPIV1 substituted for their counterpart genes in an HPIV3 background, wherein the phenotype of the resultant chimeric clone is further modified by *ts*, *ca* or *att* mutation(s) encoded within one or more of the N, P, L, M, HN, F, C, D and V genes. Various combinations from a menu of possible mutations disclosed herein can be made to calibrate a vaccine virus to a selected level of attenuation, immunogenicity and stability, e.g., to achieve a satisfactorily attenuated and immunogenic, chimeric virus having immunological characteristics representative of multiple PIV strains. In one aspect, recombinant PIVs are provided wherein at least one attenuating mutation occurs in the PIV polymerase gene L (as exemplified by the recombinants r942, r992, r1558, r942/992, r942/1558, r992/1558, or r942/992/1558 described in the Examples below) incorporated in a chimeric PIV background. For example, useful chimeric PIV recombinants within this aspect of the invention will have one or more genes or gene segments of the HN and/or F glycoprotein genes from, e.g., HPIV1 substituted for their counterpart gene(s) in a heterologous background,

e.g., in an HPIV3 clone, and will further incorporate one or more attenuating mutations, eg., nucleotide substitutions resulting in an amino acid change in the polymerase gene (such as change from Tyr to His at position 942, a change from Leu to Phe at position 992, and/or a change from Thr to Ile at position 1558) from a biologically derived PIV mutant. One such chimeric, attenuated recombinant exemplified hereinbelow is rPIV3-1.cp45L, a derivative of rPIV3-1 which incorporates all three L gene mutations specified above from JS cp45.

Yet additional mutations which can be incorporated in a chimeric PIV background for developing vaccine strains will be selected from biologically derived mutations in other genes, or will be created *de novo* in a recombinant genome by standard site directed mutagenesis or other purely recombinant mutagenic methods. Target genes for adopting biologically derived mutations or creating novel mutations in a recombinant PIV in this context include the nucleocapsid protein N, phosphoprotein P, large polymerase subunit L, matrix protein M, hemagglutinin-neuraminidase protein HN, fusion protein F and the C, D and V ORF products. Also targeted are extragenic sequences, eg., sequences in the 3' leader or trailer regions of a PIV genome. Exemplary mutations identified and incorporated in non-chimeric, recombinant PIV, described above will thus be readily incorporated within a chimeric PIV background, eg., as exemplified by rPIV3-1. These exemplary mutations include one or more mutations in the N protein, including specific mutations at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45. In more detailed aspects, these mutations are represented as Val₉₆ to Ala or Ser₃₈₉ to Ala. Also desired for incorporation in chimeric PIV recombinants are mutations in the C protein, eg., a mutation at a position corresponding to Ile₉₆ of JS cp45, preferably represented by a substitution of Ile₉₆ to Thr, as described above. Further exemplary mutations for incorporation in a chimeric PIV background include one or more mutations in the F protein, for example adopted from JS cp45 at a position corresponding to residues Ile₄₂₀ or Ala₄₅₀, eg., substitutions Ile₄₂₀ to Val or Ala₄₅₀ to Thr. Yet additional chimeric PIV

recombinants within the invention will adopt one or more amino acid substitutions in the HN protein, for example a mutation at a position corresponding to residue Val₃₈₄ of JS cp45, such as Val₃₈₄ to Ala. Yet additional chimeric recombinants will incorporate one or more mutations in an extragenic sequence, eg., a 3' leader sequence of the recombinant genome or antigenome. Exemplary mutations in this context include mutations in the 3' leader occurs at one or more positions corresponding to nucleotide 23, 24, 28, and/or 45 of JS cp45, for example a T to C change at nucleotide 23, a C to T change at nucleotide 24, a G to T change at nucleotide 28, or a T to A change at nucleotide 45. Yet additional extragenic mutations for incorporation within a chimeric PIV background include one or more mutations in a N gene start sequence, as exemplified herein by a mutation in the N gene start sequence at a position corresponding to nucleotide 62 of JS cp45, such as a A to T change. These exemplary mutations evaluated and combined into recombinant PIV in the Examples below will be readily incorporated within a chimeric PIV recombinant using the methods and tools provided herein, and will specify, individually and/or in combination, desired phenotypic changes to yield yet additional attenuated chimeric vaccine strains within the invention.

In additional combinatorial mutation designs, modified PIVs are provided which incorporate one or more of the foregoing *ts*, *ca* or *att* mutations adopted from biologically derived PIV or generated recombinantly in a PIV clone of the invention, in combination with another, distinct mutation disclosed herein (e.g., a deletion, addition, or rearrangement of a PIV N, P, L, M, HN, F, C, D or V gene or gene segment, or a gene or gene segment from another source such as RSV or measles virus). Also in this case, various combinations from a menu of mutations disclosed herein can be made to calibrate the vaccine virus to a selected level of attenuation, immunogenicity and stability. Thus, recombinant PIVs are provided which exhibit at least one attenuating mutation from a biologically derived PIV mutant, e.g., a mutation in the PIV polymerase gene L as found in JS cp45, or a recombinantly

generated mutation, and which further incorporates one or more additional changes selected from, e.g., substitution or introduction of a heterologous gene or gene segment from a non-PIV source (e.g., an immunogenic RSV or measles gene or epitope, or a gene encoding a cytokine), a change in the order of viral genes to alter expression levels, removal of gene overlap, substitution of a PIV genome promoter with its antigenome counterpart, shortening, lengthening or removal of intergenic regions, e.g., to increase capacity for inserting foreign sequences, mutations in cis-acting signals to reduce or increase RNA replication or transcription, insertion of unique restriction sites, or deletion of even entire, non-essential genes, among other changes.

For vaccine use, virus produced according to the present invention can be used directly in vaccine formulations, or lyophilized, as desired, using lyophilization protocols well known to the artisan. Lyophilized virus will typically be maintained at about 4°C. When ready for use the lyophilized virus is reconstituted in a stabilizing solution, e.g., saline or comprising SPG, Mg⁺⁺ and HEPES, with or without adjuvant, as further described below.

PIV vaccines of the invention contain as an active ingredient an immunogenically effective amount of PIV produced as described herein. The modified virus may be introduced into a host with a physiologically acceptable carrier and/or adjuvant. Useful carriers are well known in the art, and include, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration, as mentioned above. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, and the like. Acceptable adjuvants include incomplete Freund's

adjuvant, MPL™ (3-o-deacylated monophosphoryl lipid A; RIBI ImmunoChem Research, Inc., Hamilton, MT) and IL-12 (Genetics Institute, Cambridge MA), among many other suitable adjuvants well known in the art.

5 Upon immunization with a PIV composition as described herein, via aerosol, droplet, oral, topical or other route, the immune system of the host responds to the vaccine by producing antibodies specific for PIV virus proteins, e.g., F and HN glycoproteins. As a result of the vaccination with
10 an immunogenically effective amount of PIV produced as described herein, the host becomes at least partially or completely immune to PIV infection, or resistant to developing moderate or severe PIV infection, particularly of the lower respiratory tract.

15 The host to which the vaccines are administered can be any mammal which is susceptible to infection by PIV or a closely related virus and which host is capable of generating a protective immune response to the antigens of the vaccinizing strain. Accordingly, the invention provides
20 methods for creating vaccines for a variety of human and veterinary uses.

 The vaccine compositions containing the PIV of the invention are administered to a host susceptible to or otherwise at risk for PIV infection to enhance the host's own
25 immune response capabilities. Such an amount is defined to be a "immunogenically effective dose." In this use, the precise amount of PIV to be administered within an effective dose will depend on the host's state of health and weight, the mode of administration, the nature of the formulation, etc., but will
30 generally range from about 10^3 to about 10^6 plaque forming units (PFU) or more of virus per host, more commonly from about 10^4 to 10^5 PFU virus per host. In any event, the vaccine formulations should provide a quantity of modified PIV of the invention sufficient to effectively protect the host patient
35 against serious or life-threatening PIV infection.

 The PIV produced in accordance with the present invention can be combined with viruses of other PIV serotypes or strains to achieve protection against multiple PIV

serotypes or strains. Alternatively, protection against multiple PIV serotypes or strains can be achieved by combining protective epitopes of multiple serotypes or strains engineered into one virus, as described herein. Typically
5 when different viruses are administered they will be in admixture and administered simultaneously, but they may also be administered separately. Immunization with one strain may protect against different strains of the same or different serotype.

10 In some instances it may be desirable to combine the PIV vaccines of the invention with vaccines which induce protective responses to other agents, particularly other childhood viruses. In another aspect of the invention the PIV can be employed as a vector for protective antigens of other
15 pathogens, such as respiratory syncytial virus (RSV) or measles virus, by incorporating the sequences encoding those protective antigens into the PIV genome or antigenome which is used to produce infectious PIV, as described herein. The cloning of RSV cDNA and other disclosure relevant to the
20 invention is described in copending United States patent applications Serial Nos. 08/534,768, 60/021,773, 08/720,132, 60/046,141, 60/047,634, and 08/892,403, and PCT patent application PCT/US97/12269, each incorporated herein by reference.

25 Single or multiple administrations of the vaccine compositions of the invention can be carried out. In neonates and infants, multiple administration may be required to elicit sufficient levels of immunity. Administration should begin within the first month of life, and at intervals throughout
30 childhood, such as at two months, six months, one year and two years, as necessary to maintain sufficient levels of protection against native (wild-type) PIV and/or other subject viral infections. Similarly, adults who are particularly
35 susceptible to repeated or serious PIV infection, for example health care workers, day care workers, family members of young children, the elderly, and individuals with compromised cardiopulmonary function, may require multiple immunizations to establish and/or maintain protective immune responses.

Levels of induced immunity provided by the vaccines of the invention can be monitored by measuring amounts of neutralizing secretory and serum antibodies. Based on these measurements, vaccine dosages can be adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine viruses may be advantageous for different recipient groups. For example, an engineered PIV strain expressing an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

In yet another aspect of the invention the PIV is employed as a vector for transient gene therapy of the respiratory tract. According to this embodiment the recombinant PIV genome or antigenome incorporates a sequence which is capable of encoding a gene product of interest. The gene product of interest is under control of the same or a different promoter from that which controls PIV expression. The infectious PIV produced by coexpressing the recombinant PIV genome or antigenome with the N, P, L and other desired PIV proteins, and containing a sequence encoding the gene product of interest, is administered to a patient. Administration is typically by aerosol, nebulizer, or other topical application to the respiratory tract of the patient being treated. Recombinant PIV is administered in an amount sufficient to result in the expression of therapeutic or prophylactic levels of the desired gene product. Representative gene products which may be administered within this method are preferably suitable for transient expression, including, for example, interleukin-2, interleukin-4, gamma-interferon, GM-CSF, G-CSF, erythropoietin, and other cytokines, glucocerebrosidase, phenylalanine hydroxylase, cystic fibrosis transmembrane conductance regulator (CFTR), hypoxanthine-guanine phosphoribosyl transferase, cytotoxins, tumor suppressor genes, antisense RNAs, and vaccine antigens.

The following examples are provided by way of illustration, not limitation.

EXAMPLE IConstruction of Plasmid p218(131)
Encoding Negative sense PIV Genomic RNA

5 A full cDNA clone designated p218(131) (Fig. 1; SEQ
ID NO: 1) (deposited under the terms of the Budapest Treaty
with the American Type Culture Collection (ATCC) of 12301
Parklawn Drive, Rockville, Maryland 20852, U.S.A., and granted
the designation 97991) was constructed to encode the complete
10 15462 nt genomic sequence of HPIV3 JS strain. A hepatitis
delta ribozyme was placed abutting the 3' end of the genomic
sequence such that self-cleavage would yield the 3' end of
HPIV3 (Perrotta and Been, Nature 350: 434-436, (1991),
incorporated herein by reference in its entirety). A T7
15 transcription terminator was placed following the delta
ribozyme. The T7 promoter was placed adjacent to the 5' end
of the genomic sequence such that the 5' terminal nucleotide
of the HPIV3 genome was the first nucleotide synthesized. In
this configuration, the cDNA encodes a complete negative-sense
20 copy of PIV3 genomic RNA containing the correct genomic
termini without any additional heterologous nucleotides.

 The HPIV3 cDNA was assembled from 14 overlapping
subclones (termed A*-L, which letters in parentheses designate
individual plasmids and do not refer to specific viral genes)
25 constructed by reverse transcription (RT) and polymerase chain
reaction (PCR) of RNA isolated from virions purified by
sucrose gradient centrifugation (Stokes et al., supra, 1992;
Stokes et al., supra, 1993, each incorporated herein by
reference in its entirety). The subclones spanned the
30 following nucleotides of genomic RNA (numbered with the 3' end
designated as position 1): 1-2058 (A*), 1874-3111 (A'),
3086-5140 (C), 4348-5276 (C'), 5072-6695 (D*), 5904-8532 (E),
7806-9898 (F), 9632-10740 (F'), 9760-10955 (G), 10862-11925
(H), 11835-12868 (I), 12426-13677 (J), 13630-14496 (K), and
35 14467-15462 (L). Each fragment was cloned into pBluescript
KSII (Stratagene, La Jolla, CA) using conventional cloning
techniques and was sequenced completely.

Plasmid p(L) was then subjected to site-directed mutagenesis to introduce the T7 promoter via a single-stranded DNA intermediate according to the MUTA-GENE procedure (BioRad, Hercules, CA). The T7 promoter was positioned so that transcription initiates at the precise 5' end of the HPIV3 genome using the negative-sense mutagenic primer:

5'-AATACGACTCACTATA*ACCAAACAAGAGAAG-3' (SEQ ID NO: 2; T7

sequences are italicized, HPIV3-specific sequences are underlined, and the 5'-end HPIV3 nucleotide, genome position 15462, is indicated by an asterisk). This modified p(L) was designated p(L*). Plasmid p(E) was modified to yield p(E*) by the same method using the negative-sense mutagenic

oligonucleotide 5'-CCAAGTACTATGAGATGCTTGATT-3' (SEQ ID NO: 3) to insert three nucleotide substitutions (underlined) into the HN gene at HPIV3 position 7903, 7913, 7915 (Fig. 1). These substitutions removed an *Hga* I site, inserted a *Sca* I site, and modified amino acid 370 of the encoded HN protein such that the epitope recognized by monoclonal antibodies (mAb) 423/6 and 170/7 was ablated (van Wyke Coelingh et al., J. Virol. 61:1473-1477, (1987), incorporated herein by

reference). The p(E*) to p(K) subclones were assembled in a step-wise fashion into the p(L*) plasmid to give p(E*FF'GHIJKL*) (Fig. 1A). This plasmid includes HPIV3 nucleotides 5904-15462 with the T7 promoter adjacent to nucleotide 15462 at the 5' end of the genome. Subclones p(A*) to p(E) were assembled into a second, overlapping subclone, p(A*A'CC'D'E) which contained HPIV3 nucleotides 1-8532.

Both subclones p(E*FF'GHIJKL*) and p(A*A'CC'D'E) were sequenced completely. In addition to the introduced point mutations described above, the cDNA differed from the authentic JS HPIV3 sequence (Stokes et al., supra, 1992) by a single nucleotide substitution at position 1615 which was within the N gene and caused a substitution at amino acid 506 in the encoded protein. Three other nucleotide substitutions were found at positions 10355, 11333, and 15248 in the L gene which did not change the encoded protein (Fig. 2). These three noncoding changes were retained as additional sequence markers to identify recombinant virus (designated rPIV)

derived from cDNA, and the mutation in the N gene was corrected as described later.

Subclone p(A*A'CC'D'E) was then modified to insert the hepatitis delta virus ribozyme and T7 terminator adjacent to HPIV3 position 1. An HPIV3 minigenome in which the 3' end of the HPIV3 genome (GGT↓GGG) (underlined) was generated through self-cleavage of a flanking hepatitis delta virus antigenomic ribozyme (shown in part in bold-type) was previously constructed (Dimock and Collins, J. Virol. 67: 2772-2778, (1993); Perrotta and Been, supra, (1991), each incorporated herein by reference in its entirety). The ribozyme in turn was followed by a T7 transcription terminator. This minigenome cDNA was used as a template in a PCR reaction which modified the sequence adjacent to the ribozyme cleavage site to be a *Sma* I site (CCC↓GGG) and placed an *Apa*I site (GGGCC↓C) on the downstream side of the T7 terminator. The PCR product was cloned into pKSII which had been digested with *Bss*HII and made blunt-ended by filling in, yielding p218.

p218 was designed such that any sequence could be introduced into the opened *Sma* I site by blunt-end ligation and its RNA transcript would be cleaved at the delta ribozyme cut site (NNN↓GGG). The p(A*A'CC'D'E) subclone was digested with *Hga* I and *Sal* I (8533), which released the HPIV3 cDNA, and was filled in with dNTPs and T4 DNA polymerase to give blunt termini. The *Hga* I site is 10 nucleotides upstream of HPIV3 position 1 and, when digested and filled in, leaves a blunt terminus beginning with HPIV3 position 1. The modified *Hga* I-*Sal* I fragment was gel purified and cloned into the *Sma* I site of p218. The mutation in the N gene (T at nt 1615) was corrected to the JS wt sequence (A at nt 1615) (see GenBank accession #Z11575, incorporated herein by reference) using Kunkel mutagenesis (Kunkel et al., Methods Enzymol. 154: 367-382, (1987), incorporated herein by reference in its entirety). This plasmid was designated p218(A*A'CC'D'E) (Fig. 1).

The *Xho* I-*Ngo* MI fragment of p(E*FF`GHIJKL*), which contained the T7 promoter and the HPIV3 cDNA from nucleotides 7438-15462, was cloned into the *Xho* I-*Ngo* MI window of p218(A*A`CC`D*E) (Fig. 1). This joined the two fragments of HPIV3 cDNA at HPIV3 nucleotide 7438, yielding a plasmid containing a full-length HPIV3 cDNA encoding a negative-sense genomic RNA with the above-mentioned three desired mutations in the HN gene and three incidental mutations in the L gene. The final construct, designated p218(131) (Fig. 1; SEQ ID NO: 1), was then sequenced in its entirety by automated sequencing at the NCI Frederick Cancer Research and Development Center (Frederick, MD) using the Taq DYE Deoxy Terminator cycle sequencing kit (ABI, Foster City, CA). This identified an additional change in the HN gene, namely a change of C to T in the HN gene at position 7593 which changed HN amino acid 263 from threonine to isoleucine that is also indicated in Fig. 2.

EXAMPLE II

Transcription and RNA Replication System for HPIV3

The present example describes compositions and methods for producing a reconstituted transcription and RNA replication system for human parainfluenza virus type 3 (HPIV3). This exemplary system was developed using components expressed intracellularly from transfected plasmids driven by a T7 RNA polymerase supplied by a vaccinia virus recombinant. The system is based on a negative-sense analog of HPIV3 genomic RNA in which the viral genes were deleted and replaced with a polynucleotide encoding bacterial chloramphenicol acetyl transferase (CAT). The N, P and L proteins are expressed from cotransfected plasmids so as to direct efficient transcription and RNA replication. Transcription according to this example yields subgenomic polyadenylated mRNA, which can be readily isolated, e.g., by oligo(dT) chromatography. RNA replication according to this example yields mini-antigenome and progeny minigenome, which are shown

to be encapsidated based on resistance to digestion with micrococcal nuclease.

A) Viruses and cells

A vaccinia virus recombinant, vTF7-3, that expresses bacteriophage T7 RNA polymerase, was provided as described by Fuerst et al. (Proc. Natl. Acad. Sci. U.S.A. **83**: 8122-8126, 1986, incorporated herein by reference in its entirety). HEP-2 monolayer cultures were maintained at 37°C in 5% CO₂ with OptiMEM 1 (Life Technologies, Gaithersburg, MD) supplemented with 2% fetal bovine serum (FBS), 50 µg/ml gentamicin sulfate and 2mM glutamine.

B) cDNAs

cDNAs corresponding to ORFs of the N, P, and L genes of the JS strain of HPIV3 (GenBank #Z11575; Stokes et al., 1992) were individually cloned into the Nco I-Sal I window of plasmid pTM-1, in which transcription is mediated by T7 RNA polymerase and translation by an internal ribosome entry site preceding the foreign ORF (Elroy-Stein et al., Proc. Natl. Acad. Sci. U.S.A. **86**: 6126-6130 (1989), incorporated herein by reference in its entirety). Each gene was first modified by polymerase chain reaction (PCR) to place an Nco I or Nco I-compatible site at the translational start site and a Sal I site on the downstream end.

The plasmid p(131), which is similar to p218(131) except that it lacks the hepatitis delta virus ribozyme, was used as a template for each PCR. The primers used to amplify the N ORF were CCCTATAATTTCAAC**ATG**TTGAGCCTATTTG (SEQ ID NO: 4; forward primer relative to positive-sense) and GATTAAAATGTTGGT**CGACTTAGTTGCTTCC** (SEQ ID NO: 5; italics represent restriction enzyme sites, and the translational start site is in bold). The PCR product, a 1578 bp fragment flanked by an Afl III and Sal I site, was cloned into the Nco I-Sal I window of pTM-1 to yield pTM(N).

The primers used to amplify the PIV3 phosphoprotein (P) ORF were 5'-CCATAGAGAGTCC**ATG**GAAAGCGATGCTAAAACTATC-3' (SEQ ID NO:6; forward primer) and 5'-CGGTGTCGTTTCTTTGTCGACTCATTGGCAATTGTTG-3' (SEQ ID NO:7; reverse primer). A full-length cDNA of JS strain of genomic RNA

(p131) was used as template for the PCR. The resultant PCR product was an 1851 bp fragment flanked by an *Nco* I and *Sal* I restriction site (in italics). The PCR product was then cloned into the *Nco* I-*Sal* I window of pTM-1 to yield pTM(P).

5 A second PCR was performed to amplify the PIV3 phosphoprotein P ORF without the C ORF. p131 was again used as template cDNA. A different forward primer and the same reverse primer were used to amplify the PIV3 P ORF without C; 5'-CCATAGAGAGTCCATGGAAAGCGACGCTAAAACTATC-3' (SEQ ID NO: 74; forward primer) and 5'-CGGTGTCGTTTCTTTGTGCGACTCATTGGCAATTGTTG-3' (SEQ ID NO:7; reverse primer). The resultant PCR product was an 1851 bp fragment flanked by an *Nco* I and *Sal* I restriction site (designated by italics). The underlined nucleotide in the forward primer represents a nucleotide substitution which is silent in the P ORF but changes the start codon of the C ORF to threonine. The next start codon for the C ORF is more than 400 nucleotides downstream. Thus, only the P protein would be produced. The PCR product was then cloned into the *Nco* I-*Sal* I window of pTM-1 to yield a second plasmid, pTM(P no C).

The L ORF of HPIV3 was cloned into pTM-1 in three parts: the ends were derived from PCR products and the main body was a restriction fragment from p218(131). The upstream end of the L ORF was amplified using the primers 25 GCAAAGCGTGCCCGGGCCATGGACACTGAATCTAACAATGGC (SEQ ID NO: 8) and GAAATTCCTTAATCGATTCTCTAGATTC (SEQ ID NO: 9). This yielded the 1,020-bp PCR product L1 in which positions 8625-9645 of the full-length genome were flanked by *Sma* I and *Nco* I sites on the upstream end and a *Cla* I site on the downstream end (all three sites are italicized). The downstream end of the L ORF was amplified using the primers CCCATCAACTGTAACATACGTAAGAAAGAC (SEQ ID NO: 10) and GGTTAGGATATGTGCGACATTGTATTTATG (SEQ ID NO: 11). This yielded the 1,733-bp PCR product L2 in which positions 13,645-15,378 of the full-length genome were flanked by a *SnaB* I and *Sal* I site (italicized). Plasmid p(131) was digested with *Cla* I and *Pst* I to yield the 4,487-bp fragment L middle containing positions 9,630-14,120 of the full-length genome. L1 and L middle were joined at the common *Cla* I site

and cloned into the *Sma* I-*Pst* I window of pBluescript to yield p(L1+L middle). The L2 fragment was then cloned into the *Pst* I-*Sal* I window of p(L1+L middle) to yield the complete L ORF flanked by *Nco* I and *Sal* I. This was then cloned into the *Nco* I-*Sal* I window of pTM-1 to yield pTM(L). The sequences of PCR-generated regions of pTM(N) (SEQ ID NO: 20), pTM(P) (SEQ ID NO: 21), and pTM(L) (SEQ ID NO: 22) were confirmed by the dideoxynucleotide sequencing method.

To increase the efficiency of T7 transcription, certain modifications were made to a cDNA construct encoding a negative-sense PIV minigenome, called PIV3-CAT(-) (Dimock and Collins, *J. Virol.* 67: 2772-2778 (1993), incorporated herein by reference in its entirety). PIV3-CAT(-) includes the 3'-terminal 111 nucleotides and 5'-terminal 115 nucleotides of the HPIV3 genome fused to a negative-sense copy of the CAT ORF. This cDNA was designed to yield, upon linearization with *Hga*I and transcription with T7 RNA polymerase, a minigenome containing the exact correct ends of the HPIV3 genome. Two successive rounds of PCR, using mutagenic oligonucleotides which added successive extensions to the cDNA end, were used to replace the *Hga*I site with the hepatitis delta ribozyme (Perotta and Been, *Nature* 350: 434-436 (1991), incorporated herein by reference in its entirety), such that self-cleavage generates the correct 3' HPIV3 genomic end. A T7 transcriptional termination signal was inserted immediately after the ribozyme (Fig. 3) to yield PIV3-CAT-delta.

PIV3-CAT-delta cDNA was modified by PCR mutagenesis to insert one, two or three G residues between the T7 promoter and the 5' end of the minigenome, using restriction sites flanking the trailer and T7 promoter. This modification yielded increased efficiency of T7 transcription. In preliminary experiments, the minigenome containing two G residues, called PIV3-CAT-GG, was the most active in the expression of CAT in the reconstituted transcription and replication system described below, and was used for all subsequent derivatives.

The PIV3-CAT-GG cDNA was further modified by overlapping PCR mutagenesis to introduce modifications

simultaneously at two sites, as follows. First, the sequence T₇CT, which contains tandem transcriptional termination motifs for the vaccinia virus early-stage RNA polymerase (T₅NT) (Yuen and Moss, Proc. Natl. Acad. Sci. U.S.A. 84: 6417-6421 (1987), incorporated herein by reference in its entirety), was inserted into the positive-sense strand cDNA strand between the delta ribozyme and T7 transcriptional terminator (Fig. 1). This motif, and a second motif described herein below, were added to prevent promiscuous transcription of the CAT gene by the vaccinia virus early RNA polymerase. Second, the minigenome cDNA insert was modified at the junction between the N nontranslated region and CAT gene to contain (i) the insertion of a second vaccinia virus termination motif (T₅AT) into the positive-sense cDNA strand at positions 103 to 109 relative to the encoded minigenome, and (ii) the insertion of 0 to 6 G residues (negative-sense) at minigenome position 112 (Fig. 3). The overlapping PCR mutagenesis involved three sets of reactions (PCR 1, 2 and 3) performed as follows. PCR 1 was a set of seven parallel reactions [PCR1 (0) to (+6)] which used the PIV3-CAT-GG cDNA as a template and the following two mutagenic oligonucleotides as primers: the forward primer was: ⁻¹¹¹GGGGTTATGCTACTGCAGGCTTTTTTCTCCCTTAGCCATCCG⁻⁶² (SEQ ID NO: 12) and the reverse primer was: ¹²⁴CTCCATTCTAGA(N)TTATAAAAATTATAGAGTCCCC⁹⁰ (SEQ ID NO: 13). The bold sequence in the first oligonucleotide is the upstream tandem vaccinia terminator, and the bold sequence in the second oligonucleotide is the second terminator. This reaction amplified the ribozyme and adjacent leader region and inserted the mutations described above. PCR 2 was a single reaction that used the PIV3-CAT-GG cDNA as a template, and a forward primer that hybridized in plasmid sequence upstream of a unique NgoMI site (Fig. 3), and a reverse primer complementary to the forward primer of reaction one. Thus, the products of PCR 1 and 2 overlapped at this latter sequence. The products of PCR 1 (0) to (+6) and PCR 2 were gel purified. The products of PCR1 (0) to (+6) were each mixed separately with an aliquot of PCR 2 product and amplified in a third reaction (PCR3 (0) to (+6)) which also

contained the forward primer of PCR 2 and the reverse primer of PCR 1. The products of PCR3 (0) to (+6) were digested with *Ngo*MI, which cuts in plasmid-specific sequence, and *Xba*I, which cuts at the upstream end of the CAT gene (Fig. 3), and cloned into the *Ngo*MI-*Xba*I window of PIV3-CAT-GG. This resulted in a panel of cDNAs encoding minigenomes which were named according to the number of inserted G residues: PIV3-CAT 0 to PIV3-CAT +6. The structures of all DNA regions derived from PCR were confirmed by dideoxynucleotide sequencing.

10 C) Transfection

HEp-2 cells were grown to 90% confluence in 6 well plates. Each well of a six-well plate (1.5×10^6 cells) was transfected with 0.4 μ g pTM(P), 0.4 μ g pTM(N), 0.05 μ g pTM(L), and 0.4 μ g minigenome plasmid. The plasmids were added to 0.1 ml of OptiMEM (Life Technologies) and mixed with 0.1 ml of OptiMEM containing 12 μ l of LipofectACE (Life Technologies). After an incubation period of approximately 15 minutes at room temperature, 0.8 ml of OptiMEM 1 containing 2% calf serum and 1.5×10^7 pfu of vTF7-3 was added to each well. The plates were incubated at 37°C for 12 hours after which the media was replaced with fresh OptiMEM 1 containing 2% fetal bovine serum. The cells were then incubated at 37°C for a total of 48 hours and harvested for RNA analysis and CAT assay. Each minigenome was represented in triplicate (3 wells) which was scraped into the medium and pooled.

25 D) CAT assay

An aliquot representing 3.33% (1.5×10^5 cells) of each pooled sample of harvested cells described above was removed for CAT assay. The aliquot was centrifuged at 1,000 rpm for 5 minutes and the supernatant discarded. The cell suspension was washed with 1 ml of 40 mM Tris, pH 7.5, 1 mM EDTA, 150 mM NaCl and resuspended in 50 μ l 0.25 M Tris, pH 7.5. Lysate was prepared by three cycles of freezing and thawing and clarified by centrifuging at 8,000 rpm for 5 minutes. 1 μ l of lysate was assayed for the ability to acetylate D-threo-[dichloroacetyl 1- 14 C]chloramphenicol (Amersham) using a conventional assay (Gorman et al., Mol. Cell. Biol. 2: 1044-1051 (1982), incorporated herein by

reference in its entirety). Acetylation was visualized by thin-layer chromatography and quantified by phosphoimager analysis (Molecular Dynamics, Sunnyvale, CA).

E) RNA analysis

5 The remaining cell harvest of each pooled sample was divided into three equal parts for isolation of encapsidated RNA, total RNA, and mRNA. The three aliquots were centrifuged at 1,000 rpm for five minutes and the supernatants discarded. Two aliquots of cell suspension were resuspended in 50 μ l of
10 RSB (10mM NaCl, 10mM Tris, pH 7.5, 1.5mM MgCl₂) containing 1% Triton X-100, 0.5% DOC. 50 μ l of 10 mM Tris 7.5, 1 mM CaCl₂, and 20 μ g (1mg/ml stock) of micrococcal nuclease was then added to one aliquot, and the other received the same mixture without micrococcal nuclease (Baker & Moyer, J. Virol. 62:
15 834-838 (1988), incorporated herein by reference in its entirety). The purpose of the micrococcal nuclease was to destroy nonencapsidated RNA, and the conditions used had been optimized in preliminary experiments. The mixtures were incubated at 30°C for 30 min and the RNA was isolated with
20 Trizol (Life Technologies) according to the procedure of the supplier. The third aliquot of cell suspension was processed for RNA purification with Trizol and the purified RNA was separated by oligo(dT) cellulose chromatography into polyadenylated and nonpolyadenylated fractions (Grosfeld et
25 al., J. Virol. 69: 5677-5686 (1995), incorporated herein by reference in its entirety). RNA samples were run on 1.5% agarose gels containing 0.44 M formaldehyde, transferred to nitrocellulose (Chomczynski, Anal. Biochem. 201: 134-139 (1992), incorporated herein by reference in its entirety),
30 hybridized with strand specific riboprobes, and quantified by phosphoimager analysis.

EXAMPLE III

Construction and Expression of Modified PIV3 Minigenomes

35 In the present example, a panel of cDNAs was constructed to encode PIV3 minigenomes which differed in length by single nucleotide increments. Transcription and RNA

replication in this reconstituted system were the most efficient for the minigenome whose length was an even multiple of six. In this context, members of the *Paramyxovirus* and *Morbillivirus* genera typically abide by a "rule of six," i.e., genomes (or minigenomes) replicate efficiently only when their nucleotide length is a multiple of six (thought to be a requirement for precise spacing of nucleotide residues relative to encapsidating NP protein). However, the present findings illustrate that minigenomes whose lengths were one nucleotide greater than or less than an even multiple of six were surprisingly active, especially in RNA replication.

A panel of seven cDNAs was constructed to encode seven PIV3-CAT minigenomes, called PIV3-CAT 0 to +6, that differ in length by single-nucleotide increments (Fig. 3). Each minigenome is a short negative-sense analog of HPIV3 genomic RNA in which the viral genes had been deleted and replaced with a negative-sense copy of the CAT ORF. The CAT ORF is flanked by nontranslated segments of the N and L genes, the N GS and L GE transcription motifs, and the 3' leader and 5' trailer extragenic termini of genomic RNA. The 5' end of each PIV3-CAT minigenome is defined by the adjacent promoter for T7 RNA polymerase. This promoter was constructed to contribute an extension of two nonviral G residues to the 5' end of the encoded minigenome, as described above. The presence of additional G residues adjacent to the nontranscribed core of the T7 promoter improves its transcriptional efficiency, and preliminary work showed that presence of two G residues provided the highest levels of activity in the reconstituted system described below. These two G residues are not included in minigenome length calculations. The 3' minigenome end is created by self-cleavage by an abutting hepatitis delta virus ribozyme, which would generate the correct 3'-terminal nucleotide. The seven PIV3-CAT minigenomes differ by single-nucleotide increases in length due to the insertion of 0 to 6 G residues at the junction between the N nontranslated region and the CAT gene (Fig. 3), and range in length from 898 to 904 nucleotides. The PIV3-CAT +2 minigenome is the only one which

is an even multiple of six (it should be noted that each minigenome contained two additional nonviral G residues at the 5' end contributed by the T7 promoter, but it is assumed that these would be lost during intracellular HPIV3-mediated RNA replication).

Each PIV3-CAT cDNA was transfected into HEp-2 cells that had been infected with vTF7-3, a vaccinia virus recombinant that expresses T7 RNA polymerase. Plasmids encoding the N, P and L proteins under the control of the T7 promoter were transfected in parallel. The P cDNA had been modified by site-directed mutagenesis to eliminate the translational start site of the C ORF, as described above. Cells were harvested at 48 h post-infection. An aliquot of the cell suspension was processed for CAT enzyme assay (Fig. 4). The remaining cells were divided into three equal aliquots and processed for RNA analysis as described below.

The minigenome cDNA was further modified to contain two tandem vaccinia virus early-gene transcription termination motifs (T₇NT) in the positive-sense plasmid strand upstream of the PIV3-CAT insert, and a third one (T₇AT) in the same strand immediately upstream of the CAT ORF (Fig. 3). These were designed to minimize promiscuous transcription of the CAT ORF by vaccinia virus polymerase (Grosfeld et al. (1995), *supra*). CAT expression was reproducibly detected when each of PIV3-CAT minigenomes was complemented by the N, P and L plasmids (Fig. 4), and detection of CAT was dependent on all three PIV3 proteins. However, expression was much higher for PIV3-CAT+2, which has a nucleotide length that is an even multiple of six. Preferred ratios and amounts of the minigenome and support plasmids were determined based on CAT enzyme expression.

EXAMPLE IV

Synthesis of Positive-Sense RNAs by PIV Minigenomes

Transcription and replication of PIV minigenomes was confirmed by detection of RNA products of both processes. As described in the foregoing Example, three equal aliquots of cell suspension were taken for RNA analysis. One aliquot was

used for oligo(dT) analysis, as described below. The other two aliquots were lysed with detergent and incubated with micrococcal nuclease or mock-treated. RNA was then isolated, separated by electrophoresis on formaldehyde agarose gels, transferred to nitrocellulose, and analyzed by hybridization with negative-sense CAT riboprobe. RNA from micrococcal-treated and mock-treated lysates are shown in Fig. 5A upper and lower panels, respectively.

Analysis of RNA from mock-treated lysates showed that complementation of each minigenome with the N, P and L plasmids resulted in the synthesis of a band of RNA which was very similar in size to a marker consisting of RNA expressed by the 931-nucleotide RSV-CAT C2 minigenome (Grosfeld et al. (1995), supra). Phosphorimagery analysis is shown in Fig. 5B. Little or no RNA was detected when the N or L plasmids were omitted, confirming that these RNAs are products of the reconstituted PIV3 polymerase.

Each PIV3-CAT minigenome is expected to encode two positive-sense RNAs, namely the mini-antigenome and the subgenomic CAT mRNA. Each mini-antigenome is expected to be the exact complement of its minigenome, which was 898 to 904 nucleotides in length. The predicted subgenomic mRNA is defined by the GS and GE signals, and is expected to be 804 nucleotides in length and contain a polyA tail of 100 to 200 nucleotides.

Detection of a single gel band of positive-sense RNA in Fig. 5A (lower panel) suggested that the antigenome and mRNA were not resolved by gel electrophoresis. Accordingly, treatment with micrococcal nuclease was used to identify antigenome RNA, since the antigenome (and genome) but not mRNA would be encapsidated and resistant to digestion. The use of micrococcal nuclease for this purpose is well established (Baker & Moyer (1988), supra), and the conditions selected were verified with RSV minireplicons and shown to completely degrade mRNA contained in the HEp-2 cell lysates. Residual RNA was purified and analyzed by Northern blot analysis with negative-sense riboprobe (Fig. 5A, upper panel) and quantitated by phosphorimagery (Fig. 5B; note that in this

analysis the micrococcal-treated and untreated RNA amounts were normalized separately). These investigations revealed the presence of a population of protected RNA corresponding to the positive-sense encapsidated mini-antigenome. Among
5 several experiments, this protected RNA accounted for approximately 3 to 15% of the positive-sense RNA.

For both the total and the micrococcal-resistant RNA, accumulation was greatest in the case of the +2 minigenome, which is 900 nucleotides in length and thus a
10 multiple of six. However, substantial amounts of RNA also accumulated in the case of the minigenomes which did not exhibit a length corresponding to a multiple of six nucleotides, in particular minigenomes +1 and +3 which were one nucleotide longer or shorter than the +2 minigenome. In
15 fact, the amount of encapsidated antigenome produced by the +1 and +3 minigenomes was 85% and 72% that of the +2 minigenome (Fig. 5B). Even the least efficient minigenome, the +5 minigenome, was 20% as active as the +2 minigenome as determined by measurement of accumulated encapsidated RNA. In
20 the case of measurements to detect total positive-sense RNA, the +1 and +3 minigenomes produced 52% and 45% as much total RNA as the +2 minigenome.

To confirm the presence of subgenomic mRNA, the final aliquot of harvested cell suspension was processed for
25 RNA purification. The RNA was then subjected to oligo(dT) chromatography. RNAs which failed to bind, and those which bound and were eluted in low salt buffer, were analyzed by Northern blot hybridization (Fig. 6A) and phosphorimager (Fig. 6B; note that in this case the bound and unbound are
30 normalized together relative to the bound RNA of the +2 minigenome). These assays showed that approximately 64% of positive-sense RNA was polyadenylated, as expected for subgenomic mRNA. The accumulation of mRNA was greatest for the +2 minigenome. However, substantial amounts of mRNA also
35 were observed for the other minigenomes. The amount of mRNA synthesized by the +1 and +3 minigenomes was 30% and 20% respectively compared to that synthesized by the +2

minigenome, and was approximately 13% for the least active minigenomes.

EXAMPLE V

5 Synthesis of Negative Sense RNA by PIV Minigenomes

The various PIV3-CAT minigenomes described in the foregoing examples directed synthesis of mRNA and positive-sense encapsidated mini-antigenome, the latter
10 representing the first step in RNA replication. The second step in RNA replication involves synthesis of encapsidated progeny minigenome from the mini-antigenome product. To evaluate this latter process, the samples of RNA from
15 mock-treated and nuclease-treated lysates described in the preceding Example were analyzed by Northern blot hybridization with positive-sense CAT riboprobe (Fig. 7A) and quantitated by phosphorimagery (Fig. 7B).

Analysis of RNA from mock-treated lysates (Fig. 7A, lower panel) showed that considerable amounts of minigenome
20 accumulated intracellularly in all samples, including negative controls in which the N or L support plasmid was omitted. The analyses described in Figs. 5A-B and 6A-B showed that the synthesis of positive-sense RNA was insignificant under these conditions. Therefore, the minigenome observed in the absence
25 of N or L could not be the product of RNA replication mediated by the reconstituted HPV3 polymerase, and instead must be the product of T7 transcription of transfected plasmid.

Minigenome produced by the reconstituted HPV3 polymerase is expected to be encapsidated, whereas much of the
30 minigenome produced by T7 RNA polymerase is expected to be unencapsidated. Therefore, RNA from the same micrococcal nuclease-treated samples described for Figs. 5A-B were used to prepare a second blot, which was hybridized with positive-sense CAT riboprobe (Fig. 7A, upper panel). This
35 showed that all minigenome RNA accumulated in the absence of the N protein was degraded (Fig. 7A, upper panel, lane 1), as expected. Essentially all of the minigenome which accumulated in the absence of L was also sensitive to degradation (Fig.

7A, upper panel, lane 2). Plasmid-derived minigenome synthesized in the absence of L, and in the presence of N and P alone, did not appear to occur efficiently.

When the complete set of three support plasmids was present, significant amounts of micrococcal nuclease-resistant minigenome RNA accumulated for each of the minigenomes (Fig. 7A, upper panel). As was the case with the positive-sense RNAs, the greatest amount of progeny minigenome was observed with the +2 minigenome. The +1 and +3 minigenomes were next in abundance, with levels of genomic RNA that were 67% and 42% of that of the +2 minigenome.

The foregoing examples demonstrate that the HPIV3 N, P and L proteins were necessary and sufficient for efficient transcription and RNA replication. The very robust nature of transcription and RNA replication mediated by the reconstituted PIV3 polymerase confirmed the functionality of the encoded proteins. It is further expected that inclusion of additional viral proteins within the expression system will augment or modify these processes. Coexpression of PIV C, D and potentially V, within the compositions and methods of the invention will be useful, e.g., to augment and/or modify RNA replication. For this purpose, plasmids will be constructed and assayed according to the foregoing methods to achieve coexpression of one or more of these elements to determine their effects on PIV transcription and RNA replication, as well as on PIV phenotype in suitable infection models.

EXAMPLE VI

Construction of Infectious, Recombinant PIV from cDNA

The following examples describe production of infectious, recombinant PIV (rPIV) by intracellular co-expression of four plasmid-borne cDNAs. These cDNAs separately encode a complete HPIV3 genome and the HPIV3 nucleocapsid protein N, the phosphoprotein P, and the polymerase protein L.

A) Viruses and cells

Modified vaccinia strain Ankara (MVA), the vaccinia virus recombinant that expresses bacteriophage T7 RNA polymerase, was provided according to Wyatt et al., Virology 210:202-205, 1995, incorporated herein by reference in its entirety. HEp-2 monolayer cultures were maintained at 37° C in 5% CO₂ with OptiMEM 1 (Life Technologies) supplemented with 2% FBS, 50µg/ml gentamicin sulfate and 2mM glutamine. The JS wt strain of HPIV3 and its attenuated *ts* derivative, JS cp45, were propagated in LLC-MK2 cells as described by Hall et al., Virus Res. 22: 173-184, (1992), incorporated herein by reference in its entirety.

B) cDNAs

The full cDNA clone designated p218(131) (Fig. 1, SEQ ID NO: 1) encoding a complete genomic sequence of HPIV3 was constructed as described above. A second cDNA clone designated p3/7(131) (SEQ ID NO: 14) (deposited under the terms of the Budapest Treaty with the ATCC, and granted the designation 97990) was constructed to encode a complete antigenomic sequence of HPIV3. p3/7(131) differs from p218(131) in that the positions of the T7 promoter and the ribozyme/T7 terminator relative to the HPIV3 cDNA insert have been interchanged. Thus, the first nucleotide synthesized in p3/7(131) is the 5' end of the antigenome, namely the positive-sense complement of genome position 1, and the 3' antigenome end defined by ribozyme cleavage is the complement of genome position 15462. A third clone designated p3/7(131)2G (SEQ ID NO: 15) (deposited under the terms of the Budapest Treaty with the ATCC, and granted the designation 97989) was also constructed identical to p3/7(131), except that two G residues were inserted between the 5' end of the antigenome and the T7 promoter.

For construction of p/37(131)2G and p3/7(131), the two plasmids p(A*A'CC'D*E) and p(E*FF'GHIJKL*) were modified and joined to encode the complete positive-sense antigenome of HPIV3. First, the T7 terminator and delta ribozyme abutting the 3' end of HPIV3 in p(A*A'CC'D*E) were replaced by a T7 promoter using PCR (see Fig. 8). The positive-sense primer

5'-GGCCCGTCGACGCGTAATACGACTCACTATAGGACCAAACAAGAG-3' (SEQ ID NO: 18) placed the T7 promoter (bold) adjacent to the 3' leader of HPIV3 (HPIV3 sequence underlined) with 2 G residues inserted between these two elements to improve transcription. A unique *Mlu* I site (italicized) was placed upstream of the T7 promoter for cloning purposes (Fig. 8). The negative-sense primer 5'-¹²²⁴CGGCATCACGTGCTAC¹²⁰⁹-3' (SEQ ID NO: 19) spanned nt 1209-1224 of the HPIV3 N gene and included a unique *Pml* I site (italicized) present in the natural HPIV3 sequence. Both the PCR product and the parent template p218(A*A'CC'D'E) were digested with *Mlu* I and *Pml* I and the PCR product was then cloned into the *Mlu* I-*Pml* I window. A second PCR reaction was done using the same negative-sense primer and a positive-sense primer without 2G residues inserted between the T7 promoter and 3' end of HPIV3. These plasmids were designated p(Left + 2G) and p(Left +). The construction of p(Left + 2G) is illustrated in Fig. 8, the construction of p(Left +) followed the same strategy.

Plasmid p(E*FF'GHIJKL*) was modified by PCR to place the delta ribozyme and T7 terminator adjacent to the 5' end of HPIV3 (Fig. 9). The positive-sense primer:

5'-GGATTTGCGCGC¹⁴⁸¹³AATTAAATCATCTGG¹⁴⁸²⁸-3' (SEQ ID NO: 16)

introduced a *BssH* II site (italicized) just upstream of a unique *Swa* I site (italicized, underlined) in the L gene of HPIV3 (HPIV3 specific sequence is underlined). The negative-sense primer:

5'-CCCAGGTTCGGACCGCGAGGAGGTGGAGATGCCATGCCAGCCC¹⁵⁴⁶²ACCA
AAACAAGAGAAGAACTCTGTTTGG¹⁵⁴³⁵-3' (SEQ ID NO: 17) placed a

portion of the delta ribozyme (bold) which included a naturally-occurring unique *Rsr* II site (italicized) adjacent to the 5' end of the HPIV3 genome (negative-sense underlined). This PCR product was digested with *BssH* II and *Rsr* II and cloned into the *BssH* II-*Rsr* II window of plasmid p3/7 (Fig. 9). Plasmid p3/7 is identical to p218 except that the delta ribozyme and T7 terminator are in the reverse orientation. The p3/7 plasmid containing the above-mentioned insertion was designated pPIV3-3/7 and contained the complete delta ribozyme and T7 terminator adjacent to 5' terminal 651 nucleotides of

HPIV3. The *Swa*I and *Ngo*MI fragment of plasmid pPIV3-3/7 was then isolated and cloned into the *Swa*I-*Ngo*MI window of p(E*FF⁻GHIJKL*). The resulting plasmid, designated p(Right +) placed the complete delta ribozyme and T7 terminator adjacent to the 5' end of HPIV3 (see Fig. 9). The *Xho*I -*Ngo*MI fragment of p(Right +) was cloned into the *Xho*I -*Ngo*MI window of p(Left +) and p(Left + 2G) resulting in plasmids p3/7(131) (SEQ ID NO: 14) and p3/7(131 2G) (SEQ ID NO: 15), respectively (Fig. 10). These each encode the complete positive-sense analog of HPIV3 antigenomic RNA, with the latter cDNA containing two G residues adjacent to the T7 promoter for improved transcriptional efficiency.

C) Transfection

HEp-2 cells were grown to 90% confluence in six well plates. Each well of a six-well plate (1.5×10^6 cells) was transfected with the three previously-described support plasmids, 0.4 μ g pTM(P), 0.4 μ g pTM(N), 0.05 μ g pTM(L), together with 5 μ g of full-length genomic or antigenomic HPIV3 cDNA. The plasmids were added to 0.2 ml of OptiMEM 1 (Life Technologies) containing 12 μ l of LipofectACE (Life Technologies). After an incubation period of approximately 15 minutes at room temperature, 0.8 ml of OptiMEM 1 containing 2% fetal bovine serum and 1.5×10^7 pfu of MVA-T7 was added to each well. The cultures were incubated at 32°C for 12 hours, after which the media was replaced with fresh OptiMEM 1 containing 2% fetal bovine serum. The cultures were incubated at 32°C for three additional days then harvested and passaged (referred to as passage 1) onto fresh HEp-2 monolayers. These passage 1 cultures were incubated at 32°C for five days, and virus present in the culture was harvested, passaged once in LLC-MK2 cultures, and characterized by hemagglutination-inhibition (HAI) as described (van Wyke Coelingh et al., Virology 143: 569-582, (1985) (incorporated herein by reference in its entirety) to determine if it possessed the monoclonal antibody resistant mutation (MARM) that marked virus recovered from cDNA.

D) Sequencing of Recombinant Virus

The presence of nucleotide sequence markers in the HN and L genes of recombinant PIV was determined by RT-PCR of RNA isolated from recovered virions. 1 ml of rPIV (1 X 10⁵ pfu/ml, passage level 2) was precipitated with 200 µl 25% polyethylene glycol by incubation on ice for one hour and centrifuging at 12,000g for 15 minutes. The RNA was purified with TRIzol reagent (Life Technologies) following the manufacturer's recommended procedure. RT-PCR was performed with the Superscript kit (Life Technologies) following the manufacturer's recommended protocol. Control reactions were identical except that reverse transcriptase was omitted from the reaction to confirm that the PCR products were derived solely from virus RNA and not from possible contaminating cDNA plasmids. Four primer pairs were used to generate PCR products from nt 7334-8715, 9364-10854, 10939-15392, and 13623-15392. The resultant PCR products were then sequenced using cycle dideoxynucleotide sequence analysis (New England Biolabs, Beverly, MA).

EXAMPLE VIIRecovery of Recombinant Virus From cDNA Encoding
Negative-Sense Genomic RNA

Plasmid p218(131) and the three support plasmids pTM(N), pTM(P), and pTM(L) were transfected into HEp-2 cells with MVA expressing T7 RNA polymerase. A control group consisting of pTM(N), pTM(P), pTM(L), and MVA was cotransfected into HEp-2 cells without p218(131). On day four, the transfection was harvested, passaged onto fresh HEp-2 cell monolayers for five days, and passaged again for 5 days in LLC-MK2 cultures (passage 2). Virus present in the passage 2 harvest was further characterized by HAI. Cultures from the transfection group which received the three support plasmids without the full-length genomic clone p218(131) did not yield HPIV3. The rPIV recovered virus was confirmed to be HPIV3 since it reacted in the HAI assay with the mAbs 77/5, 101/1, and 454/11 which are specific for HPIV3 (Table 1). It

was presumptively identified as being cDNA-derived because it failed to react with mAbs 170/7 and 423/6, consistent with the MARM mutation which had been introduced into the cDNA.

TABLE 1

rJS-NS Contains the MARM Mutation Introduced into the
p218(131) Full-Length Negative Sense cDNA

Virus	Hemagglutination-inhibiting		Antibody Titer	(Reciprocal) of Indicated mAb	
	77/5 ¹	101/1 ¹		454/11 ¹	170/1 ²
JS wt ³	800	6400	6400	25,600	25,600
rJS-NS ⁴	3200	25,600	6400	≤ 25	≤ 25

¹ mAb 77/5 recognizes antibody epitope IIB, mAbs 101/1 and 454/11 recognize antibody epitope IIIA of HN glycoprotein, all of which were not altered in p218(131).

² Mabs 170/7 and 423/6 which both recognize antibody epitope I of JSwt, fail to recognize rJS due to the MARM mutation at this site.

³ Biologically derived wild type HP1V3 JS.

⁴ Recombinant JS virus derived from negative-sense genomic cDNA.

To confirm that rPIV was indeed recovered from cDNA, it was analyzed in parallel with wild-type JS strain HPIV3 by RT-PCR using four primer pairs flanking the seven inserted marker mutations. Each PCR product obtained was dependent upon the inclusion of RT, indicating that each was derived from RNA and not from contaminating cDNA. Cycle-sequencing of the four PCR products confirmed that the sequence of the rPIV contained each of the seven markers, sequencing data showing three of the markers is illustrated in Figure 11. The sequence differences between rPIV and JS wt in the HN gene, including nt 7903, 7913, and 7915 are readily apparent. Similar sequence analyses were carried out for the other four markers at nt positions 7593, 10355, 11333, and 15248, and the rPIV possessed each mutation.

These results demonstrate successful recovery of infectious rPIV from cDNA encoding a negative-sense genomic RNA. This differs from most published reports for recovery of nonsegmented negative strand RNA viruses, in which the cDNA used for virus recovery had been designed to encode positive-sense antigenomic RNA (Baron and Barrett, supra, 1997; Collins et al., supra, 1995; Conzelmann, supra, 1996; Garcin et al., supra, 1995; Lawson et al., supra, 1995; Radecke et al., supra, 1995; Whelan et al., supra, 1995). The recovery of infectious virus from a cDNA encoding genomic RNA had previously been reported only in the case of Sendai virus, and the efficiency of recovery was much lower than for cDNA encoding antigenomic RNA (Kato et al., 1996). In most other studies, the recovery of virus was achieved with antigenomic cDNA but not with genomic cDNA (Lawson et al., 1995; Whelan et al., 1995). A number of potential problems have been noted which may explain these refractory results, including possible annealment of cDNA-encoded genomic RNA with mRNA produced by the support plasmids; resulting in inactive hybrids (Conzelmann, supra, 1996; Lawson et al., supra, 1995). It has also been noted that T7 RNA polymerase appears to terminate preferentially at the gene junctions of genomic RNA, perhaps because the oligo U tract of the GE signal resembles

the natural signal for transcription termination by T7 RNA polymerase (Whelan et al., supra, 1995).

5

EXAMPLE VII

Recovery of Recombinant Virus From cDNA Encoding Positive-Sense Antigenomic RNA

10 As described in more detail above, p3/7(131) and
p3/7(131)2G were constructed to encode a positive-sense,
antigenome that give rise to recombinant PIV. Plasmid
p3/7(131)2G is identical to p3/7(131) but for the addition of
two G residues between the T7 promoter and the first
nucleotide of the antigenome. The addition of two G residues
15 between the T7 promoter and the first HPIV3 nucleotide
p3/7(131)2G is based on the preceding examples demonstrating
that the presence of the two added G residues (as opposed to
0, 1 or 3 added G residues) yielded substantially increased
levels of minireplicon replication.

20 The two antigenome cDNAs [p3/7(131) and p3/7(131)2G]
were transfected separately into cells together with the N, P
and L support plasmids, and were infected simultaneously with
the MVA-T7 recombinant virus using the same procedure
described above for p218(131). Infectious virus from each
25 antigenomic cDNA was recovered and was presumptively
identified as being cDNA-derived by its inability to react
with mAbs 423/6 and 170/7.

The efficiency of virus recovery was evaluated for
each of the antigenome cDNAs versus the genome cDNA p218(131).
30 Twelve transfection reactions using the negative-sense genome
cDNA p218(131) (SEQ ID NO: 1) were conducted in parallel with
twelve transfections using the positive-sense antigenome cDNA
p3/7(131)2G (SEQ ID NO: 15) to compare efficiency of virus
recovery from the two cDNAs. One ml of the transfection
35 harvest from each well was titered on LLC-MK2 cells and the
remaining 2mls were passaged (passage 1) onto fresh LLC-MK2
cells. At day five, passage 1 was harvested and titered as
described above. Recombinant virus was recovered from 12/12

wells transfected with p3/7(131)2G but from only 4/12 wells transfected with p218(131). The mean titer of virus present in culture of the passaged virus from the positive-sense antigenome was $10^{5.0}$, nearly ten-fold higher than that from the
5 negative-sense genome, which was $10^{4.1}$. However, with one additional amplification the titers became equivalent.

TABLE 2

Comparison of the Efficiency of Recovery of Recombinant Virus from cDNAs Encoding Genomic or Antigenomic RNA, the latter with or without 2G residues adds to the T7 promoter.

Experiment#	Transfected cDNA	Sense of Encoded RNA	Rescue Efficiency ¹	Mean Titer (log ₁₀ pfu/ml) of Recovered Virus ²
#1	p3/7(131)2G ³ p218(131)	Antigenomic	12/12 (100%)	5.0 ± 0.27
		Genomic	4/12 (33%)	4.1 ± 0.27
#2	p3/7(131)2G ³ p3/7(131) p218(131)	Antigenomic	6/6 (100%)	6.6 ± 0.18
		Antigenomic	12/12 (100%)	5.3 ± 0.12
		Genomic	5/5 (100%)	4.9 ± 0.19

72

- 1 Number of transfection cultures yielding rJS/number tested.
- 2 The mean Titer ± standard error was determined following one passage of transfection harvest for five days in LLC-MK2 cells.
3. Contains 2 G residues between the T7 promoter and the 5' end of the antigenome.

The efficiency of recovery of recombinant virus from the three full-length plasmids encoding the genomic or antigenomic HPIV3 RNAs was next studied, (i) to determine whether genomic or antigenomic cDNA is more efficient at generating recombinant virus and (ii) to determine the effect of two extra 5' terminal G residues on the yield of recombinant viruses (Table 2.) Unfortunately, it was not possible to directly titer the transfection harvest by plaque titration because residual MVA-T7 interfered with plaque formation of rJS on LLC-MK2 monolayer cultures. Therefore, we first compared the efficiency of recovery of rJS from multiple independent transfections and, as seen in other experiments, found that rJS was more frequently recovered from cultures transfected with p3/7(131)2G than from cultures transfected with p218(131). Since each transfection harvest, including those with p218(131) yielded rJS in other experiments, it was not possible to use this method of analysis to compare the relative efficiency of generating recombinant virus. We therefore estimated the relative quantity of virus present in the transfection harvest by quantitating the amount of virus following one passage of the transfection harvest (Table 2). Since each plasmid was designed to yield an identical virus (rJS), differences in titers of the passaged virus were considered a reflection of differences in the titer of virus present in transfection harvest. In the above experiments, the construct with two 5' terminal G residues was the most efficient in generating rJS. This indicated that the two added residues indeed increased efficiency of recovery, but the mechanism for this small increment in efficiency was not defined. These findings indicate a small, but measurable, advantage of using p3/7(131)2G as substrate for future experiments designed to introduce mutations into the HPIV3 cDNA. The small advantage in recovery of recombinant virus from p3/7(131)2G might be especially needed to recover viruses that have attenuating mutations that restrict replication in vitro. Although VSV has been recovered from antigenomic cDNA which included three G residues at the 5'-terminal end, the efficiency of recovery of virus from this construct was not

compared with recovery from the same construct lacking the extra residues (Whelan et al., Proc. Natl. Acad. Sci. USA 92:8388-8392 (1995), incorporated herein by reference). In contrast to the experience with Sendai virus and measles virus, it is clear that extra-viral G residues at the 5'-terminal end of the antigenome are not deleterious to recovery of recombinant virus, and, in fact, appear to be advantageous [Garcin et al., supra, 1995; Radecke et al., supra, 1995; Kato et al., 1996). The genomic and antigenomic plasmids without two 5'-terminal guanine residues appeared equally efficient in generating recombinant virus. This is in contrast to previous reports which have found genomic cDNA to be less efficient than antigenomic cDNA in generating recombinant virus (Whelan, 1995; Kato et al., 1996).

The efficiency of recovery of virus from p3/7(131) (SEQ ID NO: 14) was compared to that of p3/7(131)2G (SEQ ID NO: 15 and p218(131) (SEQ ID NO: 1) in a similar experiment as above. Recovery from p3/7(131) was comparable to that of p217(131). Recovery was more consistent and efficient with both antigenome cDNAs containing the extra Gs (Table 2).

EXAMPLE VII

Efficiency of plaque formation of rPIV at permissive and restrictive temperatures

The level of temperature sensitivity of replication *in vitro* of control viruses and rPIV derived from both p218(131) and p3/7(131)2G was determined at 32°C, 37°C, 39°C, and 40°C in LLC-MK2 monolayer cultures as previously described (Hall et al., Virus Res. 22: 173-184, (1992), incorporated herein by reference in its entirety), with the following modifications. Viruses were inoculated onto LLC-MK2 monolayers on 24-well plates in serial 10-fold dilutions allowed to adsorb for one hour at room temperature. The cultures were then overlaid with 1ml of L-15 supplemented with 2mM glutamine and 0.8% methylcellulose and the plates were then incubated for 5 days at the indicated temperature. The methylcellulose overlay was removed and the monolayer fixed

with 80% methanol at 4°C for 1h. The viral plaques present in the monolayer were scored using a mixture of two HPIV3-specific anti-HN murine mAbs 101/1 and 66/4 as ascites fluid used at a dilution of 1:500, using an immunoperoxidase method of staining specific for murine antibodies as previously described (Murphy et al., Vaccine 8: 497-502 (1990), incorporated herein by reference in its entirety).

Recombinant virus derived from either positive or negative-sense cDNA were characterized by plaque assay at 32°C, 37°C, 39°C, and 40°C to determine if they were phenotypically similar to JS wt virus. Both positive and negative-sense rPIV were comparable to the JS wt virus in their level of replication at elevated temperatures of 39°C and 40°C (Table 3). This is in contrast to the *ts* mutant JS cp45 which exhibits a 30-fold reduction in titer at 37°C and fails to produce plaques at 39°C or 40°C.

TABLE 3

The rJS Resembles its Biologically Derived Parent JS Wild-Type Virus in the Level of Replication at Restrictive Temperature (39°C - 40°C)

Virus	Virus Titer (log 10 pfu/ml)			
	32°C	37°C	39°C	40°C
rJS-PS ¹	6.1	6.1	6.1	6.6
rJS-NS ²	6.9	7.1	7.1	7.0
JS cp45 ³	6.3	4.3	<0.7	<0.7
JS wt	6.5	6.8	6.6	6.7

¹ Recombinant virus derived from the antigenomic-sense clone p3/7(131)2G

² Recombinant virus derived from the genomic-sense clone p218(131)

³ JS cp45 is a temperature sensitive mutant derived from JS wt.

The sequence of JS *cp45* has been fully determined (Stokes et al., supra, 1993) and mutations have been identified in the leader, N, P, M, F, HN, and L genes. However, it is unknown which mutation(s) are responsible for the *ca*, *att*, or *ts* phenotypes. Because exemplary rPIV of the invention demonstrate the *ts*' phenotype like the JS wt parent, *cp45* mutations among other mutations known or yet to be discovered for PIV can be introduced, alone or in combination, into the full-length cDNA to pinpoint the effects of individual mutations or combinations thereof, e.g., by evaluating replication of the recombinant virus incorporating the mutation(s) of interest at elevated temperatures. The mutation(s) thus identified can be incorporated into effective, attenuated vaccine agents as described in the Examples below. These and other mutations incorporated into recombinant PIV can be further optimized by, e.g., mutagenesis to create two or more nucleotide substitutions per codon to render a recombinant virus that is more genetically stable than a biologically derived mutant strain.

EXAMPLE VIII

Replication of rPIV in Hamsters

Thirty-six 16-week-old golden Syrian hamsters were divided into four groups of nine and inoculated intranasally with 0.1ml containing $10^{5.5}$ pfu of either rPIV recovered from negative-sense cDNA, rPIV recovered from positive-sense cDNA, JS *cp45*, or JS wt virus. On day 4, the hamsters were sacrificed and the lungs and nasal turbinates harvested. The lungs were homogenized in a 20% w/v L-15 suspension containing 2.5µg/ml amphotericin B (Quality Biologicals, Gaithersburg, MD), 200µg/ml piperacillin (Lederle Laboratories, Pearl River, NY), and 50µg/ml gentamicin (Quality Biologicals). The nasal turbinates similarly were homogenized in a 10% w/v L-15 suspension. After homogenization, the samples were aliquoted and rapidly frozen in a dry ice-ethanol bath. Virus present in the samples were titered at a later date in 96 well plates

of LLC-MK2 cells at 32°C scoring CPE at five and seven days after inoculation. The mean \log_{10} TCID₅₀/gm was calculated for each group of nine hamsters.

Table 4 illustrates that rPIV recovered from negative-sense cDNA, rPIV recovered from positive-sense cDNA replicate to substantially the same level as the JS wt in the upper and lower respiratory tract of hamsters. This is in contrast to the JS cp45 virus, which is attenuated at each site.

TABLE 4

The rJS Resembles its Biologically Derived Parent JS wt Virus in the Level of Replication in the Upper and Lower Respiratory Tract of Hamsters.

Virus	Mean Virus Titer (\log_{10} TCID ₅₀ /g) ⁵	
	Nasal Turbinates	Lungs
rJS-PS ¹	6.6 ± 0.2	4.1 ± 0.3
rJS-NS ²	6.4 ± 0.1	4.2 ± 0.2
JS cp45 ³	4.2 ± 0.2	≤ 1.4 ± 0.0
JS wt ⁴	6.3 ± 0.2	4.6 ± 0.3

¹ Recombinant virus recovered using p3/7(131)2G encoding the positive-sense HPIV3 antigenome.

² Recombinant virus recovered using p218(131) encoding the negative-sense HPIV3 genome.

³ Biologically derived *ts* mutant.

⁴ Biologically derived parent virus.

⁵ Mean titers ± standard errors for nine hamsters per group.

Thus, exemplary rPIVs of the invention can retain the replicative capacity in hamsters exhibited by the biologically derived JS wt parent strain, whereby mutations such as those present in the JS cp45 candidate vaccine that restrict replication in hamsters and other hosts, including

non-human primates and humans, can be identified and incorporated within modified rPIV strains of the invention, as described in the Examples below.

5

EXAMPLE IX

Identification of Amino Acid Substitutions in HPIV3 Specifying Attenuated Phenotypes, and Incorporation of Attenuating Mutations into Infections, Attenuated PIV Clones

10 The ability to generate infectious PIV from cDNA facilitates development of live-attenuated parainfluenza virus vaccines. More specifically, by using the methods and tools disclosed herein the genetic basis of attenuation of PIV candidate vaccines can be readily determined, and infectious
15 PIV vaccines produced from cDNA can be designed to achieve a finely calibrated level of attenuation and immunogenicity.

 In addition, the tools and methods of the invention provide for vaccine development for all three human parainfluenza viruses, HPIV1, HPIV2 and HPIV3 that are most
20 important in human disease. For example, to produce and select effective HPIV3 vaccine agents within the invention, mutations associated with desired phenotypes of biologically deriving HPIV3 candidate vaccines or the attenuated BPIV3 virus, e.g. attenuating mutations, can be identified and
25 incorporated into rPIV. Applying these methods, attenuating mutations from a large menu of candidate mutations are selected and combined to generate rPIV having a desired balance between attenuation and immunogenicity, and which retain the attenuation phenotype following replication in
30 humans.

 In the present example, the genetic bases of temperature-sensitive (ts) and in vivo attenuation (att) phenotypes of the PIV3 JS cp45 live-attenuated virus are described. Seven exemplary recombinant PIV3 viruses (three
35 single-, three double-, and one triple-lesioned virus) were recovered from full-length antigenomic cDNA and analyzed for their ts and att phenotypes. These recombinants bore one or more amino acid substitution mutations present in the L gene

of JS cp45 (alternatively referred to herein as cp45), adopted within a cDNA clone of the JS wt parent. These three exemplary, biologically derived mutations are all present in a representative strain of JS cp45 grown in Vero cells, designated JS cp45 Vero, deposited on August 21, 1997 under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., and granted the accession number ATCC VR 2588.

Analyses of exemplary PIV recombinants, presented below, demonstrate that each of the three exemplary mutations in L (Tyr₉₄₂ to His, Leu₉₉₂ to Phe, and Thr₁₅₅₈ to Ile) contribute to the ts and att phenotypes of cp45 and are useful for generating of recombinant vaccine virus.

Viruses and cells.

The PIV3 JS wt and cp45 viruses were grown in LLC-MK2 cells as described previously (Hall et al., Virus Res. 22:173-184 (1992), incorporated herein by reference). The vTF7-3 recombinant vaccinia virus is described in Fuerst et al., Virology 225: 419-422 (1996) and the modified vaccinia virus Ankara (MVA) which expresses the T7 polymerase is described in Wyatt et al., Virology 210:202-205 (1995) (each incorporated herein by reference). HEp-2 (ATCC CCL 23) and LLC-MK2 (ATCC CCL 7.1) cells were maintained in OptiMEM (Life Technologies) supplemented with 2% FBS and gentamicin sulfate (50ug/mL).

Construction of point mutations in the L gene of PIV3.

pUC19 was modified to accept a fragment of the JS wt PIV3 L gene in order to introduce point mutations into the L gene by site-directed mutagenesis. First, a unique Nhe I restriction site was introduced into pUC19 by ligating a pair of complementary oligonucleotides (5' GATCGATGCTAGCCC 3' (SEQ ID NO: 23) and 5' GATCGGGCTAGCATC 3' (SEQ ID NO: 24)) containing an Nhe I restriction site into the Hind III site of pUC19 to create pUC19 (N). The Sph I (PIV3 nt 11317) to Nhe I (PIV3 nt 14087) fragment of pTM(L), which includes the

positions where the three coding changes in cp45 occur and which can be directly introduced into the full-length PIV3 cDNA (see below), was cloned into the *Sph* I and *Nhe* I site of pUC19 (N) to create pUCL(N-S). Point mutations were
5 introduced into pUCL(N-S) using mutagenic oligonucleotides with the Transformer mutagenesis kit (Clontech, Palo Alto, CA) for the purpose of (i) creating exemplary amino acid substitutions at L protein positions 942, 992, and 1558, individually and in combination, and (ii) ablating one
10 specific naturally-occurring restriction enzyme site proximal to each codon substitution as a marker [See Table 5].

TABLE 5: Nucleotide substitutions introduced into rPIV3 that encode cp45L protein gene amino acid substitutions and, as markers, ablate naturally-occurring restriction enzyme sites.

rPIV3 designation	Amino Acid Substitution (wt to cp45)	Sequence of wt ^a	Sequence of Mutant	Restriction Enzyme site Ablated
r942	Tyr-942 to His	11468- <u>TTACATGGCCAT</u> (SEQ ID NO: 25)	11468-TCACATGGCGAT (SEQ. ID NO: 26)	Eae I
r992	Leu-992 to Phe	11618- <u>TTTGTGATTGGGC</u> (SEQ ID NO: 27)	11618- <u>TTTGTGATTGGGC</u> (SEQ. ID NO: 28)	Bsr I
r1558	Thr-1558 to Ile	13307-TGGTCCTAATACTG (SEQ. ID NO: 29)	13307-TGGGCCTAATAATCG (SEQ. ID NO: 30)	Ava II

a. The nucleotide sequence around each of the three mutated regions is shown. The first nucleotide in each provided sequence is numbered according to its position in the complete antigenomic RNA. The codon involved in each amino acid substitution is in bold. Naturally-occurring restriction enzyme sites present in the wt sequence, and which were ablated to mark the mutation, are in italics. The nucleotides that were mutated to produce an aa substitution or remove a restriction enzyme site are underlined.

Mutations introduced in pUCL(N-S) derivatives were verified by dideoxynucleotide sequencing of plasmid DNA. The *Sph* I to *Bam*HI (nt 13733) fragment of pUCL(N-S) containing the *cp45* individual L gene mutations was subcloned into the *Sph* I to *Bam*HI sites of pTM(L) to give pTM(L)-942, -992, -942/992, and -1558; the other double and triple mutations were assembled using the *Pvu* AI and *Nhe* I sites (Fig. 12). The mutant pTM(L) plasmids were each tested at permissive temperature (32°C) for the ability to direct the expression of the chloramphenicol acetyl transferase marker gene in a minireplicon system comprising a plasmid-encoded minigenome RNA and the N, P and L proteins (Durbin et al., Virology 234:74-78 (1997), incorporated herein by reference). The various mutant L plasmids supported marker gene expression to 75-106% the level of wt L, indicating that each engineered cDNA was free of significant spurious mutation (not shown). The *Sph* I to *Nhe* I fragments of each of the mutant pTM(L) plasmids were then subcloned into the *Sph* I to *Nhe* I window of the full-length PIV3 JS antigenomic cDNA p3/7(131)2G to create seven full-length PIV3 cDNA clones representing every possible combination of the three substitutions.

Recovery of recombinant PIV3 (rPIV3) bearing one, two or three *cp45* L protein substitutions.

Each full-length antigenomic cDNA bearing one or more *cp45* L gene mutations, together with the three support plasmids pTM(N), pTM(P) and pTM(L), was transfected into HEp-2 cells on 6-well plates (Costar, Cambridge, MA) using LipofectACE (Life Technologies) and MVA-T7 as described above. After incubation at 32°C for 4 days, the transfection harvest was passaged onto HEp-2 cells on 6-well plates which were incubated at 32°C for 4 days. Each passage 1 supernatant was harvested and passed onto a T-25 flask of LLC-MK2 cells, which was incubated at 32°C for 5-6 days. The passage 2 supernatant was harvested and the presence of recombinant virus was initially confirmed by immunoperoxidase staining of virus plaques (Murphy et al., Vaccine 8:497-502 (1990), incorporated herein by reference) with anti-HN monoclonal antibody (Mab) 77/5, which binds to both biologically derived and recombinant

JS PIV3, and Mab 423/6, which does not bind to cDNA-derived virus because its epitope was ablated to serve as a marker. Virus present in passage 1 was subjected to two or three rounds of plaque purification on LLC-MK2 cells as described previously. Each biologically cloned recombinant virus was amplified twice in LLC-MK2 cells at 32°C to produce virus for further characterization. Virus was concentrated from clarified medium by polyethylene glycol precipitation, and viral RNA (vRNA) was extracted with Trizol Reagent (Life Technologies). Reverse transcription (RT) was performed on vRNA using the Superscript II kit with random hexamer primers (Life Technologies). The Advantage cDNA PCR kit (Clontech, Palo Alto, CA) and sense (5' nt 11190-GCATTATCTAGATGTGTCTTCTGGTCAGAG 3' nt-11219) (SEQ ID NO: 31) and antisense (5' nt 14140-CCTGAATTATAATAATTAAGTCAGGTCCT 3' nt-14111) (SEQ ID NO: 32) primers specific for the PIV3 L gene were used to amplify the region spanning the *Sph* I to *Nhe* I fragment. The PCR fragments were analyzed by digestion with each of the restriction enzymes whose recognition sites had been ablated during insertion of the three cp45 mutations in L (see Table 5).

Efficiency of plaque formation (EOP) at permissive and restrictive temperatures of rPIV3 bearing one, two or three cp45 L protein amino acid substitutions.

The level of temperature sensitivity of plaque formation *in vitro* of control and recombinant viruses was determined at 32°C, 37°C, 38°C, 39°C, 40°C, and 41°C in LLC-MK2 monolayer cultures and plaques were enumerated by hemadsorption with guinea pig red blood cells following removal of the methylcellulose overlay. Alternatively, viral plaques present in the monolayer were identified by immunoperoxidase staining with a mixture of two PIV3-specific anti-HN murine mAbs 101/1 and 454/11 diluted 1:500, (Murphy et al., supra, (1990)).

Hamster studies.

4 to 16 week-old golden Syrian hamsters in groups of six were inoculated intranasally with 0.1ml OptiMEM1 per animal containing 10^{5.5} pfu of rPIV3 JS wt, PIV3 cp45 virus, or

one of the rPIV3 containing one or more cp45 L protein substitution(s). On day 4 post-infection, the hamsters were sacrificed, the lungs and nasal turbinates were harvested, and the virus was quantified as described above. The mean log₁₀ TCID₅₀/g was calculated for each group of six hamsters.

RESULTS

Introduction of the PIV3 cp45 L protein amino acid substitutions into wt JS rPIV3.

As noted above, the three amino acid substitutions present in the L protein of cp45 (Table 5) were introduced individually or in selected combinations into the antigenomic cDNA that encodes its wt parent, PIV3 JS strain. Each introduced mutation was engineered to be marked with a silent mutation that ablated a proximal, naturally-occurring restriction enzyme site to facilitate monitoring the mutation in recovered rPIV3 (Table 5, Fig. 12). The coding change at amino acid 1558 was designed to contain two nucleotide changes in r1558, compared to the one nt substitution in cp45, to reduce the chance of reversion at this site during *in vitro* or *in vivo* replication.

Seven rPIV3s bearing one, two or all three of the amino acid substitutions from cp45 were recovered in tissue culture by transfection of each antigenomic cDNA together with the pTM(N), pTM(P) and pTM(L) support plasmids and coinfection with the vaccinia virus MVA/T7 pol recombinant (Wyatt et al., supra, (1995)). Each rPIV3 possessed the Mab resistance marker that had been deliberately introduced into the HN gene by engineering the antigenomic cDNA. The rPIV3s were biologically cloned by two or three cycles of plaque to plaque passage to ensure that each virus preparation was genetically homogeneous. This precaution was taken because vaccinia virus can mediate recombination between the antigenomic cDNA and the support plasmids.

To confirm that each of the seven rPIV3 contained the engineered mutation(s) in the L gene, RNA was purified from precipitated virions and was copied into cDNA and amplified by RT-PCR. Control reactions showed that the RT step was required for generation of RT-PCR products,

indicating that an RNA template rather than contaminating cDNA was required for the generation of the RT-PCR product. The RT-PCR products were subjected to digestion with the three restriction enzymes whose recognition sequences had been
5 ablated as markers for the inserted coding changes. As expected, the RT-PCR product of JS wt rPIV3 was cleaved the appropriate number of times by each of the three enzymes, whereas r942/992/1558 lacked each of the three sites ablated during creation of the individual cp45 coding changes. Each
10 of the other rPIV3s lacked the appropriate restriction site(s), indicating the presence of the introduced mutations.

Efficiency of plaque formation at 32°C, 37°C, 38°C, 39°C, 40°C, and 41°C of rPIV3 bearing cp45 L mutations.

The seven rPIV3s bearing the various combinations
15 of cp45 L protein amino acid substitutions were assayed for their ability to form plaques on LLC-MK2 monolayers at 32°C, 37°C, 38°C, 39°C, 40°C and 41°C. As shown in Table 6, each rPIV3 bearing a cp45 aa substitution was *ts*, whereas the JS wt rPIV3 parent was not restricted in plaque formation at any
20 temperature tested. The shut-off temperature of plaque formation of r942, r992 and r1558 was 40°C. r942 manifested a 700-fold reduction of plaque formation at 40°C, indicating that its replication was marginally reduced at this
restrictive temperature. However, the plaque size of r942
25 also was greatly reduced at 40°C, which also indicates that its replication was restricted at this temperature compared to the JS wt. r942 was completely restricted in replication at 41°C (data not shown). r992 and r1558 were greatly reduced (over a 1,000,000-fold reduction) in plaque formation at 40°C.
30 These results indicate that each of the three cp45 L gene mutations individually specifies the *ts* phenotype, although that of the r942 mutation is somewhat less restrictive. The double mutant virus r942/1558 and the triple mutant r942/992/1558 had a shut-off temperature of 39°C, while that
35 of r942/992 and cp45 was 38°C. The double mutant, r992/1558, was less *ts* than the r992 single mutant. r992/1558, like r942, was completely restricted in plaque formation at 41°C. The level of temperature sensitivity exhibited by double or

triple results from a rPIVs delicate interplay of the three mutations that cannot be predicted from level of temperature-sensitivity exhibited by the single mutants. Also, since r942/992/1558 was slightly less *ts* than cp45, other mutations outside of the L gene also likely contribute to the *ts* phenotype of cp45, therefore representing additional mutations of interest within the invention.

Table 6: The efficiency of plaque formation (EOP) at 32°C, 37°C, 38°C, 39°C and 40°C of rPIV3 bearing one, two or three cp45 L protein amino acid substitutions.

Virus ^a	Virus titer (log ₁₀)				
	pfu/ml)				
	32°C	37°C	38°C	39°C	40°C
r942	6.8	6.8 ^b	6.6 ^b	6.5 ^b	<u>4.3^b</u>
r992	6.9	6.8 ^b	6.7 ^b	6.1 ^b	<0.7
r1558	6.6 ^b	6.6 ^b	6.4 ^b	5.0 ^b	<0.7
r942/992	6.7	6.5 ^b	<u>4.5^b</u>	3.0 ^b	<0.7
r942/1558	5.2	5.0 ^b	4.0 ^b	1.0 ^b	<0.7
r992/1558	6.7	6.7 ^b	6.5 ^b	5.9 ^a	5.1 ^b
r942/992/1558	6.6	6.2 ^b	6.2 ^b	<u>2.7^b</u>	<0.7
cp45 ^b	6.6	4.8	<u>4.5^b</u>	<0.7	<0.7
rPIV3 JS	8.3	8.4	8.5	8.5	8.3

a. The cp45 virus is a biologically derived virus, and each of the other viruses tested is a recombinant.

b. Plaques were of pinpoint size.

c. Underlined numbers represent the shut-off temperature of plaque formation, which is defined as the lowest restrictive temperature at which a 100-fold reduction in titer is observed compared to the titer at 32°C.

Growth in hamsters. Groups of six Golden Syrian hamsters were inoculated intranasally with JS wt rPIV3, biologically-derived cp45, or with rPIV3 containing one or more cp45 L protein amino acid substitutions, and virus replication in the lungs and nasal turbinates was determined. In this experiment [Table 7], each of the rPIV3s bearing a single amino acid substitution was restricted in replication in the upper and lower respiratory tract [Table 7]. However, r942, the least *ts* virus, was only marginally suppressed in

replication in the upper and lower respiratory tract in a second experiment. These data demonstrate that two of the three amino acid substitutions contribute to the att phenotype when present as single-lesioned recombinant viruses. However, the 942 mutation indeed contributes to attenuation (e.g., the r942/992 is more attenuated than r992 alone). Thus, each of the amino acid substitutions in L contribute to the att phenotype either acting alone or in concert with another L amino acid mutation. Each of the double mutants was attenuated indicating that loss of any of the three L gene substitutions following replication in vivo still leaves an attenuated virus. This is a partial explanation for the previously observed high level of stability of the ts phenotype of the cp45 following replication in vivo. The triple mutant r942/992/1558 was as restricted as cp45 for replication in the upper and lower respiratory tract indicating that the three amino acid substitutions in the L protein are the major contributors to the att phenotype of cp45.

Table 7: The level of replication in the upper and lower respiratory tract of hamsters of rPIV3 bearing one, two or three cp45 L protein amino acid substitutions, compared to JS wt rPIV3 and cp45^a.

Virus	Mean virus titer (\log_{10} TCID ₅₀ /g \pm S.E ^b)	
	Nasal turbinates	Lungs
rPIV3 wt	7.4 \pm .16	5.1 \pm .49
r942	6.6 \pm .17	3.0 \pm .78
r992	4.4 \pm .16	3.1 \pm .11
r1558	3.8 \pm .40	4.3 \pm .34
r942/992	<1.5 \pm 0	<1.5 \pm 0
r942/1558	2.9 \pm .23	1.8 \pm .17
r992/1558	5.7 \pm .16	3.2 \pm .57
r942/992/1558	3.9 \pm .15	<1.5 \pm 0
cp45	4.1 \pm .27	1.6 \pm .08

a. Groups of six hamsters each were intranasally administered $10^{5.5}$ pfu of virus per animal in an 0.1ml inoculum, and lungs and nasal turbinates were harvested four days later.

b. Standard Error

c. cp45 is a biologically derived virus and the others are recombinant.

To summarize the above results, substitutions at L protein amino acid positions 992 and 1558 each specified a 1,000,000-fold reduction in plaque formation in cell culture at 40°C, while the substitution at position 942 specified a 700-fold reduction. Thus, each of the three mutations individually contributes to the *ts* phenotype. The triple recombinant which possesses all three L mutations is slightly less *ts* than cp45, suggesting that there are mutations outside of the L gene in cp45 that also might contribute to its *ts* phenotype. Two of the three individual mutations in L each contributed to restricted replication in the upper or lower respiratory tract of hamsters, which accounts for the observed stability of *ts* and *att* phenotypes of cp45 during replication *in vivo*. Importantly, the level of temperature sensitivity of recombinant vaccine strains *in vitro* was closely predictive of attenuation *in vivo*. The recombinant virus possessing all three mutations was as restricted in replication as the cp45 mutant in both the upper and lower respiratory tract of

hamsters, indicating that the L gene of the cp45 virus is a major attenuating component of this candidate vaccine strain. While each mutation on its own specifies the *ts* phenotype, when placed together they are not simply additive but instead somehow influence each other. The effect of the three mutations together in the triple mutant seemed to ameliorate rather than enhance the level of temperature-sensitivity observed in the two double mutants which were evaluated. Interestingly, this should provide an unanticipated selective pressure to maintain at least some of the cp45 L mutations, since the loss by reversion of either the 992 or 1558 substitution would increase rather than decrease the level of temperature sensitivity. Considered together, these findings indicate that the high level of the stability of the *ts* and *att* phenotypes of cp45 virus results from the contribution of multiple *ts* mutations in L to the *att* phenotype. The identification of these three mutations as the major attenuating mutations of cp45 provides the basis for monitoring virus during all stages of manufacture and following replication in humans.

It is of further interest that the tyrosine to histidine mutation at position 942, arguably the most conservative substitution of the three mutations, was the least temperature sensitive. The L polymerase of PIV3 is a large polypeptide, 2233 aa in length, and is thought to be a multifunctional protein that encodes multiple domains including those required for complex formation with the P protein, RNA binding, RNA polyadenylation, RNA transcription and RNA replication (Collins et al., supra, (1996)). The amino acid substitutions in L at positions 942 and 992 are located near regions that are well-conserved among other members of the Paramyxovirus family (Blumberg et al., Virology 164:487-497 (1982); Galinski et al., Virology 165:499-510 (1988)). The mutation at position 1558 is in a region of the polymerase which appears to share less sequence identity with other L polymerases. Although the mechanism by which the *ts* phenotype is conferred by the triple amino acid substitution in L is not known, it is likely that multiple L protein

domains and activities are affected, or that a common mechanism involving various activities of L is affected.

EXAMPLE X

Direct Identification, and Reconstitution into Recombinant Vaccine Viruses, of Mutations in a Biologically Derived, Live-Attenuated HPIV Type 3 Virus (cp45) Which Specify the Temperature-Sensitive, Cold-Adaptation and Attenuation Phenotypes

The above Examples demonstrate that each of the three amino acid substitutions in the L polymerase protein of cp45 confer the temperature-sensitive (*ts*) and attenuation (*att*) phenotypes, but not the cold-adaptation (*ca*) phenotype (see also, Skiadopoulos et al., J. Virol 72(3):1762-8, 1998). cp45 contains twelve additional mutations in other proteins (N, C, M, F and HN) or potential cis-acting sequences (the leader region and the transcription gene start {GS} signal of the N gene), and their contribution to these phenotypes has been heretofore undefined. The present Example further characterizes the genetic basis for the *ts*, *ca*, and *att* phenotypes of cp45 to provide yet additional information regarding basis for the observed high level of stability of these phenotypes following replication of cp45 in humans or non-human primates. In one aspect of this study, a recombinant cp45 (rcp45) virus containing all fifteen cp45-specific mutations was constructed, using a reverse genetics system, and was found to be essentially indistinguishable from the biologically-derived virus on the basis of plaque size, level of temperature-sensitivity, cold-adaptation, and level of replication in the upper and lower respiratory tract of hamsters. In addition, recombinant viruses containing: (1) the cp45-specific changes in the C, M, F or HN proteins, (2) the combined leader and N gene mutations, or (3) several combinations of the cp45 mutations were constructed. Analysis of these recombinant viruses showed that multiple cp45 mutations distributed throughout the genome contribute to the *ts*, *ca*, and *att* phenotypes. The mutations in C and F were not *ts* at 40°C but nonetheless

conferred the att phenotype, and they, therefore, are non-ts att mutations. The HN mutation did not confer the ca, ts or att phenotypes. Viruses possessing the 3' leader and N mutations were ts, but exhibited only marginal attenuation in the lower respiratory tract of hamsters. Recombinants possessing several combinations of mutations exhibited a greater level of temperature sensitivity than cp45, but the increased level of temperature-sensitivity was not reflected in an increase in attenuation *in vivo*. These latter findings indicate that the multiple mutations identified in cp45 are interacting to affect replication *in vitro*. The presence of multiple ts and non-ts attenuating mutations in cp45 likely contributes to its high level of attenuation and phenotypic stability. Knowledge of the phenotypes associated with the various mutations of cp45 provided herein allows for accurate monitoring of biologically derived PIV viruses and ready manipulation of recombinant virus to achieve a large assemblage of useful vaccine recombinants within the invention.

Viruses and cells.

The rPIV3s, PIV3 JS wt and cp45 viruses described in the present Example were grown in simian LLC-MK2 cells (ATCC CCL 7.1) as described above (see also, Durbin et al., Virology 235:323-332, 1997a; Hall et al., Virus Res. 22(3): 173-184, 1992; Skiadopoulos et al., J. Virol 72(3):1762-8, 1998). The modified vaccinia virus Ankara was provided as described above. HEp-2 (ATCC CCL 23) and LLC-MK2 cells were maintained in OptiMEM I (Life Technologies, Gaithersburg, MD) supplemented with 2% FBS and gentamicin sulfate (50ug/mL), or in EMEM supplemented with 10% FBS, gentamicin sulfate (50ug/mL), and 2mM glutamine. L-132 cells (ATCC CCL 5) were grown in Earl's MEM (Life Technologies) supplemented with 10% FBS, 2mM glutamine, 20 mM Hepes, 1 mM non-essential amino acids, and 100 units streptomycin-neomycin/ml.

Construction of point mutations in PIV3.

Subgenomic fragments of p3/7(131)2G, the antigenomic cDNA clone of PIV3 JS wt used above to recover infectious virus (see also, Durbin et al., Virology 235:323-332, 1997a;

Skiadopoulos et al., *J. Virol.* 72(3):1762-8, 1998), were cloned into pUC19 vectors modified to accept these fragments, using standard molecular cloning techniques. Point mutations corresponding to mutations identified in cp45, as well as mutations creating or ablating silent restriction enzyme recognition sequences (Table 8) were introduced using the Transformer Mutagenesis Kit (Clontech, Palo Alto, CA) as described previously.

Table 8: PIV3 cp45 mutations introduced into rPIV3

	seq id no	region affected	nt position ^a	sequence changes ^b	restriction marker ^c	amino acid substitution ^d
1	46 47	3' leader	20	TTGTCTGGGAAT TTGCCTGGGAAT	none	non-coding
2	46 48	3' leader	20	TTGTCTGGGAAT TTGTTTGGGAAT	none	non-coding
3	46 49	3' leader	20	TTGTCTGGGAAT TTGTCTGGTAAT	none	non-coding
4 ^e	50 51	3' leader	40	AACTTTAAATTA AACTTAAATTA	-Dra I	non-coding
5 ^e	52 53	N gene start	60	TTAAAGACATTG TTTAAGACATTG	none	non-coding
6 ^e	54 55	N	390	GCAGATGTCAAG GCAGATGCCAAG	none	Val-96 to Ala
7 ^e	56 57	N	1271	CGAATCTAAAGA CGAAGCTAAAGA	none	Ser-389 to Ala
8	58 59	C	2076	GAAATATTGATC GAAACATTGATC	-Ssp I	Ile-96 to Thr
9	60 61	M	4341	TCTCTACCCAAC TCGTTAACCAAC	+Hpa I	Pro-199 to Thr
10 ^f	62 63	F	6323	AGTACAATAGGT AGTACTGTGGGT	+Sca I	Ile-420 to Val
11	64 65	F	6419	GCACTTGATCCA ACACTGGATCCA	+Bam HI	Ala-450 to Thr
12	66 67	HN	7944	CCATCATTGTTGTTGACAA CCATCATTGTGGCTGACAA	+Bst XI	Val-384 to Ala
13	68 69	L	11468	TTACATGGCCA TCACATGGCGA	-Eae I	Tyr-942 to His
14	70 71	L	11618	TTTGGACTGGGC TTTTGATTGGGC	-Bsr I	Leu-992 to Phe
15 ^f	72 73	L	13308	GGTCCTAATACT GGGCCTAATATC	-Ava II	Thr-1558 to Ile

a. Position of the first nucleotide in the PIV3 sequence shown. b. Wild type sequence is shown above the mutant sequence. Nucleotide changes are underlined. Codon substitutions are in bold font. c. Each relevant restriction endonuclease recognition sequence is in italics; (+) indicates addition of new restriction endonuclease recognition sequence; (-) indicates ablation of a naturally occurring restriction endonuclease recognition sequence. d. Mutations are indicated as the three letter amino acid assignment of wt, followed by the amino acid position, followed by the cp45 assignment. e. These mutations were identified by Joanne Tatem (unpublished observations), the others were from Stokes et al., *Virus Res.* 30(1):43-52, 1993. f. Two nucleotides were changed in the indicated codon in order to reduce the chance of reversion to wild type sequence.

After mutagenesis, restriction endonuclease fragments were sequenced completely and cloned back into the full-length clone, p3/7(131)2G. The 3' leader and N mutations were amplified by reverse transcription (RT)-PCR directly from PIV3 cp45 virion RNA and were cloned into pLeft+2G (see
5 above), or a modified pUC19 vector for further manipulation. Combinations of mutations were constructed using standard molecular cloning techniques. The full-length plasmid clone containing the cp45 mutations, designated pFLCcp45, was
10 completely sequenced to determine if extraneous mutations had been introduced during the cloning process, but none were found.

Recovery of recombinant PIV3 (rPIV3).

Each full-length antigenomic cDNA bearing cp45
15 mutations, together with the three support plasmids pTM(N), pTM(P no C) and pTM(L), was transfected into HEP-2 cells on 6-well plates (Costar, Cambridge, MA) using LipofectACE (Life Technologies, Gaithersburg, MD) and MVA-T7 as described above. After incubation at 32°C for 4 days, the transfection harvest
20 was passaged onto LLC-MK2 cells in T-25 flasks which were incubated at 32°C for four to eight days. This harvested virus was called passage 1 and was subjected to three rounds of plaque purification on LLC-MK2 cells as described above. Each biologically-cloned recombinant virus was amplified twice
25 in LLC-MK2 cells at 32°C to produce virus for further characterization. Virus was concentrated from clarified medium by polyethylene glycol precipitation (see, Mbiguino and Menezes, J. Virol Methods 31:2-3, 1991, incorporated herein by reference in its entirety), and viral RNA (vRNA) was extracted
30 with Trizol Reagent (Life Technologies). Reverse transcription was performed on vRNA using the Superscript II Preamplification System (Life Technologies) with random hexamer primers. The Advantage cDNA PCR kit (Clontech) and sense and antisense primers specific for various portions of
35 the PIV3 genome were used to amplify fragments for restriction endonuclease analysis. The PCR fragments were analyzed by digestion with each of the restriction enzymes whose

recognition sites had been created or ablated during construction of the mutations (Table 8).

Efficiency of plaque formation (EOP) of rPIV3 bearing cp45 mutations at permissive and restrictive temperatures.

The level of temperature-sensitivity of plaque formation *in vitro* of control and recombinant viruses was determined at 32°C, 35°C, 36°C, 37°C, 38°C, 39°C, 40°C, and 41°C in LLC-MK2 monolayer cultures as described above (see also, Hall et al., Virus Res. 22(3):173-184, 1992, incorporated herein by reference in its entirety). Plaques were enumerated by hemadsorption with guinea pig red blood cells following removal of the methylcellulose overlay, or alternatively, the viral plaques present in the monolayer were identified by immunoperoxidase staining with a mixture of two PIV3-specific anti-HN murine mAbs 101/1 and 454/11 diluted 1:250 (see, Murphy et al., Vaccine 8(5):497-502, 1990; van Wyke Coelingh et al., Virology 143(2):569-582, 1985).

Evaluation of rPIV3 mutant viruses for cold-adaptation phenotype.

Growth of mutant and wt rPIV3 viruses was determined at 32°C and 20°C on confluent L-132 cell monolayers prepared in 24-well tissue culture plates. Duplicate wells of each of two plates were inoculated with 0.2 ml of each mutant or wt rPIV3 virus at a multiplicity of infection of 0.01. After one hour adsorption at room temperature, the inoculum was aspirated and the monolayers were washed with 1 ml of PBS per well. The inoculated cultures were overlaid with 0.5 ml of Earl's MEM supplemented with 10% FBS, 2mM glutamine, 20 mM Hepes, 1 mM non-essential amino acids, and 100 units streptomycin-neomycin/ml. One plate was sealed in a waterproof pouch (Kapak) and then submerged in a 20°C bath for 13 days. The duplicate plate was placed at 32°C in a CO₂ incubator for 3 days. At the end of the incubation period, virus was harvested by freeze/thawing. The titer of virus recovered from each well was determined by plaque assay in LLC-MK2 cells at 32°C using hemadsorption with guinea pig red

blood cells to visualize plaques. Two wt and two cp45 reference stocks were used as controls.

Hamster studies.

5 week-old Golden Syrian hamsters in groups of five were inoculated intranasally with 0.1 ml OptiMEM I per animal containing $10^{6.0}$ pfu of JS wt rPIV3, PIV3 cp45 virus, or one of the mutant rPIV3s. On day 4 post-infection, the hamsters were sacrificed, the lungs and nasal turbinates were harvested, and the virus was quantified as described above. The mean \log_{10} TCID₅₀/gram at 32°C was calculated for each group of hamsters.

RESULTS

Introduction of the PIV3 cp45 mutations into JS wt rPIV3.

The fifteen mutations in the 3' leader, the N GS signal, and the N, C, M, F, HN and L proteins of cp45 (Table 8) were introduced into the complete PIV3 antigenomic cDNA by site directed mutagenesis or by direct PCR amplification of a segment of cp45 cDNA bearing the desired mutations. The following antigenomic cDNAs were made (see Fig. 13): (i) rcp45 3'N, containing the four point mutations of the leader region, the point mutation in the N GS signal, and the two amino acid changes in the N protein; (ii) rcp45 C, containing the single amino acid change in the C protein; (iii) rcp45 M, containing the single amino acid change in M; (iv) rcp45 F, containing the two amino acid changes in F; (v) rcp45 HN, containing the single amino acid change in HN; (vi) rcp45 L, containing the three amino acid changes in L, as described above; (vii) rcp45 3'NL, containing the mutations from i. and vi. above; (viii) rcp45 3'NCMFHN, containing all of the mutations except for the three in L; and (ix) rcp45, containing all fifteen cp45 mutations listed in Table 8. In most cases, each cp45 change was engineered to be accompanied by one or more nearby silent changes which introduced or removed a restriction enzyme recognition site (Table 8). These served as markers to confirm the presence of the mutation in the cDNA and in recovered virus. Also, two of the amino acid coding changes (mutation numbers 10 and 15 in Table 8) were made using two

nucleotide changes rather than the single change found in cp45, reducing the possibility of same-site reversion to wt. The cp45 cDNA, which contains all fifteen of the cp45 changes in Table 8, was assembled from the same mutagenized cDNA subclones as were used to introduce cp45 changes into the other antigenomic cDNAs; it was sequenced in its entirety, and was confirmed to contain only the intended mutations. This indicated that all of the regions which had been subjected to mutagenesis or PCR and had been manipulated by cloning possessed the desired sequences and lacked other unwanted mutations. Each full-length plasmid bearing one or more of the cp45 mutations was transfected into HEp-2 cells along with support plasmids and MVA-T7 to produce recombinant PIV3 as described above. Analysis of RT-PCR fragments encompassing the mutations indicated in Table 8 were amplified from virion RNA of the various recombinant viruses indicated in Figure 13 confirmed the presence of the introduced mutations, and other unintended mutations were not found.

Plaque morphology.

Several of the recombinant viruses exhibited distinctive plaque morphologies when tested on LLC-MK2 cells. JS wt rPIV3 plaques averaged 1mm in size, and were indistinguishable in size from the biologically derived JS wt PIV3. Plaques of the cp45 and rcp45 viruses were larger than wt, averaging two- to three-fold larger in diameter than wt, and were indistinguishable from each other. This demonstrated the comparability of the biologically-derived and recombinant cp45 virus for this phenotype.

Efficiency of plaque formation of rPIV3s bearing the cp45 mutations in LLC-MK2 cells at permissive and restrictive temperatures.

The rPIV3s were assayed for their ability to form plaques at permissive and restrictive temperatures ranging from 32°C to 41°C (Table 9). Analysis of the ts phenotypes of viruses bearing individual components of cp45 revealed that the rcp45 3'N and rcp45 M viruses had a shutoff temperature of 40°C, and the rcp45 C mutant had a shutoff of 41°C. The shutoff temperature of rcp45 F and rcp45 HN mutants was

greater than 41°C. Consistent with the above results, the rcp45 L virus had a shutoff temperature of 39°C. A virus is considered to have a *ts* phenotype, for example, if its reduction of replication at 40°C (ie. titer at 32°C minus titer at 40°C) is approximately \geq 100-fold that of wt virus at 40°C. Applying this definition, the present results indicated that at least two regions of cp45 (3'N, and L) contribute to the *ts* phenotype.

Table 9: The efficiency of plaque formation (EOP) of recombinant and biologically-derived viruses at permissive and restrictive temperatures.

Virus	Mean Virus Titer ^a (log ₁₀ pfu/ml)								ts phenotype
	32°C	35°C	36°C	37°C	38°C	39°C	40°C	41°C	
rcp45 3'N	7.1	---	---	7.0	6.4	5.4	4.2	<3.7	+
rcp45 C	6.9	---	---	7.0	6.7	6.6	5.9	<3.7	-
rcp45 M	7.7	---	---	7.4	7.0	6.5	5.3	<3.7	-
rcp45 F	7.5	---	---	7.2	6.0	6.6	5.9	5.7	-
rcp45 HN	6.4	---	---	6.5	6.2	6.4	4.7	4.4	-
rcp45 L	7.3	---	---	7.2	6.7	4.0	<0.7	<0.7	+
rcp45 3'NL	7.3	5.7	<0.7	<0.7	<0.7	<0.7	<0.7	---	+
rcp45 3'NCMFHN	7.2	5.6	<0.7	2.0	2.4	<0.7	<0.7	---	+
rcp45	8.5	7.5	7.1	6.4	6.0	2.0	<0.7	<0.7	+
cp45 ^b	8.3	8.0	7.4	7.0	6.2	2.3	<0.7	<0.7	+
rPIV3 wt	7.3	7.3	7.0	7.4	7.6	7.7	6.8	6.0	-

a. Plaques were enumerated by immunoperoxidase staining after incubation for 6 days at the indicated temperature. Values are the mean of two experiments, values in bold are from a single experiment. Underlined values represent the lowest non-permissive temperature at which a 100-fold reduction of plaque efficiency occurs compared to titer at 32°C, and this is defined as the shutoff temperature of plaque formation. b. cp45 is biologically-derived and the other viruses are recombinants. c. A virus is defined as bearing the ts phenotype if its reduction of replication at 40°C (ie. titer at 32°C minus titer at 40°C) is \geq 100-fold that of wt virus at 40°C.

As shown in Table 9, *rcp45*, containing all of the *cp45* mutations, had a shut-off temperature of 38°C, which was identical to that of the biologically-derived *cp45*. These results show that the *ts* phenotype of *cp45* was successfully reproduced in *rcp45*. In addition, these results validate the sequence analysis of *cp45* and the subsequent reconstruction of mutations into recombinant virus.

The *rcp45* 3'NCMFHN virus, which is identical to *rcp45* except that it lacks the three L mutations, exhibited a shutoff temperature of 36°C. Since the L mutations are known to confer temperature-sensitivity individually and in combination, it is somewhat paradoxical that *rcp45* 3'NCMFHN was more, rather than less, *ts* than *cp45*. This implies that there is an interaction of mutations within *cp45* whereby mutations compensate for each other to give a level of temperature-sensitivity which is less than the sum of the individual components.

Virus *rcp45* 3'NL was constructed to investigate whether the L mutations interact with the leader and/or N mutations, since all of these elements are believed to interact during RNA synthesis. This virus had a shutoff temperature 36°C, compared to 40°C and 39°C for *rcp45* 3'N and *rcp45* L, respectively. These results suggest that there is an interaction between the 3'N and L mutations that results in augmentation of temperature-sensitivity. These results also provide another example in which a subset of *cp45* mutations specifies a level of temperature sensitivity greater than that observed for *rcp45* which contains the entire set of mutations

***ca* phenotype of rPIV3s bearing *cp45* mutations.**

The rPIV3s were analyzed to determine which genetic elements of *cp45* specified the *ca* phenotype (Table 10). It was demonstrated above that *rcp45* L is *ts* and *att*, but not *ca*. This indicates that the genetic elements specifying the greater part of the *ca* phenotype are located outside L, and this was confirmed in the present study. Each of the rPIVs possessing the 3' leader and N mutations were *ca* except *rcp45*

3'NCMFHN, which exhibited an intermediate phenotype. This shows that the *ca* phenotype is specified mostly within the 3'N region. The finding that the level of *ca* of viruses containing the 3'N segment is less than that of *cp45* or *rcp45* indicates that other regions of *cp45* contribute to the *ca* phenotype, even though this is not apparent from analysis of the other regions individually. The *rcp45* 3'NL virus is more *ca* than the *rcp45* 3'N virus, suggesting that the L mutations may make a modest contribution. The biologically-derived *cp45* and *rcp45* viruses exhibit comparable levels of *ca*, indicating that this phenotype, like the plaque size and *ts* phenotypes, was successfully reproduced in the recombinant *cp45* virus provided herein. Therefore, the *ca* phenotype, like the *ts* phenotype, is a composite phenotype reflecting individual contributions to the overall phenotype as well as contributions from interacting genetic elements.

Table 10 : Growth of wt and mutant PIV3s at 20°C compared to 32°C^a (cold adaptation phenotype).

Virus	20°C	32°C	<i>ca</i> Phenotype
<i>cp45</i> ^b	6.72	8.49	+
<i>rcp45</i>	6.57	8.35	+
<i>rcp45</i> 3'NL	5.02	7.23	+
<i>rcp45</i> 3'N	4.39	8.53	+
<i>rcp45</i> 3'NCMFHN	3.53	8.08	+/-
<i>rcp45</i> L	3.18	7.98	-
<i>rcp45</i> C	2.52	8.32	-
<i>rcp45</i> F	2.47	8.10	-
<i>rcp45</i> HN	2.11	8.01	-
<i>rcp45</i> M	1.76	8.21	-
PIV3 WT	2.57	8.43	-

a. Virus titer is expressed in log₁₀ PFU/mL. b. *cp45* is biologically-derived and the other viruses are recombinant. The *ca* phenotype is defined as a greater than 10-fold increase in growth at 20°C relative to wt.

Growth of the *rcp45* mutant viruses in hamsters.

Groups of five Golden Syrian hamsters were inoculated intranasally with 10⁶ TCID₅₀ of recombinant or

biologically-derived virus, and the level of virus replication in the lungs and nasal turbinates was determined four days later (Table 11). The fourth day post-inoculation has been shown previously to be the peak of virus replication in hamsters for both the wt and cp45 viruses (see, Crookshanks and Belshe, J. Med Virol 13(3):243-9, 1984, incorporated herein by reference in its entirety). The rcp45 virus was reduced approximately 40-fold in replication in the nasal turbinates and 1000-fold in the lungs, and thus was as attenuated as the biologically-derived cp45 virus. These results indicate that the attenuation phenotype of cp45 was successfully reproduced in its recombinant version. Since each phenotype of cp45 was fully reproduced in rcp45, the additional five mutations in cp45 that were not included in this Example likely made little contribution to the properties of cp45.

Table 11: Level of replication in the upper and lower respiratory tract of hamsters^a of wt and mutant PIV3s^b.

Virus	Mean virus titer (\log_{10} TCID ₅₀ /g \pm S.E ^c) in:	
	Nasal turbinates	Lungs
JS wt rPIV3	6.9 \pm 0.2	5.4 \pm 0.5
rcp45 3'N	6.5 \pm 0.2	3.9 \pm 1.1
rcp45 C	4.8 \pm 0.3	3.1 \pm 0.7
rcp45 M ^d	6.8 \pm 0.2	6.7 \pm 0.3
rcp45 F	4.6 \pm 0.2	3.4 \pm 0.6
rcp45 HN	6.3 \pm 0.2	5.3 \pm 1.0
rcp45L	4.2 \pm 0.1	2.1 \pm 0.3
rcp45 3'NL	4.7 \pm 0.2	2.1 \pm 0.3
rcp45 3'NCMFHN	5.8 \pm 0.3	2.9 \pm 0.9
rcp45	5.3 \pm 0.1	2.4 \pm 0.2
cp45	4.9 \pm 0.4	1.9 \pm 0.2

a. Groups of five hamsters each were intranasally administered 10^{6.0} TCID₅₀ of virus per animal in an 0.1ml inoculum, and lungs and nasal turbinates were harvested four days later. b. cp45 is a biologically-derived virus, the other viruses are recombinant. c. TCID₅₀, 50% tissue infectious dose \pm Standard Error. d. The virus pool used in this study was found to contain a mixed plaque phenotype. The attenuation phenotype of this mutant will be reassessed using additional virus preparations.

As demonstrated above, the mutations in the L gene of cp45 specify the majority of the attenuation phenotype of this virus. In the present Example, the contribution of the cp45 mutations outside of L as a group was examined. The rcp45 3'NCMFHN mutant was only slightly reduced in replication in the nasal turbinates, but was more than 100-fold reduced in replication in the lungs, which shows that additional attenuating mutations were present outside of the L protein. Importantly, if each of the three mutations in the L gene of rcp45 reverted to wild type sequence, the resulting virus, rcp45 3'NCMFHN, would still retain the attenuation phenotype. The rcp45 M and rcp45 HN mutant viruses were not defective for replication in the respiratory tract of hamsters, and the rcp45 3'N virus showed only a marginal decrease in replication

in the lower respiratory tract. This suggests that the mutations present in the 3' leader, in the N gene start site and the N, M and HN proteins are not attenuating in and of themselves. However, these mutations could contribute to the overall attenuation of cp45 in the context of the other cp45 mutations. Also, individual mutations within the 3'N region may have effects which are not apparent when the set of mutations is analyzed together, which can be readily determined according to the present disclosure.

Replication of the rcp45 C and rcp45 F mutant viruses was approximately 100-fold reduced in both the nasal turbinates and the lungs, demonstrating that the mutations present in the C and F proteins of cp45 confer the attenuation phenotype in hamsters, although the level of attenuation is not as great as that conferred by the cp45 L mutations. As described above, the rcp45 F and rcp45 C mutant viruses did not possess the ts phenotype, and therefore, the mutations that occur in the C and F proteins are considered to be non-ts attenuating mutations.

EXAMPLE XI

Recovery of Recombinant, Chimeric PIV3 in which the Hemagglutinin and Fusion Glycoproteins have been Substituted with

Corresponding Glycoproteins From PIV1

Within the present example, a chimeric rPIV virus is generated and selected which incorporates one or more heterologous genes or large polynucleotide sequences from one PIV into a different rPIV background. Within this aspect of the invention, individual genes or gene segments of one PIV are substituted by counterpart sequence(s) from a heterologous PIV or other source. In one embodiment described in the present Example, tools and methods are provided for substituting, e.g., the HN and/or F protective antigens of HPIV1 or HPIV2 into a recombinant HPIV3 to yield a chimeric recombinant suitable for developing live-attenuated vaccines.

Viruses and cells.

The PIV1 strain used in this study, PIV1/Washington/20993/1964 (PIV1/Wash64), was isolated in Washington DC in 1964 and was confirmed to be a virulent wild type virus in clinical studies in adult human volunteers (Murphy et al. Infect. Immun. 12:62-8 (1975), incorporated herein by reference). It was propagated in LLC-MK2 cells (ATCC CCL 7.1) in Opti-MEM I (Life Technologies) with 50 µg/ml gentamicin sulfate, 2 mM glutamine and 0.75 µg/ml trypsin (Catalog No. 3741, Worthington Biochemical Corp., Freehold, NJ). The Greer strain of human PIV2 (Catalog No. V-322-001-020, NIAID Repository, Rockville, MD) used in the hemagglutination-inhibition assay (HAI) was propagated in the same way. The JS strain of human PIV3 virus and its recombinant derivative from cDNA (rPIV3/JS) with wild type phenotype were propagated as described above. The modified vaccinia Ankara (MVA) recombinant that expresses the bacteriophage T7 RNA polymerase is described in Wyatt et al., Virology 210:202-205 (1995) (incorporated herein by reference).

HEp-2 cells were obtained from ATCC (ATCC CCL 23) and maintained in Opti-MEM I (Life Technologies) with 2% fetal bovine serum (FBS), 50 µg/ml gentamicin sulfate and 2 mM glutamine.

Construction of a cDNA encoding a complete chimeric PIV3-PIV1 antigenome.

A cDNA encoding a full-length PIV3 antigenomic RNA in which the PIV3 HN and F ORFs were replaced by their PIV1 counterparts was constructed as shown in Figure 14. cDNA clones of the HN and F genes of PIV1/Wash64 were prepared from RNA extracted from LLC-MK2 cells which had been infected with PIV1/Wash64 wild type virus. cDNA was generated using the SuperScript Preamplification System using random hexamer primers (Life Technologies). PIV1 F and HN cDNAs were amplified with Vent DNA polymerase (New England Biolabs, Beverly, MA) using gene specific primer pairs based on consensus sequences present in GenBank. All primers described below are annotated so that PIV specific sequences are

underlined, restriction sites are in italics, nt altered from wild type sequences are in lowercase, and start and stop codons are in bold. The positive sense PIV1 F primer, hybridizing to nt 69-97 upstream of the start codon, was

5 5'-GGGAAAGAAATCCAGAGACAAGAACGG-3' (SEQ ID NO: 33). The negative- sense PIV1 F primer, hybridizing to nt 36-61 downstream of the F stop codon, was

5'-GGTGAAGTTGTGGATccATTTGATTG-3' (SEQ ID NO: 34). It carries a BamH I site. The positive-sense primer for PIV1 HN was 5'-

10 CAACCTGTAAGGtAcCAGCATCCG-3' (SEQ ID NO: 35). It hybridizes to nt 13-36 upstream the HN start codon and carries a Kpn I site. The negative-sense PIV1 HN primer was

5'-GATATGGTGTTaGGcCTTGATCTGTTC-3' (SEQ ID NO: 36). It hybridizes to the last two nt of the stop codon and 25 nt

15 further downstream and carries a Stu I site. The PIV1 F cDNA was cloned as a BamH I and blunt-end fragment into BamH I-EcoR V digested pLITMUS28 (New England Biolabs), while the PIV1 HN cDNA was cloned as a Kpn I-Stu I fragment into the same vector. The nt sequences of the resulting plasmids,

20 designated as pLit.1HNwt and pLit.1Fwt (GenBank Accession number: AF016280, AF016279), were determined using the Circumvent Sequencing Kit (New England Biolabs). These two clones were modified (Fig. 14) using mutagenic PCR primers to delete their non-coding regions and to introduce new

25 restriction sites flanking their start and stop codons for the purpose of constructing the PIV3-1 chimeric HN and F genes. The sequences of positive-sense and negative-sense PIV1 F mutagenic primers were 5'-Cgcc**ATGg**AAAAATCAGAGATCCTCTTCT-3' (SEQ ID NO: 37) and 5'-Ctggatc**ct**AATTGGAGTTGTTACCCATGTA-3' (SEQ

30 ID NO: 38), respectively, the introduced restriction sites are Nco I and BamH I. The sequences of positive-sense and negative sense PIV1 HN mutagenic primers were 5'-AACc**ATGGCTG**AAAAAGGGAAAA-3' (SEQ ID NO: 39) and 5'-GGTGAaGCT**t**AAGATGTGATTTTACATATTTTA-3' (SEQ ID NO: 40),

35 respectively, the introduced restriction sites are Nco I and Hind III.

The second step involved the modification of the PIV3 HN and F genes to serve as acceptors of the PIV1 coding

regions generated above. The PIV3 F and HN genes were subcloned in several steps from the full-length clone of PIV3/JS, p3/7(131)2G to generate pLit.PIV3.HN.4 and pLit.PIV3.F.3a (Fig. 14). The PIV3 F or HN coding regions were deleted by PCR mutagenesis and replaced by restriction sites appropriate to accept the PIV1 HN and F cDNA described above. The sequences of the PIV3 F positive sense and negative sense mutagenic primers were

5'-AAATAggatccCTACAGATCATTAGATATTTAAAT-3' (SEQ ID NO: 41) and 5'-CgcCATgGTGTTTCAGTGCTTGTG-3' (SEQ ID NO: 42), respectively, the introduced restriction sites were *Bam*H I and *Nco* I. The sequences of PIV3 HN positive sense and negative sense mutagenic primers were 5'-CCACAAGCtTAATTAACCATAATATGCATCA-3' (SEQ ID NO: 43) and 5'-TTCCATggATTTGGATTGTCTATTGGGT-3' (SEQ ID NO: 44), respectively, the introduced restriction sites were *Hind* III and *Nco* I. The stop codon for the chimeric HN would be regenerated upon ligation with mutagenized PIV1 HN cassette. The PIV1 HN or F cDNAs described above were imported in as a *Nco* I-*Hind* III fragment for HN or as a *Nco* I-*Bam*H I fragment for F, which generated pLit.PIV3-1.HNhc and pLit.PIV3-1.Fhc. These two cDNAs were then joined into pSE.PIV3-1.hc, which was subsequently sequenced in its entirety. Then the *Bsp*E I-*Sph* I fragment was inserted into p3/7(131)2G to generate pFLC.2G+.hc (Fig. 14). The nucleotide sequence of this *Bsp*E I-*Sph* I fragment, containing the chimeric F and HN genes, is in GenBank (Accession No. AF016281). Importantly, the cDNA engineering was designed so that the final rPIV3-1 antigenome conformed to the rule of six (Durbin et al. *Virology* 234:74-78 (1997), incorporated herein by reference). The plasmid pFLC.2G+.hc encoding the rPIV3-1 chimera was deposited on 9/17/97 under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A.

Transfection. HEP-2 cell monolayers were grown to confluence in six-well plates (Costar Corp, Cambridge, MA) and transfections were performed as described above. Cell monolayers were infected with MVA-T7 at a multiplicity of

infection (MOI) of 3 in 0.8 ml of serum-free Opti-MEM I containing 50 µg/ml gentamicin, and 2 mM glutamine and then transfected with the three support plasmids, 0.4 µg pTM(N), 0.4 µg pTM(P), 0.05 µg pTM(L), and 5 µg the PIV3-1 antigenome cDNA. The plasmids were mixed in 0.2 ml of Opti-MEM I containing 12 µl LipofectACE (Life Technologies) and added to each well. The PIV3/JS cDNA plasmid, p3/7(131)2G, which encodes wild type PIV3/JS antigenomic RNA, was transfected in parallel as a positive control. After incubation at 32°C for 12 hours, the transfection medium was replaced with 1.5 ml of fresh Opti-MEM I supplemented with 50 µg/ml gentamicin, and cultures were incubated at 32°C for two additional days. Trypsin was added to a final concentration of 0.75 µg/ml on day 3 post transfection. Cell culture supernatants were harvested on day 4 and passaged (referred to as passage 1) onto fresh LLC-MK2 cell monolayers. After overnight adsorption, the medium was replaced with fresh Opti-MEM I with 0.75 µg/ml trypsin. Passage 1 cultures were incubated at 32°C for 4 days, and the amplified virus was harvested and further passaged on LLC-MK2 cells (referred to as passage 2) for another 4 days at 32°C in the presence of 0.75 µg/ml trypsin. The recovered virus was characterized by reaction with rabbit PIV1 antiserum, guinea pig PIV2 antiserum, and two PIV3 HN monoclonal antibodies in a hemagglutination-inhibition (HAI) assay as described previously (van Wyke Coelingh et al., supra, (1985)) with the exception that chicken red blood cells (RBCS) were used for viruses possessing the PIV1 HN glycoproteins, whereas guinea pig RBCs were used for PIV3. The HAI was performed to determine if rPIV3 possessed the monoclonal antibody resistant mutation (MARM) that marked virus recovered from cDNA or if rPIV3-1 possesses the PIV1 HN.

Nucleotide sequence analysis. The chimeric cDNA construct pSE.PIV3-1.hc was sequenced with the Circumvent Sequencing kit (New England Biolabs) before the final assembly into full-length clone pFLC.2G+.hc. For analysis of RNA, the appropriate PIVs were amplified in T75 flasks of LLC-MK2 cells. Virus was harvested on day 5 post-infection and concentrated by polyethylene glycol precipitation (Mbiguino et

al., J. Virol. Methods 31:161-70 (1991)). Viral RNA was extracted from the viral pellets and used in reverse transcription (RT) with the Superscript Preamplification System (Life Technologies). RT-PCR was performed using the Advantage cDNA synthesis kit (Clontech) and PIV1 or PIV3 specific primer pairs. RT-PCR products were gel purified by electrophoresis onto, and elution from, strips of NA45 DEAE nitrocellulose membrane (Schleicher & Schuell, Keene, NH), and were sequenced.

Replication of PIVs in LLC-MK2 cells. Plaque enumeration on LLC-MK2 monolayers was performed as previously described except that trypsin was added in the case of the PIV1 and rPIV3-1 viruses (Hall et al. Virus Res. 22:173-84 (1992)). Briefly, serially-diluted virus was inoculated onto LLC-MK2 monolayers in six-well plates; virus was adsorbed for 1 hour at 32°C; the cultures were overlaid with L-15 medium (Life Technologies) containing 0.8% agarose with or without added trypsin and incubated at 32°C for 6 days; and plaques were identified by hemadsorption (HAD) with guinea pig red blood cells following the removal of the agarose overlay.

Growth of the PIV viruses in tissue culture was evaluated by infecting confluent LLC-MK2 monolayers on twelve-well plates with virus at a MOI of 0.01. After adsorption at 32°C for 1 hour, the inoculum was replaced with 1.5 ml/well Opti-MEM I supplemented with gentamicin (50 µg/ml) and trypsin (0.75 µg/ml), and further incubated at 32°C for 6 days. At 24 hour intervals, 0.3 ml medium was removed from each well and 0.3 ml fresh medium with trypsin was added back. The titer of virus was determined at 32°C by hemadsorption assay on LLC-MK2 cell monolayers using fluid overlay as previously described (Hall et al. Virus Res. 22:173-84 (1992)), and the titers were expressed as log₁₀TCID₅₀/ml.

Replication of PIVs in the respiratory tract of hamsters. Golden Syrian hamsters in groups of 12 were each inoculated intranasally with 0.1 ml of L-15 medium containing 10⁵ pfu of rPIV3/JS, rPIV3-1, or PIV1/Wash64. On days 4 and 5 post-inoculation, six hamsters from each group were sacrificed, and their lungs and nasal turbinates harvested,

and homogenized, and virus present in the samples was titered on LLC-MK2 cell monolayers at 32°C. The titers were expressed as mean log₁₀ TCID₅₀/g for each group of six hamsters.

5

RESULTS

Construction of a cDNA clone encoding a full-length, chimeric PIV3-1 antigenomic RNA yielded recovery of the Chimeric Virus rPIV3-1.

As noted above, the final construct of the PIV3 and PIV1 chimeric cDNA, in which the ORFs of the JS wild type PIV3 HN and F glycoprotein genes were replaced by those of PIV1/Wash64 coding sequences (Fig. 14) encodes a PIV3-1 chimeric antigenomic RNA of 15,516 nt, which conforms to the rule of six (Durbin et al., Virology 234:74-78 (1997)). The pFLC.2G+.hc cDNA encoding the chimeric PIV3-1 antigenome was transfected onto HEp-2 cells together with the N, P and L support plasmids. The p3/7(131)2G cDNA encoding the JS wt PIV3 antigenome was transfected in parallel to generate a rPIV3 control parental virus. Virus was recovered from each transfection following the second amplification on LLC-MK2 cells, and studies were initiated to confirm that each recombinant virus was derived from cDNA.

Recombinant viruses rPIV3-1 and rPIV3 were first characterized for the presence of the PIV1 or PIV3 HN glycoprotein by HAI assay with serotype-specific anti-HN monoclonal or polyclonal antibodies. As shown in Table 12, rPIV3 reacted with only one of the two PIV3 mAbs as expected, whereas its biologically derived parent PIV3/JS reacted with both. This confirmed that rPIV3 contained the introduced MARM mutation that marks this virus as being derived from cDNA. The rPIV3-1 virus reacted with antibodies specific to the PIV1 HN glycoprotein, but not to ones specific to HN of PIV3 or PIV2, showing that the virus contained the PIV1 HN gene as expected.

Table 12. rPIV3-1 possesses the HN glycoprotein of PIV1

Virus	Hemagglutination-inhibition titer ^a (reciprocal) of indicated monoclonal antibody or polyclonal antiserum			
	PIV1 ^b antiserum	PIV2 ^b antiserum	α -PIV3 mAb 423/6 ^c	α -PIV3 mAb 77/5 ^c
PIV1/Wash64	256	32 ^d	≤ 50	≤ 50
rPIV3-1	64	≤ 2	≤ 50	≤ 50
rPIV3/JS	4	≤ 2	≤ 50	3,200
PIV3/JS	8	≤ 2	12,800	6,400
PIV2/Greer	8	512	≤ 50	≤ 50

^aChick red blood cells (RBC) were used in HAI assay for PIV1, PIV2, and rPIV3-1 and guinea pig RBCs were used for PIV3/JS and rPIV3/JS.

^bPIV1 rabbit antiserum was purchased from Denka Seiken Co. Ltd., Japan (Catalog No. 410-701), and PIV2 guinea pig antiserum was obtained from NIAID repository, Rockville, MD (Catalog No. V-322-50-558).

^cBiologically derived PIV3/JS contains epitopes recognized by both mAb 423/6 and 77/5, whereas rPIV3/JS was engineered to lack reactivity with mAb 423/6.

^dThe PIV2 antiserum had some reactivity with PIV1 virus, and therefore is not completely type specific.

It was next confirmed that the rPIV3-1 virus contained the engineered, chimeric PIV3-1 HN and F genes. As designed, the genetic structure of rPIV3-1 was unique in four junction regions when compared with either of its parents, PIV1/Wash64 or rPIV3/JS (boxed in Figure 15A). These regions are the transition points at which the sequence switches from the PIV3 non-coding region to the PIV1 coding region and then from the PIV1 coding region back to the PIV3 non-coding region. Using the primer pair A specific to PIV3 M and L genes, or primer pair B specific to the PIV1 M and the very 3'-end of HN gene, RT-PCR products were generated for virion-derived RNAs from rPIV3-1, rPIV3/JS, and PIV1/Wash64. Control reactions showed that the RT step was required for generation of RT-PCR products, indicating that an RNA template, rather than contaminating DNA, was required to produce the RT-PCR product. An early indication that rPIV3-1

was indeed a chimeric virus came from the finding that only the PIV3-specific primer pair A generated the expected 4.6 kb cDNA product that spans the F and HN genes (Fig. 15B). Thus, while rPIV3-1 virus contains only HPIV1-specific HN glycoprotein (See Table 12), the non-coding regions are specific to PIV3. Conversely, the PIV1 specific primer pair B amplified an appropriately sized product from PIV1 control but not from rPIV3-1. Restriction digestion analysis also demonstrated that rPIV3-1 RT-PCR product had unique restriction patterns different from that of rPIV3/JS and PIV1/Wash64 and appropriate for its predicted sequence.

The nt sequence of the 4.6 kb RT-PCR product of rPIV3-1 was determined in its four regions (Fig. 15 A) and compared with that of rPIV3/JS and PIV1/Wash64 (Fig. 16). The rPIV3-1 sequence was completely in agreement with the cDNA from which it was derived. Examination of the sequence alignment of the Region I-IV for the three RT-PCR products illustrates that rPIV3-1 contains the PIV1 F and HN glycoprotein ORFs with altered start and stop codons and flanked by the 5' and 3' non-coding regions of PIV3. Examples of sequencing ladders spanning the Region III and IV of rPIV3-1 (Fig. 17), compared in parallel with rPIV3/JS or PIV1/Wash64, were evaluated, and this analysis confirmed that rPIV3-1 is a recombinant chimeric virus whose structure is completely in agreement with the cDNA from which it was generated.

Trypsin-dependence and cytopathicity of rPIV3-1 in vitro.

PIV1, like Sendai virus but contrary to PIV3, requires trypsin for cleavage of its F glycoprotein in order to undergo multicycle replication on continuous lines of tissue culture cells (Frank et al. J. Clin. Microbiol. 10:32-6 (1979)). In addition, PIV1 is a non-cytopathic virus whereas PIV3 readily produces extensive CPE (Collins et al. In Fields Virology, 3rd ed., 1:1205-43 (1996)). rPIV3-1, rPIV3 and PIV1/Wash64 were compared on the basis of these properties. rPIV3-1, like PIV1/Wash64, had a higher HA titer using chicken, rather than guinea pig (Table 13), RBCs. rPIV3-1,

like its PIV1/Wash64 parent, required trypsin for efficient replication in cultures with fluid overlay as well as for efficient plaque formation. rPIV3-1 produced plaques at 32°C, 37°C or 40°C with similar efficiency. It is therefore evident that rPIV3-1 possesses the F glycoprotein of the PIV1 parent virus, and it is not temperature sensitive. On the other hand, rPIV3-1 produced CPE, as indicated by the cell rounding and detaching in the virus infected monolayers, almost to the same extent as its PIV3 parent suggesting that this biological property is a function of its PIV3 genes, which lie outside of the HN and F coding regions. Thus, rPIV3-1 possesses biological properties from both parents which is consistent with the findings above demonstrating that it is a chimeric virus. This exemplary recombinant, chimeric virus within the invention was deposited on May 21, 1998 under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC) of 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A.

Table 13. Comparison of the HA titer and the infectivity and cytopathicity of parental and chimeric PIVs^a

5		HA titer using indicated RBC		Infectious titer ^b (Log ₁₀ TCID ₅₀ /ml) using CPE or HAD as endpoint				PFU/ml ^c (Log ₁₀)	
10	Virus								
		Chicken	Guinea Pig	CPE		HAD		Trypsin	
				Trypsin		Trypsin			
15				-	+	-	+	-	+
	PIV1/Wash64	16	8	≤2.5	≤2.5	4.8 ^d	6.3	≤0.7 ^e	5.8
20	rPIV3-1	64	16	≤2.5	5.8	5.5 ^d	7.8	≤0.7 ^e	7.1
	rPIV3/JS	0	8	4.5	7.3	5.0 ^d	7.5	5.0	6.2

^aVirus stocks were grown in LLC-MK2 cells which were infected at an MOI of 0.01 and incubated for 6 days in the presence (PIV1/Wash64, rPIV3-1) or absence (rPIV3/JS) of 0.75 µg/ml trypsin. The resulting virus stocks were assayed by the tests below in the presence or absence of trypsin as indicated.

^bThe TCID₅₀ assay was read at 6 days by direct visualization of CPE or by hemadsorption (HAD).

^cPlaques were visualized by HAD after six days of incubation.

^dThe HAD of PIV3-infected monolayers was grossly apparent whereas that of PIV1 and rPIV3-1 was observable only under the microscope in which single cells with RBC adsorbed were observed.

^eThe lowest level of virus detectable was 10^{0.7}/ml.

Comparison of the level of replication of rPIV3-1 and its parental viruses in LLC-MK2 cells and hamsters.

The multicycle replication of rPIV3, rPIV3-1, and PIV1 Wash/64 viruses was evaluated following inoculation of LLC-MK2 tissue culture cells at a MOI of 0.01 (Fig. 18). It can be seen that the kinetics and magnitude of replication of the three viruses are very similar. This indicates that the substitution of the HN and F genes of PIV1 for those of PIV3 did not attenuate the virus for replication *in vitro*. It was next determined whether rPIV3-1 was attenuated *in vivo*, specifically for replication in the upper and lower respiratory tracts of hamsters (TABLE 14). The observed level of replication of rPIV3-1 was similar to, if not slightly higher than, either parent in the upper and lower respiratory tract of hamsters.

Table 14. Level of replication of parental and chimeric PIVs in the upper and lower respiratory tract of hamsters^a

Virus	Virus titer (mean log ₁₀ TCID ₅₀ /gram±S.E.) in indicated tissue			
	Nasal Turbinates		Lungs	
	Day 4	Day 5	Day 4	Day 5
PIV1/Wash64	5.2±.24	5.2±.12	5.0±.31	5.0±.38
rPIV3-1	4.9±.23	6.2±.17	5.8±.15	6.0±.09
rPIV3/JS	4.5±.09	5.0±.18	5.1±.26	5.0±.32

^aHamsters were infected intranasally with 10^{5.0} TCID₅₀ per animal of the indicated virus, and lungs and nasal turbinates were removed on day 4 or 5 after infection. The titers are means of six animals per day and are expressed as mean log₁₀TCID₅₀/gram ± standard error.

In summary, the present Example demonstrates successful recovery of a rPIV3-1 chimeric virus in which the ORFs of the PIV1 HN and F glycoproteins were substituted for those of rPIV3. This chimeric virus replicated like its wild type PIV1 and PIV3 parental viruses *in vitro* and *in vivo*, demonstrating that the substitution of the glycoprotein ORFs did not result in attenuation of rPIV3-1. This successful recovery of a recombinant PIV3 which bears the HN and F glycoproteins of PIV1 is surprising because the two viruses, representing distinct serotypes, are not closely related. In particular, it is remarkable that the chimeric recombinant grows as well as the two parents. Notably, chimeric recombinant viruses possessing a substitution in the glycoprotein gene have also been recovered for vesicular stomatitis virus (VSV) (Lawson et al. Proc. Natl. Acad. Sci. USA 92:4477-81 (1995)). Specifically, the VSV G glycoprotein gene of Indiana serotype was replaced by that from New Jersey serotype which share only 50% amino acid sequence identity. In contrast to rPIV3-1, the chimeric recombinant VSV_{I/NJ} replicates to only 10% the level of recombinant VSV_I or biologically derived, VSV.

In the present Example, the HN and F glycoproteins have 43 and 47% sequence identity, respectively, between PIV1

and PIV3. The transfer of the two glycoproteins together would, of course, obviate glycoprotein-to-glycoprotein incompatibility (Tanabayashi, K. and Compans, R.W. J. Virol. 70:6112-18 (1996)). On the other hand, it is generally thought that the glycoproteins interact with the M protein (which is 63% identical between PIV1 and PIV3) through their cytoplasmic (CT) or transmembrane (TM) domains, and that this interaction is important in virion morphogenesis and structure. In this regard, the degree of sequence identity between the HN and F proteins of the two serotypes in the TM and CT domains is low indeed: 30% and 22%, respectively for the TM domain, and 9 and 11% respectively for the CT domain. In light of this low level of sequence relatedness, we have also pursued a parallel strategy of constructing chimeric glycoproteins in which the PIV1 ectodomain of each glycoprotein was fused to the PIV3 TM and CT domains. Regarding possible interaction with the M protein or other internal proteins, it might be that a conserved structure, such as a constellation of charged amino acids, is important rather than a conserved sequence. Alternatively, it might be that interaction of the TM and CT domains of the glycoproteins with internal proteins is not as critical as has been previously thought. It will be possible to examine these factors more closely using the methods and tools provided herein. For example, these factors will be further elucidated by work in progress employing the methods described above to construct a PIV3 virus bearing HN and F of PIV2.

It was expected that rPIV3-1 would require trypsin for efficient replication in tissue culture since this is a property conferred by the PIV1 F glycoprotein, and this was found to be the case. However, it was interesting to observe that rPIV3-1 caused CPE that more closely resembled that of PIV3 parent virus, indicating that a PIV3 gene(s) other than HN or F specifies this phenotype. These roles will also be further elucidated using the methods and tools provided herein to exchange additional gene(s) between the non-cytopathic PIV1 and the cytopathic PIV3.

EXAMPLE XIIRecovery of Live-Attenuated Chimeric Recombinant PIV Encoding
the Internal Proteins of PIV Type 3 and the Surface
Glycoproteins of PIV Type 1

5 In the present Example, a derivative of rPIV3-1
carrying the three temperature-sensitive and attenuating amino
acid coding changes found in the L gene of the live-attenuated
cp45 PIV3 candidate vaccine virus, termed rPIV3-1.cp45L, is
shown to exhibit a temperature sensitive phenotype with a
10 shut-off temperature of 38°C, similar to that of the
recombinant rPIV3cp45L which possesses the same three
mutations. rPIV3-1.cp45L is attenuated in the respiratory
tract of hamsters to the same extent as rPIV3cp45L. Infection
of hamsters with rPIV3-1.cp45L generates a moderate level of
15 hemagglutination-inhibiting antibodies against wt PIV1 and
induces complete resistance to challenge with wild type PIV1.
This demonstrates that attenuated chimeric PIV according to
the invention are capable of inducing a highly effective
immune response against PIV1. This disclosure also confirms
20 the above described data demonstrating that the surface
glycoproteins of parainfluenza viruses are sufficient to
induce a high level of resistance to homologous virus
challenge. Unexpectedly, infection with recombinant chimeric
virus rPIV3-1.cp45L or rPIV3-1, each bearing the surface
25 glycoprotein genes of PIV1 and the internal genes of PIV3,
also induces a moderate level of resistance to replication of
PIV3 challenge virus. This indicates that the internal genes
of PIV3 can independently induce protective immunity against
PIV3 in rodents. Thus, a reverse genetics system for PIV3 as
30 disclosed herein successfully produces live attenuated PIV1
vaccine candidates that are attenuated and protective in
accepted model subjects.

Viruses and cells.

35 The wt PIV1 strain used in this study is
PIV1/Washington/20993/1964 (PIV1/Wash64) (see, eg., Murphy et
al., Infect. Immun. 12:62, 1975 (incorporated herein by
reference in its entirety). Chimeric rPIV3-1, recovered from
chimeric PIV3 cDNA in which the PIV3 F and HN ORFs were

replaced with those of PIV1/Wash64, as described above and in Tao et al., J. Virol. 72:2955, 1998 (incorporated herein by reference in its entirety). These viruses were propagated in LLC-MK2 cells (ATCC CCL 7.1) in Opti-MEM I (Life Technologies, Gaithersburg, MD) with 50 µg/ml gentamicin sulfate, and 0.75 µg/ml trypsin (Catalog No. 3741, Worthington Biochemical Corp., Freehold, NJ). Trypsin is included because the F glycoprotein of PIV1, but not that of PIV3, is dependent on exogenous trypsin for cleavage when grown in cell culture under these conditions. The wt JS strain of human PIV3 virus and its recombinant derivative from cDNA (rPIV3/JS) were propagated as described above and in Durbin et al., Virology 235:323, 1997 (incorporated herein by reference in its entirety). The propagation of cp45, an attenuated derivative of wt PIV3/JS (see above; and Karron et al., J. Infect. Dis. 171:1107, 1995 (incorporated herein by reference in its entirety)), and rPIV3cp45L, a recombinant PIV3 carrying the three ts mutations found in the L gene of cp45, were propagated as described above and in Skiadopoulos et al., J. Virol 72:1762, 1998 (incorporated herein by reference in its entirety). The modified vaccinia Ankara (MVA) recombinant that expresses the bacteriophage T7 RNA polymerase is described in Virology 210: 202, 1995 (incorporated herein by reference in its entirety).

Hep-2 cells, which are used in transfection, were obtained from ATCC (ATCC CCL 23) and maintained in Opti-MEM I with 2% fetal bovine serum (FBS), 50 µg/ml gentamicin sulfate.

Introduction of L mutations into rPIV3-1 antigenomic cDNA.

The three L mutations of cp45 present in the pTM(L)942/992/1558 plasmid, described above (see also, Skiadopoulos et al., J. Virol 72:1762, 1998, were introduced into chimeric cDNA pFLC.2G+.hc (described above; see also, Tao et al., J. Virol 72:2955, 1998), as a 2.8 kb SphI-NheI fragment (nt 11313 to 14092 in PIV3 antigenomic cDNA) to generate the full-length pFLC.2G+.hc.cp45L bearing the PIV1 F and HN ORFs and the three cp45 L gene mutations (Fig. 19).

The specific mutations present in pTM(L)942/992/1558 are indicated in the legend to Figure 19.

Transfection. HEp-2 cell monolayers in six-well plates were grown to confluence and transfections were performed as described above (see also, Tao et al., J Virol 72:2955, 1998). Trypsin was added to a final concentration of 0.75 μ g/ml on day 3 post transfection prior to harvesting on day 4. Cell culture supernatants were clarified and passaged (referred to as passage 1) onto fresh LLC-MK2 cell monolayers. After overnight adsorption, the medium was replaced with fresh Opti-MEM I with 0.75 μ g/ml trypsin. Passage 1 cultures were incubated at 32°C for 4 days, and the virus present in the supernatant was harvested and passaged again under the same conditions (referred to as passage 2). Virus present in the passage 2 harvest was tested for the presence of the PIV1 HN protein by hemagglutination-inhibition (HAI) assay as described above (see also Tao et al., J Virol 72:2955, 1998).

Replication of PIVs in LLC-MK2 at various temperatures.

Plaque enumeration on LLC-MK2 monolayers was performed as described above, with 0.75 μ g/ml trypsin added to the agarose overlay in the case of PIV1, rPIV3-1, and rPIV3-1.cp45L (see also, Tao et al., J Virol 72:2955, 1998). After incubation at various temperatures for 6 days, the agarose overlay was removed and plaques were identified by hemadsorption (HAD) with guinea pig erythrocytes (RBCs).

Replication of PIVs in the respiratory tract of hamsters.

Groups of five hamsters were inoculated intranasally with 0.1 ml of L15 medium containing 10^6 plaque forming units (PFU) of rPIV3/JS, rPIV3cp45L, cp45, PIV1/Wash64, rPIV3-1, or rPIV3-1.cp45L. Hamsters were sacrificed on day 4 post-infection, and their lungs and nasal turbinates were harvested and homogenized. Virus present in the tissue samples was titrated on LLC-MK2 cell monolayers at 32°C as described above and in Tao et al., J Virol 72:2955, 1998. The titers are expressed as reciprocal mean \log_{10} TCID₅₀/gram of tissue for each group.

Immunization and challenge studies in hamsters.

Groups of ten hamsters were immunized intranasally with 10^6 PFU of virus per animal, as described above. Serum was collected for HAI assay prior to infection and on day 33. The level of HAI antibodies present in the sera of each group of 10 hamsters was determined using PIV1/Wash64 and PIV3/JS as antigens, and the HAI titers determined are presented as mean \log_2 (see also, Tao et al., J Virol 72:2955, 1998).

Thirty-five days post-immunization, five hamsters from each group were challenged intranasally with 10^6 PFU of either PIV1/Wash64 or rPIV3/JS. Nasal turbinates and lungs of these challenged hamsters were harvested four days post challenge. Virus titers in tissue samples were determined on LLC-MK2 monolayers as described above and in Tao et al., J Virol 72:2955, 1998, and the titers are presented as mean \log_{10} TCID₅₀/gram of tissue.

RESULTS

Recovery and characterization of the recombinant chimeric virus rPIV3-1.cp45L.

As noted above, the cDNA clone pFLC.2G+.hc, a full-length antigenomic cDNA of PIV3 in which the ORFs encoding the F and HN glycoproteins have been replaced by those of PIV1, was modified by introduction of three amino acid coding changes (designated 942, 992 and 1558, according to amino acid position in the L protein) identified in the L gene of cp45 and shown to be independent *ts* and attenuating mutations (Fig. 19; see also, Skiadopoulos et al., J Virol 72:1762, 1998). Each coding change was marked by the co-introduction of contiguous translationally silent nt substitutions that ablate a naturally-occurring restriction site (Fig. 19; Table 8). The final full-length plasmid construct, pFLC.2G+.hc.cp45L (Fig. 19), encodes a PIV3-1 chimeric antigenomic RNA of 15516 nt in length and conforms to the rule of six (see, Durbin et al., Virology 234:74, 1997, incorporated herein by reference in its entirety). The authenticity of pFLC.2G+.hc.cp45L was confirmed by digestion with appropriate restriction enzymes.

The pFLC.2G+.hc.cp45L cDNA was transfected into HEP-2 cells together with the PIV3 N, P and L support plasmids and infected with MVA-T7 as described above and in Tao et al., J Virol 72:2955, 1998). Virus recovered after two passages on LLC-MK2 cells, termed rPIV3-1.cp45L, was biologically cloned by plaque-to-plaque-to-plaque passage, and amplified virus was analyzed to confirm that it possessed the PIV1 glycoproteins and the three introduced mutations in L. First, the presence of the PIV1 HN protein in rPIV3-1.cp45L was confirmed by reactivity with PIV1 specific antibodies in HAI assay as described above and in Tao et al., J Virol 72:2955, 1998). The presence of the chimeric PIV3-1 HN and F genes as well as the introduced L gene mutations in rPIV3-1.cp45L genomic RNA was confirmed by restriction enzyme digestion or nucleotide sequence analysis of RT-PCR products generated from virion RNA as described above and in Tao et al., J Virol 72:2955, 1998. These data confirmed that rPIV3-1.cp45L is a recombinant chimeric virus bearing the three codon substitutions of the L gene of cp45.

rPIV3-1.cp45L is temperature sensitive.

The three L gene mutations of cp45 were shown above to confer the *ts* phenotype when introduced into wt PIV3 (see also, Skiadopoulos et al., J Virol 72:1762, 1998). To evaluate whether their presence in the chimeric virus would have the same effect, the efficiency of plaque formation of rPIV3-1.cp45L was determined at various temperatures. As shown in Table 15, the three L mutations indeed conferred the *ts* phenotype to the chimeric virus. The level of temperature sensitivity specified by the cp45 L mutations in the recombinant viruses rPIV3cp45L and rPIV3-1.cp45L was equivalent (Table 15), indicating that the effect of the mutations is independent of the PIV3 or PIV1 HN and F glycoproteins. The level of temperature sensitivity of rPIV3cp45L and rPIV3-1.cp45L was comparable to that of the biologically derived cp45 virus, despite the fact that the latter virus possesses mutations outside of L (see, Stokes et al., Virus Res 30:43, 1993, incorporated herein by reference in its entirety).

Table 15. The recombinant chimeric rPIV3-1.cp45L candidate vaccine virus is temperature sensitive

Virus ^a	Virus titer ^b at indicated temperatures (log ₁₀ PFU/ml)					
	32°C	36°C	37°C	38°C	39°C	40°C
rPIV3/JS	7.4	7.2	7.2	7.2	7.2	7.2
PIV1/Wash64 ^c	7.5	7.6	7.5	7.5	7.4	7.2
rPIV3-1	7.5	7.5	7.5	7.2	6.0	6.1
PIV3cp45 ^c	7.4	6.9	6.8	4.7 ^d	<0.7	<0.7
rPIV3cp45L	7.9	7.4	7.7	5.3	1.2	<0.7
rPIV3-1.cp45L	8.1	8.0	8.2	6.1	<0.7	<0.7

^a Virus nomenclature: rPIV3/JS, recombinant wt PIV3 strain JS; PIV1/Wash64, biologically-derived wt PIV1; rPIV3-1, recombinant chimeric PIV3 in which the F and HN ORFs have been replaced with those of PIV1/Wash64; PIV3cp45, biologically-derived cp45 candidate vaccine virus; rPIV3cp45L, recombinant PIV3 containing the three L gene mutations of cp45; rPIV3-1.cp45L, recombinant chimeric rPIV3-1 containing the three L gene mutations of cp45.

^b Virus titers were determined using LLC-MK2 monolayers in 12-well plates. Titters are the average of two assays.

^c Biologically-derived viruses. All others are recombinant viruses.

^d The shut-off temperature, i.e. the lowest restrictive temperature at which a two log₁₀ reduction in virus titer is observed, of each ts virus is indicated in bold.

Level of replication of rPIV3-1.cp45L in hamsters.

The three L gene mutations of cp45 were shown above to confer attenuation of virus replication in the upper and lower respiratory tract of hamsters when introduced into wt PIV3 (see also, Skiadopoulos et al., *J Virol* 72:1762, 1998). Their effect on the chimeric virus was evaluated by intranasal infection of hamsters, as shown in Table 16. These findings indicate that rPIV3-1.cp45L indeed was attenuated at both sites and, furthermore, that its level of attenuation was comparable to that of rPIV3cp45L. Thus, the ability of the cp45 L mutations to confer attenuation, like temperature sensitivity, is independent of the antigenic specificity of the surface glycoproteins.

Table 16. The recombinant chimeric rPIV3-1.cp45L candidate vaccine virus is attenuated in the respiratory tract of hamsters ^a

Virus	Virus titer in indicated tissue (log ₁₀ TCID ₅₀ /g±S.E) ^b	
	Nasal turbinates	Lungs
rPIV3-1.cp45L	4.6±0.3	1.9±0.4
rPIV3-1	6.0±0.3	6.3±0.4
rPIV3cp45L	3.0±0.3	<1.2
rPIV3/JS	5.7±0.3	5.0±0.3

^a Groups of five hamsters were infected intranasally with indicated viruses at a dosage of 10⁶ PFU per hamster. On day 4 post infection, the tissue samples were harvested and assayed for virus.

^b Virus titers are given as Log₁₀TCID₅₀ per gram of tissue.

Infection with rPIV3-1 or rPIV3-1.cp45L, containing the internal proteins of PIV3 and the glycoproteins of PIV1, confers resistance to PIV1 challenge in hamsters.

The chimeric rPIV3-1 virus and its attenuated rPIV3-1.cp45L derivative were evaluated for immunogenicity and protective efficacy in hamsters. As shown in Table 17, infection with either virus induced HAI antibodies against PIV1, but not PIV3, confirming that these chimeric viruses possess the PIV1 HN glycoprotein and are highly immunogenic. The level of HAI antibodies induced by rPIV3-1.cp45L was two-fold less than that by rPIV3-1, which indicates that its attenuation resulted in a modest decrease in immunogenicity. Similarly, rPIV3 and rPIV3cp45L induced HAI antibodies against PIV3, but not PIV1, and the level induced by the attenuated virus was approximately two-fold lower. Despite the restricted replication in hamsters of the recombinant viruses bearing the cp45 L mutations, infection with either rPIV3cp45L or rPIV3-1.cp45L induced complete resistance to replication of challenge virus bearing homologous glycoproteins.

Table 17. The recombinant chimeric rPIV3-1.cp45L candidate vaccine virus induces complete resistance to PIV1 and partial resistance to PIV3 upon challenge in hamsters ^a

Virus used for immunization	Origin of glycoproteins	Post-immunization HAI titer (log ₂ Reciprocal±S.E.)	Virus Titer in Indicated Tissue ^c (log ₁₀ TCID ₅₀ /Gram±S.E.)			
			Replication of PIV1 challenge virus		Replication of PIV3 challenge virus	
			Nasal Turbinates	Lungs	Nasal Turbinates	Lungs
Control ^b	-	≤1	5.0±0.3	4.6±0.5	5.5±0.4	5.0±0.7
rPIV3-1.cp45L	PIV1	6.9±0.5	≤1.2	≤1.2	1.9±0.6	1.7±0.5
rPIV3-1	PIV1	7.9±0.4	≤1.2	≤1.2	2.9±0.3	2.6±0.4
rPIV3cp45L	PIV3	≤1	4.6±0.2	2.8±0.7	≤1.2	≤1.2
rPIV3/JS	PIV3	≤1	4.6±0.4	2.4±0.5	≤1.2	≤1.2
PIV3cp45 ^d	PIV3	≤1	4.9±0.2	2.5±0.8	≤1.2	≤1.2

123

^a Groups of 10 hamsters were immunized intranasally with 10⁶ PFU of indicated viruses. Post-immunization sera were collected on days 33, two days prior to challenge (see ^c).

^b Hamsters in control group were not inoculated.

^c Five weeks after immunization, five hamsters from each group were challenged intranasally with 10⁶ PFU of indicated virus. Tissue samples were harvested 4 days post challenge. Viruses present in tissue samples were titrated on LLC-MK2 monolayers and the data are presented as log₁₀TCID₅₀/gram of tissue ± standard error.

^d Biologically-derived virus.

Infection with rPIV3-1.cp45L also confers resistance to PIV3.

Information on the role of the non-HN or F glycoproteins of PIVs (i.e., the internal proteins) in resistance is limited. The disclosure and use of rPIV3-1 and rPIV3-1.cp45L herein provides an opportunity to examine the role that internal proteins play in resistance to challenge with PIV3, since the only genes shared by immunizing and challenge viruses are the internal protein genes. PIV3 challenge virus replication was significantly restricted in both the upper and lower respiratory tracts by prior infection of hamsters with rPIV3-1 or rPIV3-1.cp45L (Table 17). Thus, these data indicate that the internal proteins of PIV3, like the HN and F proteins, are capable of inducing partial resistance to replication of the challenge PIV3.

Among the findings demonstrated by the immunogenicity and efficacy studies above a particularly unexpected finding was that infection with wt or attenuated rPIV3 induced a 100-fold reduction in the replication of PIV1 challenge virus in the lungs. Thus, infection with one serotype of PIV provided significant protection against a heterologous serotype. This was unexpected in part because previous studies indicated that infection of animals with one type of human PIV did not induce significant heterologous protection against a PIV belonging to a different human serotype, conforming to a general belief that immunity to human PIV infections was largely type-specific (see, eg., Cook et al., Amer. Jour. Hyg. 77:150, 1963; Ray et al., J. Infect. Dis. 162:746, 1990, each incorporated herein by reference in its entirety).

The present Example demonstrates successful exploitation of novel methods and reagents developed for generating PIV3 vaccines to further provide rapid, rational development of live attenuated candidate vaccines for PIV1. A cDNA encoding infectious PIV3 was modified by substitution of the ORFs encoding the PIV1 HN and F protective antigens for their PIV3 counterparts. Subsequently, attenuating mutations, exemplified by three attenuating mutations present in the L

gene of the cp45 PIV3, were incorporated within this modified chimeric PIV3-PIV1 cDNA. From this cDNA, a recombinant virus was recovered bearing the HN and F genes of PIV1, the internal proteins of PIV3, and the PIV3 cp45 L gene mutations. This recombinant, rPIV3-1.cp45L, was temperature sensitive, highly attenuated in hamsters, and highly efficacious against PIV1 challenge in hamsters. The level of temperature sensitivity, attenuation, and immunogenicity exhibited by rPIV3-1.cp45L was comparable to that of cp45 PIV3, indicating that the phenotypes specified by the set of cp45 L gene mutations are independent of the HN and F surface glycoproteins. These findings, which represent the first live attenuated PIV1 vaccine candidate generated by reverse genetics, provide a generally successful scheme for developing vaccines against PIV1.

Little information is known concerning the role that internal proteins of parainfluenza viruses play in resistance to reinfection with homologous virus. Infection with vaccinia recombinants expressing N, epitopes within N, or M reportedly induce resistance to replication of challenge virus, but the magnitude of the resistance reported is less than that induced by vaccinia recombinants bearing HN or F glycoproteins (see, eg., Kast et al., Proc. Natl. Acad. Sci. USA 88:2283, 1991; Sakaguchi et al., J. Gen. Virol. 74:479, 1993; Thomson et al., J. Immunol. 157:822, 1996, each incorporated herein by reference in its entirety). These studies suggested that the internal proteins were making only minor contributions to resistance to reinfection. Therefore, the present disclosure presents unexpected results by showing that prior infection of hamsters with rPIV3-1.cp45L or rPIV3-1 induced about 250-to 4000-fold reduction of replication of PIV3 in both the nasal turbinates and lungs. These two chimeric recombinant viruses differ from the PIV3 challenge virus in that they possess the HN and F glycoproteins of PIV1 rather than PIV3, but they share all other genes with the challenge virus. The HN and F glycoproteins of PIV1 share 47% and 43% sequence identity with those of PIV3, respectively. Although it is likely that the shared internal proteins are mediating the observed

resistance, it is also possible that shared protein sequences between PIV1 and PIV3 F and HN glycoproteins are contributing to the observed immunity. For example, there are 5 stretches in HN and 2 stretches in F extending at least 9 amino acid residues in length that are shared between PIV1 and PIV3 and have the potential to act as protective CTL epitopes. It is reasonable to consider that the shared internal proteins are contributing to the restriction of replication of wt PIV3 challenge virus, since this level of cross-immunity has not been seen in previous studies (see, eg., Cook et al., Amer. Jour. Hyg. 77:150, 1963; Ray et al., J. Infect. Dis. 162:746, 1990, incorporated herein by reference in its entirety).

The finding that the internal PIV3 proteins of the rPIV3-1 and rPIV3-1.cp45L chimeras conferred resistance to PIV3 challenge demonstrates that attenuated derivatives of PIV3 can be used as vectors for PIV1 and PIV2 protective antigens. Following the teachings of the invention, immunization with one PIV3-based live-attenuated vaccine virus can restrict the replication of other PIV3-based vaccine viruses administered subsequently, thereby decreasing the immunogenicity of the second virus. Since PIV3, like RSV, induces significant illness in early infancy, a combined RSV-PIV3 vaccine for use in the very young 2- to 4-week old infant is therefore an important aspect of the invention (see, eg., Collins et al., Fields Virology 3rd ed. Philadelphia: Lippincott-Raven Publishers, 1205(1), 1996; Reed et al., J. Infect. Dis. 175:807, 1997, each incorporated herein by reference in its entirety). According to this aspect of the invention, immunization with a PIV1-PIV2 vaccine will be preferably initiated at about 6 months of age, since most PIV1 and PIV2 disease occurs after the age of six months. In the possible circumstance that immunization with rPIV3 cp45 significantly inhibits replication of a chimeric recombinant PIV3-1 vaccine virus with which it shares internal protein genes, successful immunization with a recombinant PIV3-1 vaccine may be compromised. In this event, a trivalent PIV vaccine will be administered simultaneously rather than sequentially, thereby preventing the above noted inhibition.

The disclosure herein that infection with a vaccine or wt PIV3 would induce a 100-fold reduction of pulmonary virus replication of the heterologous wt PIV1 was clearly unexpected, in part because the human PIV viruses are
5 serologically distinct by neutralization assay, and previous studies in hamsters found that prior infection with one type of PIV failed to induce resistance to challenge with a high dose of a different PIV type (see eg., Cook et al., Amer. Jour. Hyg. 77:150, 1963; Ray et al., J. Infect. Dis. 162:746,
10 1990; Cook et al., Amer. Jour. Hyg. 69:250, 1959). Furthermore, there is little epidemiological data documenting that prior infection with one PIV significantly modifies subsequent infection with a heterotypic PIV.

In summary, the present Example shows that rPIV3 was
15 successfully converted into a vaccine for PIV1 by substituting the ORFs encoding the F and HN glycoproteins and introducing known attenuating mutations into the PIV3 internal genes. Thus, the extensive methods and reagents provided herein can be applied directly and predictably to attenuating the PIV3
20 backbone of the rPIV3-1 chimeric virus, as well as for generating live-attenuated PIV2 vaccine viruses.

The foregoing disclosure makes it possible to exploit the reagents and methods provided herein to develop a broad assemblage of PIV and related vaccines. In this context,
25 recovery of live, immunogenic chimeras between PIV3 and PIV2 exemplifies powerful new tools for developing a range of recombinant PIV viruses for vaccine use. In conjunction with this work, identification and characterization of the genetic basis for attenuation of naturally occurring PIV mutants,
30 e.g., cp45 and BPIV3 vaccine candidates, following the teachings of the present disclosure also enables development of a large host of recombinant vaccine viruses and subviral particles. In particular, desired mutations present in biologically derived mutant viruses will be readily identified
35 and selected by their introduction, singly and in combination, into a wild type, partially attenuated, or chimeric PIV background, as shown in the Examples above. These findings will expand the menu of exemplary, attenuating mutations

within the invention which can introduced into PIV clones to calibrate the level of attenuation and immunogenicity in vaccine recombinants. Biologically derived mutations can also be introduced within PIV clones having different types of mutations, e.g., mutations involving alterations, deletions, or substitutions of a gene or gene segment. Within this aspect of the invention, recombinant PIV are provided which have a selected gene deletion, addition, or substitution, such as rPIV having a deletion of the C, D or V ORF(s). Such alternatively mutated clones can be further modified according to the present disclosure by introducing one or more mutations specifying a *ts*, *ca* or *att* phenotype adopted from a biologically derived mutant PIV, as exemplified by the PIV recombinants r942, r992, r1558, r942/992, r992/1558, or r942/1558, and r942/992/1558. In additional aspects of the invention, biologically derived mutations will be combined with *de novo* attenuating mutations not found in nature, as exemplified by attenuating gene deletions, e.g., of the C, D and/or V ORFs. Other types of mutations disclosed herein conferring desired phenotypic characteristics will also be combined with biologically derived, attenuating mutations, similar to the range of combinatorial mutations disclosed for recombinant RSV vaccine strains in United States Patent Application Serial No. 08/892,403, filed July 15, 1997 (incorporated herein by reference). Comparable mutations can be readily introduced, e.g., into a chimeric virus, to achieve a desired levels of attenuation and immunogenicity in a chimeric vaccine strain. In this manner, a large menu of mutations are provided within the invention that are useful to engineer a wide assemblage of live attenuated rPIV vaccines having a desired balance of attenuation and immunogenicity, along with other desired phenotypic characteristics.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: The Government of the United States
- (B) STREET: Box OTT
- (C) CITY: Bethesda
- (D) STATE: Maryland
- (E) COUNTRY: United States of America
- (F) POSTAL CODE (ZIP): 20892
- (G) TELEPHONE:
- (H) TELEFAX:
- (I) TELEX:

(ii) TITLE OF INVENTION: PRODUCTION OF ATTENUATED PARAINFLUENZA
VIRUS VACCINES FROM CLONED NUCLEOTIDE SEQUENCE

(iii) NUMBER OF SEQUENCES: 74

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(v) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: WO
- (B) FILING DATE: 22-MAY-1998
- (C) CLASSIFICATION:

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 60/059,385
- (B) FILING DATE: 19-SEP-1997

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 60/047,575
- (B) FILING DATE: 23-MAY-1997

(vii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: King, Jeffrey J.
- (B) REGISTRATION NUMBER: 38,515
- (C) REFERENCE/DOCKET NUMBER: 17634-000320PC

(viii) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 206-467-9600
- (B) TELEFAX: 415-576-0300

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15669 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```
GCTGGGTACC GGGCCGTCG ACGCGTATAT AGTTCCTCCT TTCAGCAAAA AACCCCTCAA      60
GACCCGTTTA GAGGCCCAA GGGGTTATGC TACTGCAGGC TCTCCCTTAG CCATCCGAGT      120
```

GGACGTGCGT CCTCCTTCGG ATGCCCAGGT CGGACCGCGA GGAGGTGGAG ATGCCATGCC	180
GACCCACCAA ACAAGAGAAG AAAGTTGTCT GGGAATATAA ATTTAACTTT AAATTAAGTT	240
AGGATTAAAG ACATTGACTA GAAGGTCAAG AAAAGGGAAC TCTATAATTT CAAAAATGTT	300
GAGCCTATTT GATACATTTA ATGCACGTAG GCAAGAAAAC ATAACAAAAT CAGCCGGTGG	360
AGCTATCATT CCTGGACAGA AAAATACTGT CTCTATATTC GCCCTTGGAC CGACAATAAC	420
TGATGATAAT GAGAAAATGA CATTAGCTCT TCTATTTCTA TCTCATTAC TAGATAATGA	480
GAAACAACAT GCACAAAGGG CAGGGTTCTT GGTGTCTTTA TTGTCAATGG CTTATGCCAA	540
TCCAGAGCTC TACCTAACAA CAAATGGAAG TAATGCAGAT GTCAAGTATG TCATATACAT	600
GATTGAGAAA GATCTAAAAC GGCAAAAGTA TGGAGGATTT GTGGTTAAGA CGAGAGAGAT	660
GATATATGAA AAGACAACTG ATTGGATATT TGGAAGTGAC CTGGATTATG ATCAGGAAAC	720
TATGTTGCAG AACGGCAGGA ACAATTCAAC AATTGAAGAC CTTGTCCACA CATTGCGGTA	780
TCCATCATGT TTAGGAGCTC TTATAATACA GATCTGGATA GTTCTGGTCA AAGCTATCAC	840
TAGTATCTCA GGGTTAAGAA AAGGCTTTTT CACCCGATTG GAAGCTTCA GACAAGATGG	900
AACAGTGCAG GCAGGGCTGG TATTGAGCGG TGACACAGTG GATCAGATTG GGTCAATCAT	960
GCGGTCTCAA CAGAGCTTGG TAACTCTTAT GGTTGAAACA TTAATAACAA TGAATACCAG	1020
CAGAAATGAC CTCACAACCA TAGAAAAGAA TATACAAATT GTTGGCAACT ACATAAGAGA	1080
TGCAGGTCTC GCTTCATTCT TCAATACAAT CAGATATGGA ATTGAGACCA GAATGGCAGC	1140
TTTGACTCTA TCCACTCTCA GACCAGATAT CAATAGATTA AAAGCTTTGA TGGAAGTGTG	1200
TTTATCAAAG GGACCACGCG CTCCTTTTCAT CTGTATCCTC AGAGATCCTA TACATGGTGA	1260
GTTCGCACCA GGCAACTATC CTGCCATATG GAGCTATGCA ATGGGGGTGG CAGTTGTACA	1320
AAATAGAGCC ATGCAACAGT ATGTGACGGG AAGATCATAT CTAGACATTG ATATGTTCCA	1380
GCTAGGACAA GCAGTAGCAC GTGATGCCGA AGCTCAAATG AGCTCAACAC TGGAAGATGA	1440
ACTTGGAGTG ACACACGAAT CTAAAGAAAG CTTGAAGAGA CATATAAGGA ACATAAACAG	1500
TTCAGAGACA TCTTTCCACA AACCGACAGG TGGATCAGCC ATAGAGATGG CAATAGATGA	1560
AGAGCCAGAA CAATTCGAAC ATAGAGCAGA TCAAGAACAA AATGGAGAAC CTCAATCATC	1620
CATAATTCAA TATGCCTGGG CAGAAGGAAA TAGAAGCGAT GATCAGACTG AGCAAGCTAC	1680
AGAATCTGAC AATATCAAGA CCGAACAACA AAACATCAGA GACAGACTAA ACAAGAGACT	1740
CAACGACAAG AAGAAACAAA GCAGTCAACC ACCCACTAAT CCCACAAACA GAACAAACCA	1800
GGACGAAATA GATGATCTGT TTAACGCATT TGGAAGCAAC TAATCGAATC AACATTTTAA	1860
TCTAAATCAA TAATAAATAA GAAAACTTA GGATTAAAGA ATCCTATCAT ACCGGAATAT	1920
AGGGTGGTAA ATTTAGAGTC TGCTTGAAAC TCAATCAATA GAGAGTTGAT GGAAAGCGAT	1980
GCTAAAAACT ATCAAATCAT GGATTCTTGG GAAGAGGAAT CAAGAGATAA ATCAACTAAT	2040
ATCTCCTCGG CCCTCAACAT CATTGAATTC ATACTCAGCA CCGACCCCCA AGAAGACTTA	2100
TCGGAAAACG ACACAATCAA CACAAGAACC CAGCAACTCA GTGCCACCAT CTGTCAACCA	2160
GAAATCAAAC CAACAGAAAC AAGTGAGAAA GATAGTGGAT CAACTGACAA AAATAGACAG	2220

TCTGGGTCAT CACACGAATG TACAACAGAA GCAAAAGATA GAAATATTGA TCAGGAAACT	2280
GTACAGAGAG GACCTGGGAG AAGAAGCAGC TCAGATAGTA GAGCTGAGAC TGTGGTCTCT	2340
GGAGGAATCC CCAGAAGCAT CACAGATTCT AAAAATGGAA CCCAAAACAC GGAGGATATT	2400
GATCTCAATG AAATTAGAAA GATGGATAAG GACTCTATTG AGGGGAAAAT GCGACAATCT	2460
GCAAATGTTC CAAGCGAGAT ATCAGGAAGT GATGACATAT TTACAACAGA ACAAAGTAGA	2520
AACAGTGATC ATGGAAGAAG CCTGGAATCT ATCAGTACAC CTGATACAAG ATCAATAAGT	2580
GTTGTACTG CTGCAACACC AGATGATGAA GAAGAAATAC TAATGAAAA TAGTAGGACA	2640
AAGAAAAGTT CTTCAACACA TCAAGAAGAT GACAAAAGAA TTA AAAAAGG GGGAAAAGGG	2700
AAAGACTGGT TTAAGAAATC AAAAGATACC GACAACCAGA TACCAACATC AGACTACAGA	2760
TCCACATCAA AAGGGCAGAA GAAAATCTCA AAGACAACAA CCACCAACAC CGACACAAAG	2820
GGGCAACAG AAATACAGAC AGAATCATCA GAAACACAAT CCTCATCATG GAATCTCATC	2880
ATCGACAACA ACACCGACCG GAACGAACAG ACAAGCACAA CTCCTCCAAC AACAACTTCC	2940
AGATCAACTT ATACAAAAGA ATCGATCCGA ACAAACCTCTG AATCCAAACC CAAGACACAA	3000
AAGACAAATG GAAAGGAAAG GAAGGATACA GAAGAGAGCA ATCGATTTAC AGAGAGGGCA	3060
ATTACTCTAT TGCAGAATCT TGGTGAATT CAATCCACAT CAAAACCTAGA TTTATATCAA	3120
GACAAACGAG TTGTATGTGT AGCAAATGTA CTAAACAATG TAGATACTGC ATCAAAGATA	3180
GATTTCTCTG CAGGATTAGT CATAGGGGTT TCAATGGACA ACGACACAAA ATTAACACAG	3240
ATACAAAATG AAATGCTAAA CCTCAAAGCA GATCTAAAGA AAATGGACGA ATCACATAGA	3300
AGATTGATAG AAAATCAAAG AGAACAACCTG TCATTGATCA CGTCACTAAT TTCAAATCTC	3360
AAAATTATGA CTGAGAGAGG AGGAAAGAAA GACCAAAATG AATCCAATGA GAGAGTATCC	3420
ATGATCAAAA CAAAATTGAA AGAAGAAAAG ATCAAGAAGA CCAGGTTTGA CCCACTTATG	3480
GAGGCACAAG GCATTGACAA GAATATACCC GATCTATATC GACATGCAGG AGATACACTA	3540
GAGAACGATG TACAAGTTAA ATCAGAGATA TTAAGTTCAT ACAATGAGTC AAATGCAACA	3600
AGACTAATAC CAAAAAAGT GAGCAGTACA ATGAGATCAC TAGTTGCAGT CATCAACAAC	3660
AGCAATCTCT CACAAAGCAC AAAACAATCA TACATAAAG AACTCAAACG TTGCAAAAAT	3720
GATGAAGAAG TATCTGAATT AATGGACATG TTCAATGAAG ATGTCAACAA TTGCCAATGA	3780
TCCAACAAAG AAACGACACC GAACAAACAG ACAAGAAACA ACAGTAGATC AAAACCTGTC	3840
AACACACACA AAATCAAGCA GAATGAAACA ACAGATATCA ATCAATATAC AAATAAGAAA	3900
AACTTAGGAT TAAAGAATAA ATTAATCCTT GTCCAAAATG AGTATAACTA ACTCTGCAAT	3960
ATACACATTC CCAGAATCAT CATTCTCTGA AAATGGTCAT ATAGAACCAT TACCACTCAA	4020
AGTCAATGAA CAGAGGAAAG CAGTACCCCA CATTAGAGTT GCCAAGATCG GAAATCCACC	4080
AAAACACGGA TCCCGGTATT TAGATGTCTT CTTACTCGGC TTCTTCGAGA TGGAACGAAT	4140
CAAAGACAAA TACGGGAGTG TGAATGATCT CGACAGTGAC CCGAGTTACA AAGTTTGTGG	4200
CTCTGGATCA TTACCAATCG GATTGGCTAA GTACACTGGG AATGACCAGG AATTGTGACA	4260
AGCCGCAACC AACTGGATA TAGAAGTGAG AAGAACAGTC AAAGCGAAAG AGATGGTTGT	4320

TTACACGGTA	CAAAATATAA	AACCAGAACT	GTACCCATGG	TCCAATAGAC	TAAGAAAAGG	4380
AATGCTGTTC	GATGCCAACA	AAGTTGCTCT	TGCTCCTCAA	TGTCTTCCAC	TAGATAGGAG	4440
CATAAAATTT	AGAGTAATCT	TCGTGAATTG	TACGGCAATT	GGATCAATAA	CCTTGTTCAA	4500
AATTCCTAAG	TCAATGGCAT	CACTATCTCT	ACCCAACACA	ATATCAATCA	ATCTGCAGGT	4560
ACACATAAAA	ACAGGGGTTC	AGACTGATTC	TAAAGGGATA	GTTCAAATTT	TGGATGAGAA	4620
AGGCGAAAAA	TCACTGAATT	TCATGGTCCA	TCTCGGATTG	ATCAAAAGAA	AAGTAGGCAG	4680
AATGTACTCT	GTTGAATACT	GTAAACAGAA	AATCGAGAAA	ATGAGATTGA	TATTTTCTTT	4740
AGGACTAGTT	GGAGGAATCA	GTCTTCATGT	CAATGCAACT	GGGTCCATAT	CAAAAACACT	4800
AGCAAGTCAG	CTGGTATTCA	AAAGAGAGAT	TTGTTATCCT	TTAATGGATC	TAAATCCGCA	4860
TCTCAATCTA	GTTATCTGGG	CTTCATCAGT	AGAGATTACA	AGAGTGGATG	CAATTTTCCA	4920
ACCTTCTTTA	CCTGGCGAGT	TCAGATACTA	TCCTAATATT	ATTGCAAAAG	GAGTTGGGAA	4980
AATCAAACAA	TGGAAC TAGT	AATCTCTATT	TTAGTCCGGA	CGTATCTATT	AAGCCGAAGC	5040
AAATAAAGGA	TAATCAAAAA	CTTAGGACAA	AAGAGGTCAA	TACCAACAAC	TATTAGCAGT	5100
CACACTCGCA	AGAATAAGAG	AGAAGGGACC	AAAAAAGTCA	AATAGGAGAA	ATCAAAACAA	5160
AAGGTACAGA	ACACCAGAAC	AACAAAATCA	AAACATCCAA	CTCACTCAAA	ACAAAAATTC	5220
CAAAAGAGAC	CGGCAACACA	ACAAGCACTG	AACACAATGC	CAACTTCAAT	ACTGCTAATT	5280
ATTACAACCA	TGATCATGGC	ATCTTTCTGC	CAAATAGATA	TCACAAAAC	ACAGCACGTA	5340
GGTGTATTGG	TCAACAGTCC	CAAAGGGATG	AAGATATCAC	AAAAC TTTGA	AACAAGATAT	5400
CTAATTTTGA	GCCTCATACC	AAAAATAGAA	GACTCTAACT	CTTGTGGTGA	CCAACAGATC	5460
AAGCAATACA	AGAAGTTATT	GGATAGACTG	ATCATCCCTT	TATATGATGG	ATTAAGATTA	5520
CAGAAAGATG	TGATAGTAAC	CAATCAAGAA	TCCAATGAAA	ACACTGATCC	CAGAACAAAA	5580
CGATTCTTTG	GAGGGGTAAT	TGGAACCATT	GCTCTGGGAG	TAGCAACCTC	AGCACAAATT	5640
ACAGCGGCAG	TTGCTCTGGT	TGAAGCCAAG	CAGGCAAGAT	CAGACATCGA	AAAAC TCAAA	5700
GAAGCAATTA	GGGACACAAA	CAAAGCAGTG	CAGTCAGTTC	AGAGCTCCAT	AGGAAATTTA	5760
ATAGTAGCAA	TTAAATCAGT	CCAGGATTAT	GTTAACAAAG	AAATCGTGCC	ATCGATTGCG	5820
AGGCTAGGTT	GTGAAGCAGC	AGGACTTCAA	TTAGGAATTG	CATTAACACA	GCATTACTCA	5880
GAATTAACAA	ACATATTTGG	TGATAACATA	GGATCGTTAC	AAGAAAAAGG	AATAAAATTA	5940
CAAGGTATAG	CATCATTATA	CCGCACAAAT	ATCACAGAAA	TATTCACAAC	ATCAACAGTT	6000
GATAAATATG	ATATCTATGA	TCTGTTATTT	ACAGAATCAA	TAAAGGTGAG	AGTTATAGAT	6060
GTTGACTTGA	ATGATTACTC	AATCACCTC	CAAGTCAGAC	TCCCTTTATT	AACTAGGCTG	6120
CTGAACACTC	AGATCTACAA	AGTAGATTCC	ATATCATATA	ACATCCAAAA	CAGAGAATGG	6180
TATATCCCTC	TTCCCAGCCA	TATCATGACG	AAAGGGGCAT	TTCTAGGTGG	AGCAGACGTC	6240
AAAGAATGTA	TAGAAGCATT	CAGCAGCTAT	ATATGCCCTT	CTGATCCAGG	ATTTGTATTA	6300
AACCATGAAA	TAGAGAGCTG	CTTATCAGGA	AACATATCCC	AATGTCCAAG	AACAACGGTC	6360
ACATCAGACA	TTGTTCCAAG	ATATGCATTT	GTCAATGGAG	GAGTGGTTGC	AAACTGTATA	6420

ACAACCACCT GTACATGCAA CGGAATTGGT AATAGAATCA ATCAACCACC TGATCAAGGA	6480
GTAAAAATTA TAACACATAA AGAATGTAGT ACAATAGGTA TCAACGGAAT GCTGTTCAAT	6540
ACAAATAAAG AAGGAACTCT TGCATTCTAT ACACCAAATG ATATAACACT AAACAATTCT	6600
GTTGCACTTG ATCCAATTGA CATATCAATC GAGCTCAACA AGGCCAAATC AGATCTAGAA	6660
GAATCAAAAG AATGGATAAG AAGGTCAAAT CAAAAACTAG ATTCTATTGG AAATTGGCAT	6720
CAATCTAGCA CTACAATCAT AATTATTTTG ATAATGATCA TTATATTGTT TATAATTAAT	6780
ATAACGATAA TTACAATTGC AATTAAGTAT TACAGAATTC AAAAGAGAAA TCGAGTGGAT	6840
CAAAATGACA AGCCATATGT ACTAACAAAC AAATAACATA TCTACAGATC ATTAGATATT	6900
AAAATTATAA AAAACTTAGG AGTAAAGTTA CGCAATCCAA CTCTACTCAT ATAATTGAGG	6960
AAGGACCCAA TAGACAAATC CAAATTCGAG ATGGAATACT GGAAGCATAC CAATCACGGA	7020
AAGGATGCTG GTAATGAGCT GGAGACGTCT ATGGCTACTC ATGGCAACAA GCTCACTAAT	7080
AAGATAATAT ACATATTATG GACAATAATC CTGGTGTTAT TATCAATAGT CTTCATCATA	7140
GTGCTAATTA ATTCCATCAA AAGTGAAAAG GCCCACGAAT CATTGCTGCA AGACATAAAT	7200
AATGAGTTTA TGGAAATTAC AGAAAAGATC CAAATGGCAT CGGATAATAC CAATGATCTA	7260
ATACAGTCAG GAGTGAATAC AAGGCTTCTT ACAATTCAGA GTCATGTCCA GAATTACATA	7320
CCAATATCAT TGACACAACA GATGTCAGAT CTTAGGAAAT TCATTAGTGA AATTACAATT	7380
AGAAATGATA ATCAAGAAGT GCTGCCACAA AGAATAACAC ATGATGTAGG TATAAAACCT	7440
TTAAATCCAG ATGATTTTTG GAGATGCACG TCTGGTCTTC CATCTTTAAT GAAAACCTCA	7500
AAAATAAGGT TAATGCCAGG GCCGGGATTA TTAGCTATGC CAACGACTGT TGATGGCTGT	7560
GTTAGAATC CGTCTTTAGT TATAAATGAT CTGATTTATG CTTATACCTC AAATCTAATT	7620
ACTCGAGGTT GTCAGGATAT AGGAAAATCA TATCAAGTCT TACAGATAGG GATAATAACT	7680
GTAAACTCAG ACTTGGTACC TGACTTAAAT CCTAGGATCT CTCATACCTT TAACATAAAT	7740
GACAATAGGA AGTCATGTTT TCTAGCACTC CTAAATATAG ATGTATATCA ACTGTGTTCA	7800
ACTCCCAAAG TTGATGAAAG ATCAGATTAT GCATCATCAG GCATAGAAGA TATTGTACTT	7860
GATATTGTCA ATTATGATGG TTCAATCTCA ACAACAAGAT TTAAGAATAA TAACATAAGC	7920
TTTGATCAAC CATATGCTGC ACTATACCCA TCTGTTGGAC CAGGGATATA CTACAAAGGC	7980
AAAATAATAT TTCTCGGGTA TGGAGGTCTT GAACATCCAA TAAATGAGAA TGTAATCTGC	8040
AACACAACCTG GGTGCCCCGG GAAAACACAG AGAGACTGTA ATCAAGCATC TCATAGTACT	8100
TGGTTTTTCAG ATAGGAGGAT GGTCAACTCC ATCATTGTTG TTGACAAAGG CTTAACTCA	8160
ATTCCAAAAT TGAAAGTATG GACGATATCT ATGCGACAAA ATTACTGGGG GTCAGAAGGA	8220
AGGTTACTTC TACTAGGTAA CAAGATCTAT ATATATACAA GATCTACAAG TTGGCATAGC	8280
AAGTTACAAT TAGGAATAAT TGATATTACT GATTACAGTG ATATAAGGAT AAAATGGACA	8340
TGGCATAATG TGCTATCAAG ACCAGGAAAC AATGAATGTC CATGGGGACA TTCATGTCCA	8400
GATGGATGTA TAACAGGAGT ATATACTGAT GCATATCCAC TCAATCCAC AGGGAGCATT	8460
GTGTCATCTG TCATATTAGA CTCACAAAAA TCGAGAGTGA ACCCAGTCAT AACTTACTCA	8520

ACAGCAACCG AAAGAGTAAA CGAGCTGGCC ATCCTAAACA GAACACTCTC AGCTGGATAT	8580
ACAACAACAA GCTGCATTAC AACTATAAC AAAGGATATT GTTTTCATAT AGTAGAAATA	8640
AATCATAAAA GCTTAAACAC ATTTCAACCC ATGTTGTTCA AAACAGAGAT TCCAAAAAGC	8700
TGCAGTTAAT CATAATTAAC CATAATATGC ATCAATCTAT CTATAATACA AGTATATGAT	8760
AAGTAATCAG CAATCAGACA ATAGACAAAA GGGAAATATA AAAAAGTTAG GAGCAAAGCG	8820
TGCTCGGGAA ATGGACACTG AATCTAACAA TGGCACTGTA TCTGACATAC TCTATCCTGA	8880
GTGTCACCTT AACTCTCCTA TCGTTAAAGG TAAAATAGCA CAATTACACA CTATTATGAG	8940
TCTACCTCAG CTTTATGATA TGGATGACGA CTCAATACTA GTTATCACTA GACAGAAAAT	9000
AAAACTTAAT AAATTGGATA AAAGACAACG ATCTATTAGA AGATTAAAAT TAATATTAAC	9060
TGAAAAAGTG AATGACTTAG GAAAATACAC ATTTATCAGA TATCCAGAAA TGTCAAAAGA	9120
AATGTTCAAA TTATATATAC CTGGTATTAA CAGTAAAGTG ACTGAATTAT TACTTAAAGC	9180
AGATAGAACA TATAGTCAAA TGAAGTATGG ATTAAGAGAT CTATGGATTA ATGTGCTATC	9240
AAAATTAGCC TCAAAAAATG ATGGAAGCAA TTATGATCTT AATGAAGAAA TTAATAATAT	9300
ATCGAAAGTT CACACAACCT ATAAATCAGA TAAATGGTAT AATCCATTCA AAACATGGTT	9360
TACTATCAAG TATGATATGA GAAGATTACA AAAAGCTCGA AATGAGATCA CTTTAAATGT	9420
TGGGAAGGAT TATAACTTGT TAGAAGACCA GAAGAATTTT TTATTGATAC ATCCAGAATT	9480
GGTTTTGATA TTAGATAAAC AAAACTATAA TGGTTATCTA ATTACTCCTG AATTAGTATT	9540
GATGTATTGT GACGTAGTCG AAGGCCGATG GAATATAAGT GCATGTGCTA AGTTAGATCC	9600
AAAATTACAA TCTATGTATC AGAAAGGTAA TAACCTGTGG GAAGTGATAG ATAAATTGTT	9660
TCCAATTATG GGAGAAAAGA CATTTGATGT GATATCGTTA TTAGAACCAC TTGCATTATC	9720
CTTAATTCAA ACTCATGATC CTGTTAAACA ACTAAGAGGA GCTTTTTTTAA ATCATGTGTT	9780
ATCCGAGATG GAATTAATAT TTGAATCTAG AGAATCGATT AAGGAATTTT TGAGTGTAGA	9840
TTACATTGAT AAAATTTTAG ATATATTTAA TAAGTCTACA ATAGATGAAA TAGCAGAGAT	9900
TTTCTCTTTT TTTAGAACAT TTGGGCATCC TCCATTAGAA GCTAGTATTG CAGCAGAAAA	9960
GGTTAGAAAA TATATGTATA TTGGAAAACA ATTAAAATTT GACACTATTA ATAAATGTCA	10020
TGCTATCTTC TGTACAATAA TAATTAACGG ATATAGAGAG AGGCATGGTG GACAGTGGCC	10080
TCCTGTGACA TTACCTGATC ATGCACACGA ATTCATCATA AATGCTTACG GTTCAAACCTC	10140
TGCGATATCA TATGAAAATG CTGTTGATTA TTACCAGAGC TTTATAGGAA TAAAATTCAA	10200
TAAATTCATA GAGCCTCAGT TAGATGAGGA TTTGACAATT TATATGAAAG ATAAAGCATT	10260
ATCTCCAAAA AAATCAAATT GGGACACAGT TTATCCTGCA TCTAATTTAC TGTACCGTAC	10320
TAACGCATCC AACGAATCAC GAAGATTAGT TGAAGTATTT ATAGCAGATA GTAAATTTGA	10380
TCCTCATCAG ATATTGGATT ATGTAGAATC TGGGGACTGG TTAGATGATC CAGAATTTAA	10440
TATTTCTTAT AGTCTTAAAG AAAAAGAGAT CAAACAGGAA GGTAGACTCT TTGCAAAAAT	10500
GACATACAAA ATGAGAGCTA CACAAGTTTT ATCAGAGACC CTAATTGCAA ATAACATAGG	10560
AAAATTCTTT CAAGAAAATG GGATGGTGAA GGGAGAGATT GAATTACTTA AGAGATTAAC	10620

AACCATATCA	ATATCAGGAG	TTCCACGGTA	TAATGAAGTG	TACAATAATT	CTAAAAGCCA	10680
TACAGATGAC	CTTAAAACCT	ACAATAAAAT	AAGTAATCTT	AATTTGTCTT	CTAATCAGAA	10740
ATCAAAGAAA	TTTGAATTCA	AGTCAACGGA	TATCTACAAT	GATGGATACG	AGACTGTGAG	10800
CTGTTTCCTA	ACAACAGATC	TCAAAAAATA	CTGTCTTAAT	TGGAGATATG	AATCAACAGC	10860
TCTATTTGGA	GAAACTTGCA	ACCAAATATT	TGGATTAAAT	AAATTGTTTA	ATTGGTTACA	10920
CCCTCGTCTT	GAAGGAAGTA	CAATCTATGT	AGGTGATCCT	TACTGTCCTC	CATCAGATAA	10980
AGAACATATA	TCATTAGAGG	ATCACCCCTGA	TTCTGGTTTT	TACGTTTCATA	ACCCAAGAGG	11040
GGGTATAGAA	GGATTTTGTC	AAAAATTATG	GACACTCATA	TCTATAAGTG	CAATACATCT	11100
AGCAGCTGTT	AGAATAGGCG	TGAGGGTGAC	TGCAATGGTT	CAAGGAGACA	ATCAAGCTAT	11160
AGCTGTAACC	ACAAGAGTAC	CCAACAATTA	TGACTACAGA	GTTAAGAAGG	AGATAGTTTA	11220
TAAAGATGTA	GTGAGATTTT	TTGATTCATT	AAGAGAAGTG	ATGGATGATC	TAGGTCATGA	11280
ACTTAAATTA	AATGAAACGA	TTATAAGTAG	CAAGATGTTT	ATATATAGCA	AAAGAATCTA	11340
TTATGATGGG	AGAATTCTTC	CTCAAGCTCT	AAAAGCATT	TCTAGATGTG	TCTTCTGGTC	11400
AGAGACAGTA	ATAGACGAAA	CAAGATCAGC	ATCTTCAAAT	TTGGCAACAT	CATTTGCAAA	11460
AGCAATTGAG	AATGGTTATT	CACCTGTTCT	AGGATATGCA	TGCTCAATTT	TTAAGAATAT	11520
TCAACAATA	TATATTGCCC	TTGGGATGAA	TATCAATCCA	ACTATAACAC	AGAATATCAG	11580
AGATCAGTAT	TTTAGGAATC	CAAATTGGAT	GCAATATGCC	TCTTTAATAC	CTGCTAGTGT	11640
TGGGGGATTC	AATTACATGG	CCATGTCAAG	ATGTTTTGTA	AGGAATATTG	GTGATCCATC	11700
AGTTGCCGCA	TTGGCTGATA	TTAAAAGATT	TATTAAGGCG	AATCTATTAG	ACCGAAGTGT	11760
TCTTTATAGG	ATTATGAATC	AAGAACCAGG	TGAGTCATCT	TTTTTGGACT	GGGCTTCAGA	11820
TCCATATTCA	TGCAATTTAC	CACAATCTCA	AAATATAACC	ACCATGATAA	AAAATATAAC	11880
AGCAAGGAAT	GTATTACAAG	ATTCACCAAA	TCCATTATTA	TCTGGATTAT	TCACAAATAC	11940
AATGATAGAA	GAAGATGAAG	AATTAGCTGA	GTTCTTGATG	GACAGGAAGG	TAATTCTCCC	12000
TAGAGTTGCA	CATGATATTC	TAGATAATTC	TCTCACAGGA	ATTAGAAATG	CCATAGCTGG	12060
AATGTTAGAT	ACGACAAAAT	CACTAATTCG	GGTTGGCATA	AATAGAGGAG	GA CTGACATA	12120
TAGTTTGTTG	AGGAAAATCA	GTAATTACGA	TCTAGTACAA	TATGAAACAC	TAAGTAGGAC	12180
TTTGCGACTA	ATTGTAAGTG	ATAAAATCAA	GTATGAAGAT	ATGTGTTCCG	TAGACCTTGC	12240
CATAGCATTG	CGACAAAAGA	TGTGGATTCA	TTTATCAGGA	GGAAGGATGA	TAAGTGGACT	12300
TGAAACGCCT	GACCCATTAG	AATTACTATC	TGGGGTAGTA	ATAACAGGAT	CAGAACATTG	12360
TAAAATATGT	TATCTTTCAG	ATGGCACAAA	CCCATATACT	TGGATGTATT	TACCCGGTAA	12420
TATCAAAAATA	GGATCAGCAG	AAACAGGTAT	ATCGTCATTA	AGAGTTCCTT	ATTTTGGATC	12480
AGTCACTGAT	GAAAGATCTG	AAGCACAAAT	AGGATATATC	AAGAATCTTA	GTAAACCTGC	12540
AAAAGCCGCA	ATAAGAATAG	CAATGATATA	TACATGGGCA	TTGGTAATG	ATGAGATATC	12600
TTGGATGGAA	GCCTCACAGA	TAGCACAAAC	ACGTGCAAAT	TTTACACTAG	ATAGTCTCAA	12660
AATTTTAACA	CCGGTAGCTA	CATCAACAAA	TTTATCACAC	AGATTAAAGG	ATACTGCAAC	12720

TCAGATGAAA	TTCTCCAGTA	CATCATTGAT	CAGAGTCAGC	AGATTCATAA	CAATGTCCAA	12780
TGATAACATG	TCTATCAAAG	AAGCTAATGA	AACCAAAGAT	ACTAATCTTA	TTTATCAACA	12840
AATAATGTTA	ACAGGATTAA	GTGTTTTTCGA	ATATTTATTT	AGATTAAAAG	AAACCACAGG	12900
ACACAACCTT	ATAGTTATGC	ATCTGCACAT	AGAAGATGAG	TGTTGTATTA	AAGAAAGTTT	12960
TAATGATGAA	CATATTAATC	CAGAGTCTAC	ATTAGAATTA	ATTGATATC	CTGAAAGTAA	13020
TGAATTTATT	TATGATAAAG	ACCCACTCAA	AGATGTGGAC	TTATCAAAAC	TTATGGTTAT	13080
TAAAGACCAT	TCTTACACAA	TTGATATGAA	TTATTGGGAT	GATACTGACA	TCATACATGC	13140
AATTTCAATA	TGTACTGCAA	TTACAATAGC	AGATACTATG	TCACAATTAG	ATCGAGATAA	13200
TTTAAAAGAG	ATAATAGTTA	TTGCAAATGA	TGATGATATT	AATAGCTTAA	TCACTGAATT	13260
TTTGACTCTT	GACATACTTG	TATTTCTCAA	GACATTTGGT	GGATTATTAG	TAAATCAATT	13320
TGCATACACT	CTTTATAGTC	TAAAAATAGA	AGGTAGGGAT	CTCATTTGGG	ATTATATAAT	13380
GAGAACACTG	AGAGATACTT	CCCATTCAAT	ATTAAAAGTA	TTATCTAATG	CATTATCTCA	13440
TCCTAAAGTA	TTCAAGAGGT	TCTGGGATTG	TGGAGTTTTA	AACCCTATTT	ATGGTCCTAA	13500
TACTGCTAGT	CAAGACCAGA	TAAAACTTGC	CCTATCTATA	TGTGAATATT	CACTAGATCT	13560
ATTTATGAGA	GAATGGTTGA	ATGGTGTATC	ACTTGAAATA	TACATTTGTG	ACAGCGATAT	13620
GGAAGTTGCA	AATGATAGGA	AACAAGCCTT	TATTTCTAGA	CACCTTTCAT	TTGTTTGTG	13680
TTTAGCAGAA	ATTGCATCTT	TCGGACCTAA	CCTGTTAAAC	TTAACATACT	TGGAGAGACT	13740
TGATCTATTG	AAACAATATC	TTGAATTAAA	TATTAAAGAA	GACCCTACTC	TTAAATATGT	13800
ACAAATATCT	GGATTATTAA	TTAAATCGTT	CCCATCAACT	GTAACATACG	TAAGAAAGAC	13860
TGCAATCAAA	TATCTAAGGA	TTGCGGGTAT	TAGTCCACCT	GAGGTAATTG	ATGATTGGGA	13920
TCCGGTAGAA	GATGAAAATA	TGCTGGATAA	CATTGTCAAA	ACTATAAATG	ATAACTGTAA	13980
TAAAGATAAT	AAAGGGAATA	AAATTAACAA	TTTCTGGGGA	CTAGCACTTA	AGAACTATCA	14040
AGTCCTTAAA	ATCAGATCTA	TAACAAGTGA	TTCTGATGAT	AATGATAGAC	TAGATGCTAA	14100
TACAAGTGGT	TTGACACTTC	CTCAAGGAGG	GAATTATCTA	TCGCATCAAT	TGAGATTATT	14160
CGGAATCAAC	AGCACTAGTT	GTCTGAAAGC	TCTTGAGTTA	TCACAAATTT	TAATGAAGGA	14220
AGTCAATAAA	GACAAGGACA	GGCTCTTCCT	GGGAGAAGGA	GCAGGAGCTA	TGCTAGCATG	14280
TTATGATGCC	ACATTAGGAC	CTGCAGTTAA	TTATTATAAT	TCAGGTTTGA	ATATAACAGA	14340
TGTAATTGGT	CAACGAGAAT	TGAAAATATT	TCCTTCAGAG	GTATCATTAG	TAGGTAAAAA	14400
ATTAGGAAAT	GTGACACAGA	TTCTTAACAG	GGTAAAAGTA	CTGTTCAATG	GGAATCCTAA	14460
TTCAACATGG	ATAGGAAATA	TGGAATGTGA	GAGCTTAATA	TGGAGTGAAT	TAAATGATAA	14520
GTCCATTGGA	TTAGTACATT	GTGATATGGA	AGGAGCTATC	GGTAAATCAG	AAGAAACTGT	14580
TCTACATGAA	CATTATAGTG	TTATAAGAAT	TACATACTTG	ATTGGGGATG	ATGATGTTGT	14640
TTTAGTTTCC	AAAATTATAC	CTACAATCAC	TCCGAATTGG	TCTAGAATAC	TTTATCTATA	14700
TAAATTATAT	TGGAAAGATG	TAAGTATAAT	ATCACTCAAA	ACTTCTAATC	CTGCATCAAC	14760
AGAATTATAT	CTAATTTCTGA	AAGATGCATA	TTGTACTATA	ATGGAACCTA	GTGAAATTGT	14820

137

```

TTTATCAAAA CTAAAAGAT TGCTACTCTT GGAAGAAAAT AATCTATTAA AATGGATCAT 14880
TTTATCAAAG AAGAGGAATA ATGAATGGTT ACATCATGAA ATCAAAGAAG GAGAAAGAGA 14940
TTATGGAATC ATGAGACCAT ATCATATGGC ACTACAAATC TTTGGATTTC AAATCAATTT 15000
AAATCATCTG GCGAAAGAAT TTTTATCAAC CCCAGATCTG ACTAATATCA ACAATATAAT 15060
CCAAAGTTTT CAGCGAACAA TAAAGGATGT TTTATTTGAA TGGATTAATA TAACTCATGA 15120
TGATAAGAGA CATAAATTAG GCGGAAGATA TAACATATTC CCACTGAAAA ATAAGGGAAA 15180
GTTAAGACTG CTATCGAGAA GACTAGTATT AAGTTGGATT TCATTATCAT TATCGACTCG 15240
ATTACTTACA GGTCGCTTTC CTGATGAAAA ATTTGAACAT AGAGCACAGA CTGGATATGT 15300
ATCATTAGCT GATACTGATT TAGAATCATT AAAGTTATTG TCGAAAAACA TCATTAAGAA 15360
TTACAGAGAG TGTATAGGAT CAATATCATA TTGGTTTCTA ACCAAAGAAG TTAAAATACT 15420
TATGAAATTG ATCGGTGGTG CTAAATTATT AGGAATTTCC AGACAATATA AAGAACCCGA 15480
AGACCAGTTA TTAGAAAAC TACAATCAACA TGATGAATTT GATATCGATT AAAACATAAA 15540
TACATGAAG ATATATCCTA ACCTTTATCT TTAAGCCTAG GAATAGACAA AAAGTAAGAA 15600
AAACATGTAA TATATATATA CCAAACAGAG TTCTTCTCTT GTTTGGTTAT AGTGAGTCGT 15660
ATTACAATC 15669

```

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AATACGACTC ACTATAACCA AACAAGAGAA C 31

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCAAGTACTA TGAGATGCTT CATT 24

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

138

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCCTATAATT TCAACATGTT GAGCCTATTT G

31

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GATTAAAATG TTGGTCGACT TAGTTGCTTC C

31

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCATAGAGAG TCCATGGAAA GCGACGCTAA AAACATATC

38

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGTGTCGTT TCTTTGTCGA CTCATTGGCA ATTGTTG

37

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCAAAGCGTG CCCGGGCCAT GGACACTGAA TCTAACAATG GC

42

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAAATTCCTT AATCGATTCT CTAGATTC

28

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CCCATCAACT GTAACATACG TAAGAAAGAC

30

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGTTAGGATA TGTCGACATT GTATTTATG

29

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 43 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGGGTTATGC TACTGCAGGC TTTTTCCTC CCTTAGCCAT CCG

43

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:

140

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CTCCATTCTA GANTTATAAA AATTATAGAG TTCCC

35

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15660 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TAATACGACT CACTATAACC AAACAAGAGA AGAACTTGT CTGGAATAT AAATTAACT	60
TTAAATTAAC TTAGGATTAA AGACATTGAC TAGAAGGTCA AGAAAAGGGA ACTCTATAAT	120
TTCAAAAATG TTGAGCCTAT TTGATACATT TAATGCACGT AGGCAAGAAA ACATAACAAA	180
ATCAGCCGGT GGAGCTATCA TTCCTGGACA GAAAATACT GTCTCTATAT TCGCCCTTGG	240
ACCGACAATA ACTGATGATA ATGAGAAAAT GACATTAGCT CTTCTATTTT TATCTCATTC	300
ACTAGATAAT GAGAAACAAC ATGCACAAAG GGCAGGGTTC TTGGTGTCTT TATTGTCAAT	360
GGCTTATGCC AATCCAGAGC TCTACCTAAC AACAAATGGA AGTAATGCAG ATGTCAAGTA	420
TGTCATATAC ATGATTGAGA AAGATCTAAA ACGGCAAAAG TATGGAGGAT TTGTGGTTAA	480
GACGAGAGAG ATGATATATG AAAAGACAAC TGATTGGATA TTTGGAAGTG ACCTGGATTA	540
TGATCAGGAA ACTATGTTGC AGAACGGCAG GAACAATTCA ACAATTGAAG ACCTTGTTCA	600
CACATTTGGG TATCCATCAT GTTTAGGAGC TCTTATAATA CAGATCTGGA TAGTTCTGGT	660
CAAAGCTATC ACTAGTATCT CAGGGTTAAG AAAAGGCTTT TTCACCCGAT TGGAAGCTTT	720
CAGACAAGAT GGAACAGTGC AGGCAGGGCT GGTATTGAGC GGTGACACAG TGGATCAGAT	780
TGGGTCAATC ATGCGGTCTC AACAGAGCTT GGTAACCTCT ATGGTTGAAA CATTAATAAC	840
AATGAATACC AGCAGAAATG ACCTCACAAC CATAGAAAAG AATATACAAA TTGTTGGCAA	900
CTACATAAGA GATGCAGGTC TCGCTTCATT CTTCAATACA ATCAGATATG GAATTGAGAC	960
CAGAATGGCA GCTTTGACTC TATCCACTCT CAGACCAGAT ATCAATAGAT TAAAAGCTTT	1020
GATGGAAGT TATTTATCAA AGGGACCACG CGCTCCTTTC ATCTGTATCC TCAGAGATCC	1080
TATACATGGT GAGTTCGCAC CAGGCAACTA TCCTGCCATA TGGAGCTATG CAATGGGGGT	1140
GGCAGTTGTA CAAATAGAG CCATGCAACA GTATGTGACG GGAAGATCAT ATCTAGACAT	1200
TGATATGTTC CAGCTAGGAC AAGCAGTAGC ACGTGATGCC GAAGCTCAAA TGAGCTCAAC	1260

ACTGGAAGAT	GAAGCTGGAG	TGACACACGA	ATCTAAAGAA	AGCTTGAAGA	GACATATAAG	1320
GAACATAAAC	AGTTCAGAGA	CATCTTTCCA	CAAACCGACA	GGTGGATCAG	CCATAGAGAT	1380
GGCAATAGAT	GAAGAGCCAG	AACAATTCTGA	ACATAGAGCA	GATCAAGAAC	AAAATGGAGA	1440
ACCTCAATCA	TCCATAATTC	AATATGCCTG	GGCAGAAGGA	AATAGAAGCG	ATGATCAGAC	1500
TGAGCAAGCT	ACAGAATCTG	ACAATATCAA	GACCGAACAA	CAAAACATCA	GAGACAGACT	1560
AAACAAGAGA	CTCAACGACA	AGAAGAAACA	AAGCAGTCAA	CCACCCACTA	ATCCCACAAA	1620
CAGAACAAAC	CAGGACGAAA	TAGATGATCT	GTTTAAACGCA	TTTGAAGCA	ACTAATCGAA	1680
TCAACATTTT	AATCTAAATC	ATAATAAAAT	AAGAAAAACT	TAGGATTAAA	GAATCCTATC	1740
ATACCGGAAT	ATAGGGTGGT	AAATTTAGAG	TCTGCTTGAA	ACTCAATCAA	TAGAGAGTTG	1800
ATGGAAAGCG	ATGCTAAAAA	CTATCAAATC	ATGGATTCTT	GGGAAGAGGA	ATCAAGAGAT	1860
AAATCAACTA	ATATCTCCTC	GGCCCTCAAC	ATCATTGAAT	TCATACTCAG	CACCGACCCC	1920
CAAGAAGACT	TATCGGAAAA	CGACACAATC	AACACAAGAA	CCCAGCAACT	CAGTGCCACC	1980
ATCTGTCAAC	CAGAAATCAA	ACCAACAGAA	ACAAGTGAGA	AAGATAGTGG	ATCAACTGAC	2040
AAAAATAGAC	AGTCTGGGTC	ATCACACGAA	TGTACAACAG	AAGCAAAAAG	TAGAAATATT	2100
GATCAGGAAA	CTGTACAGAG	AGGACCTGGG	AGAAGAAGCA	GCTCAGATAG	TAGAGCTGAG	2160
ACTGTGGTCT	CTGGAGGAAT	CCCCAGAAGC	ATCACAGATT	CTAAAAATGG	AACCCAAAAC	2220
ACGGAGGATA	TTGATCTCAA	TGAAATTAGA	AAGATGGATA	AGGACTCTAT	TGAGGGGAAA	2280
ATGCGACAAT	CTGCAAATGT	TCCAAGCGAG	ATATCAGGAA	GTGATGACAT	ATTTACAACA	2340
GAACAAAGTA	GAAACAGTGA	TCATGGAAGA	AGCCTGGAAT	CTATCAGTAC	ACCTGATACA	2400
AGATCAATAA	GTGTTGTTAC	TGCTGCAACA	CCAGATGATG	AAGAAGAAAT	ACTAATGAAA	2460
AATAGTAGGA	CAAAGAAAAG	TTCTTCAACA	CATCAAGAAG	ATGACAAAAG	AATTAAAAAA	2520
GGGGGAAAAG	GGAAAGACTG	GTTTAAGAAA	TCAAAAGATA	CCGACAACCA	GATACCAACA	2580
TCAGACTACA	GATCCACATC	AAAAGGGCAG	AAGAAAATCT	CAAAGACAAC	AACCACCAAC	2640
ACCGACACAA	AGGGGCAAAC	AGAAATACAG	ACAGAATCAT	CAGAAACACA	ATCCTCATCA	2700
TGGAATCTCA	TCATCGACAA	CAACACCGAC	CGGAACGAAC	AGACAAGCAC	AACTCCTCCA	2760
ACAACAACCT	CCAGATCAAC	TTATACAAAA	GAATCGATCC	GAACAACTC	TGAATCCAAA	2820
CCCAAGACAC	AAAAGACAAA	TGGAAAGGAA	AGGAAGGATA	CAGAAGAGAG	CAATCGATTT	2880
ACAGAGAGGG	CAATTACTCT	ATTGCAGAAT	CTTGGTGTAA	TTCAATCCAC	ATCAAAACTA	2940
GATTTATATC	AAGACAAACG	AGTTGTATGT	GTCACAAATG	TACTAAACAA	TGTAGATACT	3000
GCATCAAAGA	TAGATTTCTT	GGCAGGATTA	GTCATAGGGG	TTTCAATGGA	CAACGACACA	3060
AAATTAACAC	AGATACAAAA	TGAAATGCTA	AACCTCAAAG	CAGATCTAAA	GAAAATGGAC	3120
GAATCACATA	GAAGATTGAT	AGAAAATCAA	AGAGAACAAC	TGTCATTGAT	CACGTCACTA	3180
ATTTCAAATC	TCAAAATTAT	GAATGAGAGA	GGAGGAAAGA	AAGACCAAAA	TGAATCCAAT	3240
GAGAGAGTAT	CCATGATCAA	AACAAAATTG	AAAGAAGAAA	AGATCAAGAA	GACCAGGTTT	3300
GACCCACTTA	TGGAGGCACA	AGGCATTGAC	AAGAATATAC	CCGATCTATA	TCGACATGCA	3360

GGAGATACAC	TAGAGAACGA	TGTACAAGTT	AAATCAGAGA	TATTAAGTTC	ATACAATGAG	3420
TCAAATGCAA	CAAGACTAAT	ACCCAAAAAA	GTGAGCAGTA	CAATGAGATC	ACTAGTTGCA	3480
GTCATCAACA	ACAGCAATCT	CTCACAAAGC	ACAAAACAAT	CATACATAAA	CGAACTCAAA	3540
CGTTGCAAAA	ATGATGAAGA	AGTATCTGAA	TTAATGGACA	TGTTCAATGA	AGATGTCAAC	3600
AATTGCCAAT	GATCCAACAA	AGAAACGACA	CCGAACAAAC	AGACAAGAAA	CAACAGTAGA	3660
TCAAAACCTG	TCAACACACA	CAAAATCAAG	CAGAATGAAA	CAACAGATAT	CAATCAATAT	3720
ACAAATAAGA	AAAACCTTAGG	ATTAAAGAAT	AAATTAATCC	TTGTCCAAAA	TGAGTATAAC	3780
TAACTCTGCA	ATATACACAT	TCCCAGAATC	ATCATTCTCT	GAAAATGGTC	ATATAGAACC	3840
ATTACCACTC	AAAGTCAATG	AACAGAGGAA	AGCAGTACCC	CACATTAGAG	TTGCCAAGAT	3900
CGGAAATCCA	CCAAAACACG	GATCCCGGTA	TTTAGATGTC	TTCTTACTCG	GCTTCTTCGA	3960
GATGGAACGA	ATCAAAGACA	AATACGGGAG	TGTGAATGAT	CTCGACAGTG	ACCCGAGTTA	4020
CAAAGTTTGT	GGCTCTGGAT	CATTACCAAT	CGGATTGGCT	AAGTACACTG	GGAATGACCA	4080
GGAATTGTTA	CAAGCCGCAA	CCAACTGGA	TATAGAAGTG	AGAAGAACAG	TCAAAGCGAA	4140
AGAGATGGTT	GTTTACACGG	TACAAAATAT	AAAACCAGAA	CTGTACCCAT	GGTCCAATAG	4200
ACTAAGAAAA	GGAATGCTGT	TCGATGCCAA	CAAAGTTGCT	CTTGCTCCTC	AATGTCTTCC	4260
ACTAGATAGG	AGCATAAAAAT	TTAGAGTAAT	CTTCGTGAAT	TGTACGGCAA	TTGGATCAAT	4320
AACCTTGTTT	AAAATTCCTA	AGTCAATGGC	ATCACTATCT	CTACCCAACA	CAATATCAAT	4380
CAATCTGCAG	GTACACATAA	AAACAGGGGT	TCAGACTGAT	TCTAAAGGGA	TAGTTCAAAT	4440
TTTGGATGAG	AAAGGCGAAA	AATCACTGAA	TTTCATGGTC	CATCTCGGAT	TGATCAAAAAG	4500
AAAAGTAGGC	AGAATGTACT	CTGTTGAATA	CTGTAAACAG	AAAATCGAGA	AAATGAGATT	4560
GATATTTTCT	TTAGGACTAG	TTGGAGGAAT	CAGTCTTCAT	GTCAATGCAA	CTGGGTCCAT	4620
ATCAAAAACA	CTAGCAAGTC	AGCTGGTATT	CAAAGAGAG	ATTTGTTATC	CTTTAATGGA	4680
TCTAAATCCG	CATCTCAATC	TAGTTATCTG	GGCTTCATCA	GTAGAGATTA	CAAGAGTGGA	4740
TGCAATTTTC	CAACCTTCTT	TACCTGGCGA	GTTGAGATAC	TATCCTAATA	TTATTGCAAA	4800
AGGAGTTGGG	AAAATCAAAC	AATGGAACATA	GTAATCTCTA	TTTTAGTCCG	GACGTATCTA	4860
TTAAGCCGAA	GCAAATAAAG	GATAATCAAA	AACTTAGGAC	AAAAGAGGTC	AATACCAACA	4920
ACTATTAGCA	GTCACACTCG	CAAGAATAAG	AGAGAAGGGA	CCAAAAAAGT	CAAATAGGAG	4980
AAATCAAAAC	AAAAGGTACA	GAACACCAGA	ACAACAAAAT	CAAAACATCC	AACTCACTCA	5040
AAACAAAAAT	TCCAAAAGAG	ACCGGCAACA	CAACAAGCAC	TGAACACAAT	GCCAACTTCA	5100
ATACTGCTAA	TTATTACAAC	CATGATCATG	GCATCTTTCT	GCCAAATAGA	TATCACAAAA	5160
CTACAGCACG	TAGGTGTATT	GGTCAACAGT	CCCAAAGGGA	TGAAGATATC	ACAAAACTTT	5220
GAAACAAGAT	ATCTAATTTT	GAGCCTCATA	CCAAAAATAG	AAGACTCTAA	CTCTTGTTGGT	5280
GACCAACAGA	TCAAGCAATA	CAAGAAGTTA	TTGGATAGAC	TGATCATCCC	TTTATATGAT	5340
GGATTAAGAT	TACAGAAAGA	TGTGATAGTA	ACCAATCAAG	AATCCAATGA	AAACACTGAT	5400
CCCAGAACAA	AACGATTCTT	TGGAGGGGTA	ATTGGAACCA	TTGCTCTGGG	AGTAGCAACC	5460

TCAGCACAAA TTACAGCGGC AGTTGCTCTG GTTGAAGCCA AGCAGGCAAG ATCAGACATC	5520
GAAAACTCA AAGAAGCAAT TAGGGACACA AACAAAGCAG TGCAGTCAGT TCAGAGCTCC	5580
ATAGGAAATT TAATAGTAGC AATTAAATCA GTCCAGGATT ATGTTAACAA AGAAATCGTG	5640
CCATCGATTG CGAGGCTAGG TTGTGAAGCA GCAGGACTTC AATTAGGAAT TGCATTAACA	5700
CAGCATTACT CAGAATTAAC AAACATATTT GGTGATAACA TAGGATCGTT ACAAGAAAAA	5760
GGAATAAAAT TACAAGGTAT AGCATCATTA TACCGCACAA ATATCACAGA AATATTCACA	5820
ACATCAACAG TTGATAAATA TGATATCTAT GATCTGTTAT TTACAGAATC AATAAAGGTG	5880
AGAGTTATAG ATGTTGACTT GAATGATTAC TCAATCACCC TCCAAGTCAG ACTCCCTTTA	5940
TTAACTAGGC TGCTGAACAC TCAGATCTAC AAAGTAGATT CCATATCATA TAACATCCAA	6000
AACAGAGAAT GGTATATCCC TCTTCCCAGC CATATCATGA CGAAAGGGGC ATTTCTAGGT	6060
GGAGCAGACG TCAAAGAATG TATAGAAGCA TTCAGCAGCT ATATATGCCC TTCTGATCCA	6120
GGATTTGTAT TAAACCATGA AATAGAGAGC TGCTTATCAG GAAACATATC CCAATGTCCA	6180
AGAACAACGG TCACATCAGA CATTGTTCCA AGATATGCAT TTGTCAATGG AGGAGTGGTT	6240
GCAAACTGTA TAACAACCAC CTGTACATGC AACGGAATTG GTAATAGAAT CAATCAACCA	6300
CCTGATCAAG GAGTAAAAAT TATAACACAT AAAGAATGTA GTACAATAGG TATCAACGGA	6360
ATGCTGTTCA ATACAAATAA AGAAGGAACT CTTGCATTCT ATACACCAA TGATATAACA	6420
CTAAACAATT CTGTTGCACT TGATCCAATT GACATATCAA TCGAGCTCAA CAAGGCCAAA	6480
TCAGATCTAG AAGAATCAAA AGAATGGATA AGAAGGTCAA ATCAAAAAT AGATTCTATT	6540
GGAAATTGGC ATCAATCTAG CACTACAATC ATAATTATTT TGATAATGAT CATTATATTG	6600
TTTATAATTA ATATAACGAT AATTACAATT GCAATTAAGT ATTACAGAAT TCAAAAGAGA	6660
AATCGAGTGG ATCAAAATGA CAAGCCATAT GTACTAACAA ACAAATAACA TATCTACAGA	6720
TCATTAGATA TTAAAATTAT AAAAACTTA GGAGTAAAGT TACGCAATCC AACTCTACTC	6780
ATATAATTGA GGAAGGACCC AATAGACAAA TCCAAATTCG AGATGGAATA CTGGAAGCAT	6840
ACCAATCACG GAAAGGATGC TGGTAATGAG CTGGAGACGT CTATGGCTAC TCATGGCAAC	6900
AAGCTCACTA ATAAGATAAT ATACATATTA TGGACAATAA TCCTGGTGTT ATTATCAATA	6960
GTCTTCATCA TAGTGCTAAT TAATTCCATC AAAAGTGAAA AGGCCACGA ATCATTGCTG	7020
CAAGACATAA ATAATGAGTT TATGGAAATT ACAGAAAAGA TCCAAATGGC ATCGGATAAT	7080
ACCAATGATC TAATACAGTC AGGAGTGAAT ACAAGGCTTC TTACAATTCA GAGTCATGTC	7140
CAGAATTACA TACCAATATC ATTGACACAA CAGATGTCAG ATCTTAGGAA ATTCATTAGT	7200
GAAATTACAA TTAGAAATGA TAATCAAGAA GTGCTGCCAC AAAGAATAAC ACATGATGTA	7260
GGTATAAAAC CTTTAAATCC AGATGATTTT TGGAGATGCA CGTCTGGTCT TCCATCTTTA	7320
ATGAAAACCTC CAAAAATAAG GTTAATGCCA GGGCCGGGAT TATTAGCTAT GCCAACGACT	7380
GTTGATGGCT GTGTTAGAAC TCCGTCTTTA GTTATAAATG ATCTGATTTA TGCTTATACC	7440
TCAAATCTAA TTAATCGAGG TTGTCAGGAT ATAGGAAAAT CATATCAAGT CTTACAGATA	7500
GGGATAATAA CTGTAAACTC AGACTTGGTA CCTGACTTAA ATCCTAGGAT CTCTCATACC	7560

TTTAACATAA	ATGACAATAG	GAAGTCATGT	TCTCTAGCAC	TCCTAAATAT	AGATGTATAT	7620
CAACTGTGTT	CAACTCCCAA	AGTTGATGAA	AGATCAGATT	ATGCATCATC	AGGCATAGAA	7680
GATATTGTAC	TTGATATTGT	CAATTATGAT	GGTTCAATCT	CAACAACAAG	ATTTAAGAAT	7740
AATAACATAA	GCTTTGATCA	ACCATATGCT	GCACTATACC	CATCTGTTGG	ACCAGGGATA	7800
TACTACAAAG	GCAAAATAAT	ATTTCTCGGG	TATGGAGGTC	TTGAACATCC	AATAAATGAG	7860
AATGTAATCT	GCAACACAAC	TGGGTGCCCC	GGGAAAACAC	AGAGAGACTG	TAATCAAGCA	7920
TCTCATAGTA	CTTGGTTTTT	AGATAGGAGG	ATGGTCAACT	CCATCATTGT	TGTTGACAAA	7980
GGCTTAAACT	CAATTCCAAA	ATTGAAAGTA	TGGACGATAT	CTATGCGACA	AAATTACTGG	8040
GGGTCAGAAG	GAAGGTTACT	TCTACTAGGT	AACAAGATCT	ATATATATAC	AAGATCTACA	8100
AGTTGGCATA	GCAAGTTACA	ATTAGGAATA	ATTGATATTA	CTGATTACAG	TGATATAAGG	8160
ATAAAATGGA	CATGGCATAA	TGTGCTATCA	AGACCAGGAA	ACAATGAATG	TCCATGGGGA	8220
CATTCATGTC	CAGATGGATG	TATAACAGGA	GTATATACTG	ATGCATATCC	ACTCAATCCC	8280
ACAGGGAGCA	TTGTGTCATC	TGTCATATTA	GACTCACAAA	AATCGAGAGT	GAACCCAGTC	8340
ATAACTTACT	CAACAGCAAC	CGAAAGAGTA	AACGAGCTGG	CCATCCTAAA	CAGAACACTC	8400
TCAGCTGGAT	ATACAACAAC	AAGCTGCATT	ACACACTATA	ACAAAGGATA	TTGTTTTTCAT	8460
ATAGTAGAAA	TAAATCATAA	AAGCTTAAAC	ACATTTCAAC	CCATGTTGTT	CAAAACAGAG	8520
ATTCCAAAAA	GCTGCGAGTTA	ATCATAATTA	ACCATAATAT	GCATCAATCT	ATCTATAATA	8580
CAAGTATATG	ATAAGTAATC	AGCAATCAGA	CAATAGACAA	AAGGGAAATA	TAAAAAACTT	8640
AGGAGCAAAG	CGTGCTCGGG	AAATGGACAC	TGAATCTAAC	AATGGCACTG	TATCTGACAT	8700
ACTCTATCCT	GAGTGTCACC	TTAACTCTCC	TATCGTTAAA	GGTAAAATAG	CACAATTACA	8760
CACTATTATG	AGTCTACCTC	AGCCTTATGA	TATGGATGAC	GACTCAATAC	TAGTTATCAC	8820
TAGACAGAAA	ATAAACTTA	ATAAATTGGA	TAAAAGACAA	CGATCTATTA	GAAGATTAAA	8880
ATTAATATTA	ACTGAAAAAG	TGAATGACTT	AGGAAAATAC	ACATTTATCA	GATATCCAGA	8940
AATGTCAAAA	GAAATGTTCA	AATTATATAT	ACCTGGTATT	AACAGTAAAG	TGACTGAATT	9000
ATTACTTAAA	GCAGATAGAA	CATATAGTCA	AATGACTGAT	GGATTAAGAG	ATCTATGGAT	9060
TAATGTGCTA	TCAAAATTAG	CCTCAAAAAA	TGATGGAAGC	AATTATGATC	TTAATGAAGA	9120
AATTAATAAT	ATATCGAAAG	TTCACACAAC	CTATAAATCA	GATAAATGGT	ATAATCCATT	9180
CAAAACATGG	TTTACTATCA	AGTATGATAT	GAGAAGATTA	CAAAAAGCTC	GAAATGAGAT	9240
CACTTTTAAT	GTTGGGAAGG	ATTATAACTT	GTTAGAAGAC	CAGAAGAATT	TCTTATTGAT	9300
ACATCCAGAA	TTGGTTTTGA	TATTAGATAA	ACAAAACAT	AATGGTTATC	TAATTACTCC	9360
TGAATTAGTA	TTGATGTATT	GTGACGTAGT	CGAAGGCCGA	TGGAATATAA	GTGCATGTGC	9420
TAAGTTAGAT	CCAAAATTAC	AATCTATGTA	TCAGAAAGGT	AATAACCTGT	GGGAAGTGAT	9480
AGATAAATTG	TTTCCAATTA	TGGGAGAAAA	GACATTTGAT	GTGATATCGT	TATTAGAACC	9540
ACTTGCAATTA	TCCTTAATTC	AACTCATGA	TCCTGTTAAA	CAACTAAGAG	GAGCTTTTTT	9600
AAATCATGTG	TTATCCGAGA	TGGAATTAAT	ATTTGAATCT	AGAGAATCGA	TTAAGGAATT	9660

TCTGAGTGTA	GATTACATTG	ATAAAATTTT	AGATATATTT	AATAAGTCTA	CAATAGATGA	9720
AATAGCAGAG	ATTTTCTCTT	TTTTTAGAAC	ATTTGGGCAT	CCTCCATTAG	AAGCTAGTAT	9780
TGCAGCAGAA	AAGGTTAGAA	AATATATGTA	TATTGGAAAA	CAATTAAAAT	TTGACACTAT	9840
TAATAAATGT	CATGCTATCT	TCTGTACAAT	AATAATTAAC	GGATATAGAG	AGAGGCATGG	9900
TGGACAGTGG	CCTCCTGTGA	CATTACCTGA	TCATGCACAC	GAATTCATCA	TAAATGCTTA	9960
CGGTTCAAAC	TCTGCGATAT	CATATGAAAA	TGCTGTTGAT	TATTACCAGA	GCTTTATAGG	10020
AATAAAATTC	AATAAATTCA	TAGAGCCTCA	GTTAGATGAG	GATTTGACAA	TTTATATGAA	10080
AGATAAAGCA	TTATCTCCAA	AAAAATCAAA	TTGGGACACA	GTTTATCCTG	CATCTAATTT	10140
ACTGTACCGT	ACTAACGCAT	CCAACGAATC	ACGAAGATTA	GTTGAAGTAT	TTATAGCAGA	10200
TAGTAAATTT	GATCCTCATC	AGATATTGGA	TTATGTAGAA	TCTGGGGACT	GGTTAGATGA	10260
TCCAGAATTT	AATATTTCTT	ATAGTCTTAA	AGAAAAAGAG	ATCAAACAGG	AAGGTAGACT	10320
CTTTGCAAAA	ATGACATACA	AAATGAGAGC	TACACAAGTT	TTATCAGAGA	CCCTACTTGC	10380
AAATAACATA	GGAAAATTCT	TTCAAGAAAA	TGGGATGGTG	AAGGGAGAGA	TTGAATTACT	10440
TAAGAGATTA	ACAACCATAT	CAATATCAGG	AGTTCCACGG	TATAATGAAG	TGTACAATAA	10500
TTCTAAAAGC	CATACAGATG	ACCTTAAAAC	CTACAATAAA	ATAAGTAATC	TTAATTTGTC	10560
TTCTAATCAG	AAATCAAAGA	AATTTGAATT	CAAGTCAACG	GATATCTACA	ATGATGGATA	10620
CGAGACTGTG	AGCTGTTTCC	TAACAACAGA	TCTCAAAAAA	TACTGTCTTA	ATTGGAGATA	10680
TGAATCAACA	GCTCTATTTG	GAGAACTTG	CAACCAAATA	TTTGGATTAA	ATAAATTGTT	10740
TAATTGGTTA	CACCCTCGTC	TTGAAGGAAG	TACAATCTAT	GTAGGTGATC	CTTACTGTCC	10800
TCCATCAGAT	AAAGAACATA	TATCATTAGA	GGATCACCTT	GATTCTGGTT	TTTACGTTCA	10860
TAACCCAAGA	GGGGGTATAG	AAGGATTTTG	TCAAAAATTA	TGGACACTCA	TATCTATAAG	10920
TGCAATACAT	CTAGCAGCTG	TTAGAATAGG	CGTGAGGGTG	ACTGCAATGG	TTCAAGGAGA	10980
CAATCAAGCT	ATAGCTGTAA	CCACAAGAGT	ACCCAACAAT	TATGACTACA	GAGTTAAGAA	11040
GGAGATAGTT	TATAAAGATG	TAGTGAGATT	TTTTGATTCA	TTAAGAGAAG	TGATGGATGA	11100
TCTAGGTCAT	GAACCTAAAT	TAAATGAAAC	GATTATAAGT	AGCAAGATGT	TCATATATAG	11160
CAAAAGAATC	TATTATGATG	GGAGAATTCT	TCCTCAAGCT	CTAAAAGCAT	TATCTAGATG	11220
TGTCTTCTGG	TCAGAGACAG	TAATAGACGA	AACAAGATCA	GCATCTTCAA	ATTTGGCAAC	11280
ATCATTTGCA	AAAGCAATTG	AGAATGGTTA	TTCACCTGTT	CTAGGATATG	CATGCTCAAT	11340
TTTTAAGAAT	ATTCAACAAC	TATATATTGC	CCTTGGGATG	AATATCAATC	CAACTATAAC	11400
ACAGAATATC	AGAGATCAGT	ATTTTAGGAA	TCCAATTGG	ATGCAATATG	CCTCTTTAAT	11460
ACCTGCTAGT	GTTGGGGGAT	TCAATTACAT	GGCCATGTCA	AGATGTTTTG	TAAGGAATAT	11520
TGGTGATCCA	TCAGTTGCCG	CATTGGCTGA	TATTAAGAAG	TTTATTAAGG	CGAATCTATT	11580
AGACCGAAGT	GTTCTTTATA	GGATTATGAA	TCAAGAACCA	GGTGAGTCAT	CTTTTTTGGA	11640
CTGGGCTTCA	GATCCATATT	CATGCAATTT	ACCACAATCT	CAAAATATAA	CCACCATGAT	11700
AAAAAATATA	ACAGCAAGGA	ATGTATTACA	AGATTCACCA	AATCCATTAT	TATCTGGATT	11760

ATTCACAAAT	ACAATGATAG	AAGAAGATGA	AGAATTAGCT	GAGTTCCTGA	TGGACAGGAA	11820
GGTAATTCTC	CCTAGAGTTG	CACATGATAT	TCTAGATAAT	TCTCTCACAG	GAATTAGAAA	11880
TGCCATAGCT	GGAATGTTAG	ATACGACAAA	ATCACTAATT	CGGGTTGGCA	TAAATAGAGG	11940
AGGACTGACA	TATAGTTTGT	TGAGGAAAAT	CAGTAATTAC	GATCTAGTAC	AATATGAAAC	12000
ACTAAGTAGG	ACTTTGCGAC	TAATTGTAAG	TGATAAAATC	AAGTATGAAG	ATATGTGTTC	12060
GGTAGACCTT	GCCATAGCAT	TGCGACAAAA	GATGTGGATT	CATTTATCAG	GAGGAAGGAT	12120
GATAAGTGGA	CTTGAAACGC	CTGACCCATT	AGAATTACTA	TCTGGGGTAG	TAATAACAGG	12180
ATCAGAACAT	TGTAAAATAT	GTTATTCTTC	AGATGGCACA	AACCCATATA	CTTGGATGTA	12240
TTTACCCGGT	AATATCAAAA	TAGGATCAGC	AGAAACAGGT	ATATCGTCAT	TAAGAGTTCC	12300
TTATTTTGGA	TCAGTCACTG	ATGAAAGATC	TGAAGCACAA	TTAGGATATA	TCAAGAATCT	12360
TAGTAAACCT	GCAAAAGCCG	CAATAAGAAT	AGCAATGATA	TATACATGGG	CATTGTGTAA	12420
TGATGAGATA	TCTTGGATGG	AAGCCTCACA	GATAGCACAA	ACACGTGCAA	ATTTTACACT	12480
AGATAGTCTC	AAAATTTTAA	CACCGGTAGC	TACATCAACA	AATTTATCAC	ACAGATTAAA	12540
GGATACTGCA	ACTCAGATGA	AATTCTCCAG	TACATCATTG	ATCAGAGTCA	GCAGATTCAT	12600
AACAATGTCC	AATGATAACA	TGTCTATCAA	AGAAGCTAAT	GAAACCAAAG	ATACTAATCT	12660
TATTTATCAA	CAAATAATGT	TAACAGGATT	AAGTGTTTTT	GAATATTTAT	TTAGATTAAA	12720
AGAAACCACA	GGACACAACC	CTATAGTTAT	GCATCTGCAC	ATAGAAGATG	AGTGTTGTAT	12780
TAAAGAAAGT	TTTAATGATG	AACATATTAA	TCCAGAGTCT	ACATTAGAAT	TAATTCGATA	12840
TCCTGAAAGT	AATGAATTTA	TTTATGATAA	AGACCCACTC	AAAGATGTGG	ACTTATCAAA	12900
ACTTATGGTT	ATTAAAGACC	ATTCTTACAC	AATTGATATG	AATTATTGGG	ATGATACTGA	12960
CATCATACAT	GCAATTTCAA	TATGTACTGC	AATTACAATA	GCAGATACTA	TGTCACAATT	13020
AGATCGAGAT	AATTTAAAAG	AGATAATAGT	TATTGCAAAT	GATGATGATA	TTAATAGCTT	13080
AATCACTGAA	TTTTTGACTC	TTGACATACT	TGTATTTCTC	AAGACATTTG	GTGGATTATT	13140
AGTAAATCAA	TTTGCATACA	CTCTTTATAG	TCTAAAAATA	GAAGGTAGGG	ATCTCATTTG	13200
GGATTATATA	ATGAGAACAC	TGAGAGATAC	TTCCCATTC	ATATTAAAAG	TATTATCTAA	13260
TGCATTATCT	CATCCTAAAG	TATTCAAGAG	GTTCTGGGAT	TGTGGAGTTT	TAAACCCTAT	13320
TTATGGTCCT	AATACTGCTA	GTCAAGACCA	GATAAAACTT	GCCCTATCTA	TATGTGAATA	13380
TTCACTAGAT	CTATTTATGA	GAGAATGGTT	GAATGGTGTA	TCACTTGAAA	TATACATTTG	13440
TGACAGCGAT	ATGGAAGTTG	CAAATGATAG	GAAACAAGCC	TTTATTTCTA	GACACCTTTC	13500
ATTTGTTTGT	TGTTTAGCAG	AAATTGCATC	TTTCGGACCT	AACCTGTTAA	ACTTAACATA	13560
CTTGGAGAGA	CTTGATCTAT	TGAAACAATA	TCTTGAATTA	AATATTAAAG	AAGACCCTAC	13620
TCTTAAATAT	GTACAAATAT	CTGGATTATT	AATTAAATCG	TTCCCATCAA	CTGTAACATA	13680
CGTAAGAAAG	ACTGCAATCA	AATATCTAAG	GATTCGCGGT	ATTAGTCCAC	CTGAGGTAAT	13740
TGATGATTGG	GATCCGGTAG	AAGATGAAAA	TATGCTGGAT	AACATTGTCA	AAACTATAAA	13800
TGATAACTGT	AATAAAGATA	ATAAAGGGAA	TAAAATTAAC	AATTTCTGGG	GACTAGCACT	13860

TAAGAACTAT CAAGTCCTTA AAATCAGATC TATAACAAGT GATTCTGATG ATAATGATAG	13920
ACTAGATGCT AATACAAGTG GTTTGACACT TCCTCAAGGA GGGAATTATC TATCGCATCA	13980
ATTGAGATTA TTCGGAATCA ACAGCACTAG TTGTCTGAAA GCTCTTGAGT TATCACAAAT	14040
TTTAATGAAG GAAGTCAATA AAGACAAGGA CAGGCTCTTC CTGGGAGAAG GAGCAGGAGC	14100
TATGCTAGCA TGTATGATG CCACATTAGG ACCTGCAGTT AATTATTATA ATTCAGGTTT	14160
GAATATAACA GATGTAATTG GTCAACGAGA ATTGAAAATA TTTCCTTCAG AGGTATCATT	14220
AGTAGGTAAA AAATTAGGAA ATGTGACACA GATTCTTAAC AGGGTAAAAG TACTGTTCAA	14280
TGGGAATCCT AATTCAACAT GGATAGGAAA TATGGAATGT GAGAGCTTAA TATGGAGTGA	14340
ATTAAATGAT AAGTCCATTG GATTAGTACA TTGTGATATG GAAGGAGCTA TCGGTAAATC	14400
AGAAGAACT GTTCTACATG AACATTATAG TGTTATAAGA ATTACATACT TGATTGGGGA	14460
TGATGATGTT GTTTTAGTTT CCAAAATTAT ACCTACAATC ACTCCGAATT GGTCTAGAAT	14520
ACTTTATCTA TATAAATTAT ATTGGAAAGA TGTAAGTATA ATATCACTCA AAACCTCTAA	14580
TCCTGCATCA ACAGAATTAT ATCTAATTTT GAAAGATGCA TATTGTACTA TAATGGAACC	14640
TAGTGAAAT GTTTTATCAA AACTTAAAAG ATTGTCATC TTGGAAGAAA ATAATCTATT	14700
AAAATGGATC ATTTTATCAA AGAAGAGGAA TAATGAATGG TTACATCATG AAATCAAAGA	14760
AGGAGAAAGA GATTATGGAA TCATGAGACC ATATCATATG GCACTACAAA TCTTTGGATT	14820
TCAAATCAAT TTAAATCATC TGGCGAAAGA ATTTTATCA ACCCCAGATC TGACTAATAT	14880
CAACAATATA ATCCAAAGTT TTCAGCGAAC AATAAAGGAT GTTTTATTTG AATGGATTAA	14940
TATAACTCAT GATGATAAGA GACATAAATT AGGCGGAAGA TATAACATAT TCCCACTGAA	15000
AAATAAGGGA AAGTTAAGAC TGCTATCGAG AAGACTAGTA TTAAGTTGGA TTTCATTATC	15060
ATTATCGACT CGATTACTTA CAGGTCGCTT TCCTGATGAA AAATTTGAAC ATAGAGCACA	15120
GACTGGATAT GTATCATTAG CTGATACTGA TTTAGAATCA TTAAAGTTAT TGTCGAAAAA	15180
CATCATTAAG AATTACAGAG AGTGTATAGG ATCAATATCA TATTGGTTTC TAACCAAAGA	15240
AGTTAAAATA CTTATGAAAT TGATCGGTGG TGCTAAATTA TTAGGAATTC CCAGACAATA	15300
TAAAGAACCC GAAGACCAGT TATTAGAAAA CTACAATCAA CATGATGAAT TTGATATCGA	15360
TTAAAACATA AATACAATGA AGATATATCC TAACCTTTAT CTTTAAGCCT AGGAATAGAC	15420
AAAAAGTAAG AAAAACATGT AATATATATA TACCAAACAG AGTTCTTCTC TTGTTTGGTG	15480
GGTCGGCATG GCATCTCCAC CTCCTCGCGG TCCGGACCTG GGCATCCGAA GGAGGACGCA	15540
CGTCCACTCG GATGGCTAAG GGAGAGCCTG CAGTAGCATA ACCCCTTGGG GCCTCTAAAC	15600
GGGTCTTGAG GGGTTTTTTG CTGAAAGGAG GAACTATATA CGCGTCGACG GGCCCCGCGC	15660

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15666 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TAATACGACT CACTATAGGA CCAAACAAGA GAAGAAACTT GTCTGGGAAT ATAAATTTAA	60
CTTTAAATTA ACTTAGGATT AAAGACATTG ACTAGAAGGT CAAGAAAAGG GAACTCTATA	120
ATTTCAAAAA TGTGTAGCCT ATTTGATACA TTTAATGCAC GTAGGCAAGA AACATAACA	180
AAATCAGCCG GTGGAGCTAT CATTCCTGGA CAGAAAAATA CTGTCTCTAT ATTCGCCCTT	240
GGACCGACAA TAACTGATGA TAATGAGAAA ATGACATTAG CTCTTCTATT TCTATCTCAT	300
TCACTAGATA ATGAGAAACA ACATGCACAA AGGGCAGGGT TCTTGGTGTC TTTATTGTCA	360
ATGGCTTATG CCAATCCAGA GCTCTACCTA ACAACAAATG GAAGTAATGC AGATGTCAAG	420
TATGTCATAT ACATGATTGA GAAAGATCTA AAACGGCAAA AGTATGGAGG ATTTGTGGTT	480
AAGACGAGAG AGATGATATA TGAAAAGACA ACTGATTGGA TATTTGGAAG TGACCTGGAT	540
TATGATCAGG AAACATATGTT GCAGAACGGC AGGAACAATT CAACAATTGA AGACCTTGTC	600
CACACATTTG GGTATCCATC ATGTTTAGGA GCTCTTATAA TACAGATCTG GATAGTTCTG	660
GTCAAAGCTA TCACTAGTAT CTCAGGGTTA AGAAAAGGCT TTTTCACCCG ATTGGAAGCT	720
TTCAGACAAG ATGGAACAGT GCAGGCAGGG CTGGTATTGA GCGGTGACAC AGTGGATCAG	780
ATTGGGTCAA TCATGCGGTC TCAACAGAGC TTGGTAACTC TTATGGTTGA AACATTAATA	840
ACAATGAATA CCAGCAGAAA TGACCTCACA ACCATAGAAA AGAATATACA AATTGTTGGC	900
AACTACATAA GAGATGCAGG TCTCGCTTCA TTCTTCAATA CAATCAGATA TGGAATTGAG	960
ACCAGAATGG CAGCTTTGAC TCTATCCACT CTCAGACCAG ATATCAATAG ATTAAAAGCT	1020
TTGATGGAAC TGTATTTATC AAAGGGACCA CGCGCTCCTT TCATCTGTAT CCTCAGAGAT	1080
CCTATACATG GTGAGTTCGC ACCAGGCAAC TATCCTGCCA TATGGAGCTA TGCAATGGGG	1140
GTGGCAGTTG TACAAAATAG AGCCATGCAA CAGTATGTGA CGGGAAGATC ATATCTAGAC	1200
ATTGATATGT TCCAGCTAGG ACAAGCAGTA GCACGTGATG CCGAAGCTCA AATGAGCTCA	1260
ACACTGGAAG ATGAACTTGG AGTGACACAC GAATCTAAAG AAAGCTTGAA GAGACATATA	1320
AGGAACATAA ACAGTTCAGA GACATCTTTC CACAAACCGA CAGGTGGATC AGCCATAGAG	1380
ATGGCAATAG ATGAAGAGCC AGAACAATTC GAACATAGAG CAGATCAAGA ACAAATGGA	1440
GAACCTCAAT CATCCATAAT TCAATATGCC TGGGCAGAAG GAAATAGAAG CGATGATCAG	1500
ACTGAGCAAG CTACAGAATC TGACAATATC AAGACCGAAC AACAAAACAT CAGAGACAGA	1560
CTAAACAAGA GACTCAACGA CAAGAAGAAA CAAAGCAGTC AACCACCCAC TAATCCCACA	1620
AACAGAACAA ACCAGGACGA AATAGATGAT CTGTTTAACG CATTTGGAAG CAACTAATCG	1680
AATCAACATT TTAATCTAAA TCAATAATAA ATAAGAAAAA CTTAGGATTA AAGAATCCTA	1740
TCATACCGGA ATATAGGGTG GTAAATTTAG AGTCTGCTTG AAACCTCAATC AATAGAGAGT	1800
TGATGGAAAG CGATGCTAAA AACTATCAAA TCATGGATTC TTGGGAAGAG GAATCAAGAG	1860
ATAAATCAAC TAATATCTCC TCGGCCCTCA ACATCATTGA ATTCATACTC AGCACCGACC	1920

CCCAAGAAGA CTTATCGGAA AACGACACAA TCAACACAAG AACCCAGCAA CTCAGTGCCA 1980
CCATCTGTCA ACCAGAAATC AAACCAACAG AAACAAGTGA GAAAGATAGT GGATCAACTG 2040
ACAAAAATAG ACAGTCTGGG TCATCACACG AATGTACAAC AGAAGCAAAA GATAGAAATA 2100
TTGATCAGGA AACTGTACAG AGAGGACCTG GGAGAAGAAG CAGCTCAGAT AGTAGAGCTG 2160
AGACTGTGGT CTCTGGAGGA ATCCCCAGAA GCATCACAGA TTCTAAAAAT GGAACCCAAA 2220
ACACGGAGGA TATTGATCTC AATGAAATTA GAAAGATGGA TAAGGACTCT ATTGAGGGGA 2280
AAATGCGACA ATCTGCAAAT GTTCCAAGCG AGATATCAGG AAGTGATGAC ATATTTACAA 2340
CAGAACAAAG TAGAAACAGT GATCATGGAA GAAGCCTGGA ATCTATCAGT ACACCTGATA 2400
CAAGATCAAT AAGTGTGTGTT ACTGTGCAA CACCAGATGA TGAAGAAGAA ATACTAATGA 2460
AAAATAGTAG GACAAAGAAA AGTTCTTCAA CACATCAAGA AGATGACAAA AGAATTAAAA 2520
AAGGGGGAAA AGGGAAAGAC TGGTTTAAGA AATCAAAAGA TACCGACAAC CAGATACCAA 2580
CATCAGACTA CAGATCCACA TCAAAAGGGC AGAAGAAAAT CTCAAAGACA ACAACCACCA 2640
ACACCGACAC AAAGGGGCAA ACAGAAATAC AGACAGAATC ATCAGAAACA CAATCCTCAT 2700
CATGGAATCT CATCATCGAC AACAACACCG ACCGGAACGA ACAGACAAGC ACAACTCCTC 2760
CAACAACAAC TTCCAGATCA ACTTATACAA AAGAATCGAT CCGAACAAAC TCTGAATCCA 2820
AACCCAAGAC ACAAAGACA AATGGAAAGG AAAGGAAGGA TACAGAAGAG AGCAATCGAT 2880
TTACAGAGAG GGCAATTACT CTATTGCAGA ATCTTGGTGT AATTCAATCC ACATCAAAAC 2940
TAGATTTATA TCAAGACAAA CGAGTTGTAT GTGTAGCAAA TGTACTAAAC AATGTAGATA 3000
CTGCATCAAA GATAGATTTC CTGGCAGGAT TAGTCATAGG GGTTC AATG GACAACGACA 3060
CAAATTAAC ACAGATACAA AATGAAATGC TAAACCTCAA AGCAGATCTA AAGAAAATGG 3120
ACGAATCACA TAGAAGATTG ATAGAAAATC AAAGAGAACA ACTGTCATTG ATCACGTCAC 3180
TAATTTCAAA TCTCAAAATT ATGACTGAGA GAGGAGGAAA GAAAGACCAA AATGAATCCA 3240
ATGAGAGAGT ATCCATGATC AAAACAAAAT TGAAAGAAGA AAAGATCAAG AAGACCAGGT 3300
TTGACCCACT TATGGAGGCA CAAGGCATTG ACAAGAATAT ACCCGATCTA TATCGACATG 3360
CAGGAGATAC ACTAGAGAAC GATGTACAAG TTAAATCAGA GATATTAAGT TCATACAATG 3420
AGTCAAATGC AACAAGACTA ATACCCAAAA AAGTGAGCAG TACAATGAGA TCACTAGTTG 3480
CAGTCATCAA CAACAGCAAT CTCTCACAAA GCACAAAACA ATCATAcata AACGAACTCA 3540
AACGTTGCAA AAATGATGAA GAAGTATCTG AATTAATGGA CATGTTCAAT GAAGATGTCA 3600
ACAATTGCCA ATGATCCAAC AAAGAAACGA CACCGAACAA ACAGACAAGA AACAACAGTA 3660
GATCAAAACC TGTCAACACA CACAAAATCA AGCAGAATGA AACAACAGAT ATCAATCAAT 3720
ATACAAATAA GAAAACTTA GGATTAAAGA ATAAATTAAT CCTTGTCCAA AATGAGTATA 3780
ACTAACTCTG CAATATACAC ATTCCCAGAA TCATCATTCT CTGAAAATGG TCATATAGAA 3840
CCATTACCAC TCAAAGTCAA TGAACAGAGG AAAGCAGTAC CCCACATTAG AGTTGCCAAG 3900
ATCGGAAATC CACCAAAACA CGGATCCCGG TATTTAGATG TCTTCTTACT CGGCTTCTTC 3960
GAGATGGAAC GAATCAAAGA CAAATACGGG AGTGTGAATG ATCTCGACAG TGACCCGAGT 4020

TACAAAGTTT	GTGGCTCTGG	ATCATTACCA	ATCGGATTGG	CTAAGTACAC	TGGAATGAC	4080
CAGGAATTGT	TACAAGCCGC	AACCAAACCTG	GATATAGAAG	TGAGAAGAAC	AGTCAAAGCG	4140
AAAGAGATGG	TTGTTTACAC	GGTACAAAAT	ATAAAACCAG	AACTGTACCC	ATGGTCCAAT	4200
AGACTAAGAA	AAGGAATGCT	GTTTCGATGCC	AACAAAGTTG	CTCTTGCTCC	TCAATGTCTT	4260
CCACTAGATA	GGAGCATAAA	ATTTAGAGTA	ATCTTCGTGA	ATTGTACGGC	AATTGGATCA	4320
ATAACCTTGT	TCAAAATTC	TAAGTCAATG	GCATCACTAT	CTCTACCCAA	CACAATATCA	4380
ATCAATCTGC	AGGTACACAT	AAAAACAGGG	GTTTCAGACTG	ATTCTAAAGG	GATAGTTCAA	4440
ATTTTGGATG	AGAAAGGCGA	AAAATCACTG	AATTTTCATGG	TCCATCTCGG	ATTGATCAAA	4500
AGAAAAGTAG	GCAGAAATGTA	CTCTGTTGAA	TACTGTAAAC	AGAAAATCGA	GAAAATGAGA	4560
TTGATATTTT	CTTTAGGACT	AGTTGGAGGA	ATCAGTCTTC	ATGTCAATGC	AACTGGGTCC	4620
ATATCAAAAA	CACTAGCAAG	TCAGCTGGTA	TTCAAAAGAG	AGATTTGTGA	TCCTTTAATG	4680
GATCTAAATC	CGCATCTCAA	TCTAGTTATC	TGGGCTTCAT	CAGTAGAGAT	TACAAGAGTG	4740
GATGCAATTT	TCCAACCTTC	TTTACCTGGC	GAGTTCAGAT	ACTATCCTAA	TATTATTGCA	4800
AAAGGAGTTG	GGAAAATCAA	ACAATGGAAC	TAGTAATCTC	TATTTTAGTC	CGGACGTATC	4860
TATTAAGCCG	AAGCAAATAA	AGGATAATCA	AAAACCTTAGG	ACAAAAGAGG	TCAATACCAA	4920
CAACTATTAG	CAGTCACACT	CGCAAGAATA	AGAGAGAAGG	GACCAAAAAA	GTCAAATAGG	4980
AGAAATCAAA	ACAAAAGGTA	CAGAACACCA	GAACAACAAA	ATCAAAACAT	CCAACCTACT	5040
CAAAACAAAA	ATTCCAAAAG	AGACCGGCAA	CACAACAAGC	ACTGAACACA	ATGCCAACTT	5100
CAATACTGCT	AATTATTACA	ACCATGATCA	TGGCATCTTT	CTGCCAAATA	GATATCACAA	5160
AACTACAGCA	CGTAGGTGTA	TTGGTCAACA	GTCCCAAAGG	GATGAAGATA	TCACAAAAC	5220
TTGAAACAAG	ATATCTAATT	TTGAGCCTCA	TACCAAAAAT	AGAAGACTCT	AACTCTTG	5280
GTGACCAACA	GATCAAGCAA	TACAAGAAGT	TATTGGATAG	ACTGATCATC	CCTTTATATG	5340
ATGGATTAAG	ATTACAGAAA	GATGTGATAG	TAACCAATCA	AGAATCCAAT	GAAAACACTG	5400
ATCCCAGAAC	AAAACGATTC	TTTGGAGGGG	TAATTGGAAC	CATTGCTCTG	GGAGTAGCAA	5460
CCTCAGCACA	AATTACAGCG	GCAGTTGCTC	TGGTTGAAGC	CAAGCAGGCA	AGATCAGACA	5520
TCGAAAAACT	CAAAGAAGCA	ATTAGGGACA	CAAACAAAGC	AGTGCAGTCA	GTTTCAGAGCT	5580
CCATAGGAAA	TTTAATAGTA	GCAATTAAAT	CAGTCCAGGA	TTATGTTAAC	AAAGAAATCG	5640
TGCCATCGAT	TGCGAGGCTA	GGTTGTGAAG	CAGCAGGACT	TCAATTAGGA	ATTGCATTAA	5700
CACAGCATTA	CTCAGAATTA	ACAAACATAT	TTGGTGATAA	CATAGGATCG	TTACAAGAAA	5760
AAGGAATAAA	ATTACAAGGT	ATAGCATCAT	TATACCGCAC	AAATATCACA	GAAATATTCA	5820
CAACATCAAC	AGTTGATAAA	TATGATATCT	ATGATCTGTT	ATTTACAGAA	TCAATAAAGG	5880
TGAGAGTTAT	AGATGTTGAC	TTGAATGATT	ACTCAATCAC	CCTCCAAGTC	AGACTCCCTT	5940
TATTAAC	TAGTGAAC	ACTCAGATCT	ACAAAGTAGA	TTCCATATCA	TATAACATCC	6000
AAAACAGAGA	ATGGTATATC	CCTCTTCCCA	GCCATATCAT	GACGAAAGGG	GCATTTCTAG	6060
GTGGAGCAGA	CGTCAAAGAA	TGTATAGAAG	CATTCAGCAG	CTATATATGC	CCTTCTGATC	6120

CAGGATTTGT	ATTAAACCAT	GAAATAGAGA	GCTGCTTATC	AGGAAACATA	TCCCAATGTC	6180
CAAGAACAAC	GGTCACATCA	GACATTGTTC	CAAGATATGC	ATTTGTCAAT	GGAGGAGTGG	6240
TTGCAAACTG	TATAACAACC	ACCTGTACAT	GCAACGGAAT	TGGTAATAGA	ATCAATCAAC	6300
CACCTGATCA	AGGAGTAAAA	ATTATAACAC	ATAAAGAATG	TAGTACAATA	GGTATCAACG	6360
GAATGCTGTT	CAATACAAAT	AAAGAAGGAA	CTCTTGCAAT	CTATACACCA	AATGATATAA	6420
CACTAAACAA	TTCTGTTGCA	CTTGATCCAA	TTGACATATC	AATCGAGCTC	AACAAGGCCA	6480
AATCAGATCT	AGAAGAATCA	AAAGAATGGA	TAAGAAGGTC	AAATCAAAAA	CTAGATTCTA	6540
TTGGAAATTG	GCATCAATCT	AGCACTACAA	TCATAATTAT	TTTGATAATG	ATCATTATAT	6600
TGTTTATAAT	TAATATAACG	ATAATTACAA	TTGCAATTAA	GTATTACAGA	ATTCAAAAAGA	6660
GAAATCGAGT	GGATCAAAAT	GACAAGCCAT	ATGTACTAAC	AAACAAATAA	CATATCTACA	6720
GATCATTAGA	TATTTAAAT	ATAAAAACT	TAGGAGTAAA	GTTACGCAAT	CCAATCTTAC	6780
TCATATAATT	GAGGAAGGAC	CCAATAGACA	AATCCAAATT	CGAGATGGAA	TACTGGAAGC	6840
ATACCAATCA	CGGAAAGGAT	GCTGGTAATG	AGCTGGAGAC	GTCTATGGCT	ACTCATGGCA	6900
ACAAGCTCAC	TAATAAGATA	ATATACATAT	TATGGACAAT	AATCCTGGTG	TTATTATCAA	6960
TAGTCTTCAT	CATAGTGCTA	ATTAATTCCA	TCAAAAGTGA	AAAGGCCAC	GAATCATTGC	7020
TGCAAGACAT	AAATAATGAG	TTTATGGAAA	TTACAGAAAA	GATCCAAATG	GCATCGGATA	7080
ATACCAATGA	TCTAATACAG	TCAGGAGTGA	ATACAAGGCT	TCTTACAATT	CAGAGTCATG	7140
TCCAGAATTA	CATACCAATA	TCATTGACAC	AACAGATGTC	AGATCTTAGG	AAATTCATTA	7200
GTGAAATTAC	AATTAGAAAT	GATAATCAAG	AAGTGCTGCC	ACAAAGAATA	ACACATGATG	7260
TAGGTATAAA	ACCTTTAAAT	CCAGATGATT	TTTGGAGATG	CACGTCTGGT	CTTCCATCTT	7320
TAATGAAAAC	TCCAAAAATA	AGGTTAATGC	CAGGGCCGGG	ATTATTAGCT	ATGCCAACGA	7380
CTGTTGATGG	CTGTGTTAGA	ACTCCGTCTT	TAGTTATAAA	TGATCTGATT	TATGCTTATA	7440
CCTCAAATCT	AATTACTCGA	GGTTGTCAGG	ATATAGGAAA	ATCATATCAA	GTCTTACAGA	7500
TAGGGATAAT	AACTGTAAAC	TCAGACTTGG	TACCTGACTT	AAATCCTAGG	ATCTCTCATA	7560
CCTTTAACAT	AAATGACAAT	AGGAAGTCAT	GTTCTCTAGC	ACTCCTAAAT	ATAGATGTAT	7620
ATCAACTGTG	TTCAACTCCC	AAAGTTGATG	AAAGATCAGA	TTATGCATCA	TCAGGCATAG	7680
AAGATATTGT	ACTTGATATT	GTCAATTATG	ATGGTTCAAT	CTCAACAACA	AGATTTAAGA	7740
ATAATAACAT	AAGCTTTGAT	CAACCATATG	CTGCACTATA	CCCATCTGTT	GGACCAGGGA	7800
TATACTACAA	AGGCAAAATA	ATATTTCTCG	GGTATGGAGG	TCTTGAACAT	CCAATAAATG	7860
AGAATGTAAT	CTGCAACACA	ACTGGGTGCC	CCGGGAAAAC	ACAGAGAGAC	TGTAATCAAG	7920
CATCTCATAG	TACTTGGTTT	TCAGATAGGA	GGATGGTCAA	CTCCATCATT	GTTGTTGACA	7980
AAGGCTTAA	CTCAATTCCA	AAATTGAAAG	TATGGACGAT	ATCTATGCGA	CAAATTACT	8040
GGGGGTCAGA	AGGAAGGTTA	CTTCTACTAG	GTAACAAGAT	CTATATATAT	ACAAGATCTA	8100
CAAGTTGGCA	TAGCAAGTTA	CAATTAGGAA	TAATTGATAT	TACTGATTAC	AGTGATATAA	8160
GGATAAAATG	GACATGGCAT	AATGTGCTAT	CAAGACCAGG	AAACAATGAA	TGTCCATGGG	8220

GACATTTCATG	TCCAGATGGA	TGTATAACAG	GAGTATATAC	TGATGCATAT	CCACTCAATC	8280
CCACAGGGAG	CATTGTGTCA	TCTGTCATAT	TAGACTCACA	AAAATCGAGA	GTGAACCCAG	8340
TCATAACTTA	CTCAACAGCA	ACCGAAAGAG	TAAACGAGCT	GGCCATCCTA	AACAGAACAC	8400
TCTCAGCTGG	ATATACAACA	ACAAGCTGCA	TTACACACTA	TAACAAAGGA	TATTGTTTTTC	8460
ATATAGTAGA	AATAAATCAT	AAAAGCTTAA	ACACATTTCA	ACCCATGTTG	TTCAAAACAG	8520
AGATTCCAAA	AAGCTGCAGT	TAATCATAAT	TAACCATAAT	ATGCATCAAT	CTATCTATAA	8580
TACAAGTATA	TGATAAGTAA	TCAGCAATCA	GACAATAGAC	AAAAGGGAAA	TATAAAAAAC	8640
TTAGGAGCAA	AGCGTGCTCG	GGAAATGGAC	ACTGAATCTA	ACAATGGCAC	TGTATCTGAC	8700
ATACTCTATC	CTGAGTGTCA	CCTTAACTCT	CCTATCGTTA	AAGGTAAAAT	AGCACAATTA	8760
CACACTATTA	TGAGTCTACC	TCAGCCTTAT	GATATGGATG	ACGACTCAAT	ACTAGTTATC	8820
ACTAGACAGA	AAATAAAACT	TAATAAATTG	GATAAAAGAC	AACGATCTAT	TAGAAGATTA	8880
AAATTAATAT	TAAGTAAAA	AGTGAATGAC	TTAGGAAAAAT	ACACATTTAT	CAGATATCCA	8940
GAAATGTCAA	AAGAAATGTT	CAAATTATAT	ATACCTGGTA	TTAACAGTAA	AGTGAAGTAA	9000
TTATTACTTA	AAGCAGATAG	AACATATAGT	CAAAATGACTG	ATGGATTAAG	AGATCTATGG	9060
ATTAATGTGC	TATCAAAATT	AGCCTCAAAA	AATGATGGAA	GCAATTATGA	TCTTAATGAA	9120
GAAATTAATA	ATATATCGAA	AGTTCACACA	ACCTATAAAT	CAGATAAATG	GTATAATCCA	9180
TTCAAAACAT	GGTTTACTAT	CAAGTATGAT	ATGAGAAGAT	TACAAAAAGC	TCGAAATGAG	9240
ATCACTTTTA	ATGTTGGGAA	GGATTATAAC	TTGTTAGAAG	ACCAGAAGAA	TTTCTTATTG	9300
ATACATCCAG	AATTGGTTTT	GATATTAGAT	AAACAAAACT	ATAATGGTTA	TCTAATTACT	9360
CCTGAATTAG	TATTGATGTA	TTGTGACGTA	GTGGAAGGCC	GATGGAATAT	AAGTGCATGT	9420
GCTAAGTTAG	ATCCAAAATT	ACAATCTATG	TATCAGAAAG	GTAATAACCT	GTGGGAAGTG	9480
ATAGATAAAT	TGTTTCCAAT	TATGGGAGAA	AAGACATTTG	ATGTGATATC	GTTATTAGAA	9540
CCACTTG CAT	TATCCTTAAT	TCAAACCTCAT	GATCCTGTTA	AACAACCTAAG	AGGAGCTTTT	9600
TTAAATCATG	TGTTATCCGA	GATGGAATTA	ATATTTGAAT	CTAGAGAATC	GATTAAGGAA	9660
TTTCTGAGTG	TAGATTACAT	TGATAAAATT	TTAGATATAT	TTAATAAGTC	TACAATAGAT	9720
GAAATAGCAG	AGATTTTCTC	TTTTTTTAGA	ACATTTGGGC	ATCCTCCATT	AGAAGCTAGT	9780
ATTGCAGCAG	AAAAGGTTAG	AAAATATATG	TATATTGGAA	AACAATTAAA	ATTTGACACT	9840
ATTAATAAAT	GTCATGCTAT	CTTCTGTACA	ATAATAATTA	ACGGATATAG	AGAGAGGCAT	9900
GGTGGACAGT	GGCCTCCTGT	GACATTACCT	GATCATGCAC	ACGAATTCAT	CATAAATGCT	9960
TACGGTTCAA	ACTCTGCGAT	ATCATATGAA	AATGCTGTTG	ATTATTACCA	GAGCTTTATA	10020
GGAATAAAAT	TCAATAAATT	CATAGAGCCT	CAGTTAGATG	AGGATTTGAC	AATTTATATG	10080
AAAGATAAAG	CATTATCTCC	AAAAAAATCA	AATTGGGACA	CAGTTTATCC	TGCATCTAAT	10140
TTACTGTACC	GTACTAACGC	ATCCAACGAA	TCACGAAGAT	TAGTTGAAGT	ATTTATAGCA	10200
GATAGTAAAT	TTGATCCTCA	TCAGATATTG	GATTATGTAG	AATCTGGGGA	CTGGTTAGAT	10260
GATCCAGAAT	TTAATATTTT	TTATAGTCTT	AAAGAAAAAG	AGATCAAACA	GGAAGGTAGA	10320

CTCTTTGCAA	AAATGACATA	CAAAATGAGA	GCTACACAAG	TTTTATCAGA	GACCCTACTT	10380
GCAAATAACA	TAGGAAAATT	CTTTCAAGAA	AATGGGATGG	TGAAGGGAGA	GATTGAATTA	10440
CTTAAGAGAT	TAACAACCAT	ATCAATATCA	GGAGTTCCAC	GGTATAATGA	AGTGTACAAT	10500
AATTCTAAAA	GCCATACAGA	TGACCTTAAA	ACCTACAATA	AAATAAGTAA	TCTTAATTTG	10560
TCTTCTAATC	AGAAATCAAA	GAAATTTGAA	TTCAAGTCAA	CGGATATCTA	CAATGATGGA	10620
TACGAGACTG	TGAGCTGTTT	CCTAACAACA	GATCTCAAAA	AATACTGTCT	TAATTGGAGA	10680
TATGAATCAA	CAGCTCTATT	TGGAGAAACT	TGCAACCAAA	TATTTGGATT	AAATAAATTG	10740
TTTAATTGGT	TACACCCTCG	TCTTGAAGGA	AGTACAATCT	ATGTAGGTGA	TCCTTACTGT	10800
CCTCCATCAG	ATAAAGAACA	TATATCATT	GAGGATCACC	CTGATTCTGG	TTTTTACGTT	10860
CATAACCCAA	GAGGGGGTAT	AGAAGGATTT	TGCAAAAAT	TATGGACACT	CATATCTATA	10920
AGTGCAATAC	ATCTAGCAGC	TGTTAGAATA	GGCGTGAGGG	TGACTGCAAT	GGTTCAAGGA	10980
GACAATCAAG	CTATAGCTGT	AACCACAAGA	GTACCCAACA	ATTATGACTA	CAGAGTTAAG	11040
AAGGAGATAG	TTTATAAAGA	TGTAGTGAGA	TTTTTTGATT	CATTAAGAGA	AGTGATGGAT	11100
GATCTAGGTC	ATGAACTTAA	ATTAAATGAA	ACGATTATAA	GTAGCAAGAT	GTTTCATATAT	11160
AGCAAAAGAA	TCTATTATGA	TGGGAGAATT	CTTCTCAAG	CTCTAAAAGC	ATTATCTAGA	11220
TGTGTCTTCT	GGTCAGAGAC	AGTAATAGAC	GAAACAAGAT	CAGCATCTTC	AAATTTGGCA	11280
ACATCATTTG	CAAAAGCAAT	TGAGAATGGT	TATTCACCTG	TTCTAGGATA	TGCATGCTCA	11340
ATTTTTAAGA	ATATTCAACA	ACTATATATT	GCCCTTGGGA	TGAATATCAA	TCCAACATA	11400
ACACAGAATA	TCAGAGATCA	GTATTTTAGG	AATCCAAATT	GGATGCAATA	TGCCTCTTTA	11460
ATACCTGCTA	GTGTTGGGGG	ATTCAATTAC	ATGGCCATGT	CAAGATGTTT	TGTAAGGAAT	11520
ATTGGTGATC	CATCAGTTGC	CGCATTGGCT	GATATTAAAA	GATTTATTAA	GGCGAATCTA	11580
TTAGACCGAA	GTGTTCTTTA	TAGGATTATG	AATCAAGAAC	CAGGTGAGTC	ATCTTTTTTG	11640
GACTGGGCTT	CAGATCCATA	TTCATGCAAT	TTACCACAAT	CTCAAAATAT	AACCACCATG	11700
ATAAAAAATA	TAACAGCAAG	GAATGTATTA	CAAGATTCAC	CAAATCCATT	ATTATCTGGA	11760
TTATTCACAA	ATACAATGAT	AGAAGAAGAT	GAAGAATTAG	CTGAGTTCCT	GATGGACAGG	11820
AAGGTAATTC	TCCCTAGAGT	TGCACATGAT	ATTCTAGATA	ATTCTCTCAC	AGGAATTAGA	11880
AATGCCATAG	CTGGAATGTT	AGATACGACA	AAATCACTAA	TTCGGGTTGG	CATAAATAGA	11940
GGAGGACTGA	CATATAGTTT	GTTGAGGAAA	ATCAGTAATT	ACGATCTAGT	ACAATATGAA	12000
ACACTAAGTA	GGACTTTGCG	ACTAATTGTA	AGTGATAAAA	TCAAGTATGA	AGATATGTGT	12060
TCGGTAGACC	TTGCCATAGC	ATTGCGACAA	AAGATGTGGA	TTCATTTATC	AGGAGGAAGG	12120
ATGATAAGTG	GACTTGAAAC	GCCTGACCCA	TTAGAATTAC	TATCTGGGGT	AGTAATAACA	12180
GGATCAGAAC	ATTGTAAAAT	ATGTTATTCT	TCAGATGGCA	CAAACCCATA	TACTTGATG	12240
TATTTACCCG	GTAATATCAA	AATAGGATCA	GCAGAAACAG	GTATATCGTC	ATTAAGAGTT	12300
CCTTATTTTG	GATCAGTCAC	TGATGAAAGA	TCTGAAGCAC	AATTAGGATA	TATCAAGAAT	12360
CTTAGTAAAC	CTGCAAAAGC	CGCAATAAGA	ATAGCAATGA	TATATACATG	GGCATTGGT	12420

AATGATGAGA	TATCTTGGAT	GGAAGCCTCA	CAGATAGCAC	AAACACGTGC	AAATTTTACA	12480
CTAGATAGTC	TCAAAATTTT	AACACCGGTA	GCTACATCAA	CAAATTTATC	ACACAGATTA	12540
AAGGATACTG	CAACTCAGAT	GAAATTCTCC	AGTACATCAT	TGATCAGAGT	CAGCAGATTC	12600
ATAACAATGT	CCAATGATAA	CATGTCTATC	AAAGAAGCTA	ATGAAACCAA	AGATACTAAT	12660
CTTATTTATC	AACAAATAAT	GTTAACAGGA	TTAAGTGTTT	TCGAATATTT	ATTTAGATTA	12720
AAAGAAACCA	CAGGACACAA	CCCTATAGTT	ATGCATCTGC	ACATAGAAGA	TGAGTGTTGT	12780
ATTAAAGAAA	GTTTTAATGA	TGAACATATT	AATCCAGAGT	CTACATTAGA	ATTAATTCGA	12840
TATCCTGAAA	GTAATGAATT	TATTTATGAT	AAAGACCCAC	TCAAAGATGT	GGACTTATCA	12900
AAACTTATGG	TTATTAAAGA	CCATTCTTAC	ACAATTGATA	TGAATTATTG	GGATGATACT	12960
GACATCATAC	ATGCAATTTT	AATATGTACT	GCAATTACAA	TAGCAGATAC	TATGTCACAA	13020
TTAGATCGAG	ATAATTTAAA	AGAGATAATA	GTTATTGCAA	ATGATGATGA	TATTAATAGC	13080
TTAATCACTG	AATTTTTGAC	TCTTGACATA	CTTGTATTTT	TCAAGACATT	TGGTGGATTA	13140
TTAGTAAATC	AATTTGCATA	CACTCTTTAT	AGTCTAAAAA	TAGAAGGTAG	GGATCTCATT	13200
TGGGATTATA	TAATGAGAAC	ACTGAGAGAT	ACTTCCCATT	CAATATTAAA	AGTATTATCT	13260
AATGCATTAT	CTCATCCTAA	AGTATTCAAG	AGGTTCTGGG	ATTGTGGAGT	TTTAAACCTT	13320
ATTTATGGTC	CTAATACTGC	TAGTCAAGAC	CAGATAAAAC	TTGCCCTATC	TATATGTGAA	13380
TATTCAC TAG	ATCTATTTAT	GAGAGAATGG	TTGAATGGTG	TATCACTTGA	AATATACATT	13440
TGTGACAGCG	ATATGGAAGT	TGCAAATGAT	AGGAAACAAG	CCTTTATTTT	TAGACACCTT	13500
TCATTTGTTT	GTTGTTTAGC	AGAAATTGCA	TCTTTCGGAC	CTAACCTGTT	AAACTTAACA	13560
TACTTGAGAG	GACTTGATCT	ATTGAAACAA	TATCTTGAAT	TAAATATTAA	AGAAGACCCT	13620
ACTCTTAAAT	ATGTACAAAT	ATCTGGATTA	TTAATTAAAT	CGTTCCCATC	AACTGTAACA	13680
TACGTAAGAA	AGACTGCAAT	CAAATATCTA	AGGATTCGCG	GTATTAGTCC	ACCTGAGGTA	13740
ATTGATGATT	GGGATCCGGT	AGAAGATGAA	AATATGCTGG	ATAACATTGT	CAAAACTATA	13800
AATGATAACT	GTAATAAAGA	TAATAAAGGG	AATAAAATTA	ACAATTTCTG	GGGACTAGCA	13860
CTTAAGAACT	ATCAAGTCCT	TAAATCAGA	TCTATAACAA	GTGATTCTGA	TGATAATGAT	13920
AGACTAGATG	CTAATACAAG	TGGTTTGACA	CTTCCTCAAG	GAGGGAATTA	TCTATCGCAT	13980
CAATTGAGAT	TATTCGGAAT	CAACAGCACT	AGTTGTCTGA	AAGCTCTTGA	GTTATCACAA	14040
ATTTTAATGA	AGGAAGTCAA	TAAAGACAAG	GACAGGCTCT	TCCTGGGAGA	AGGAGCAGGA	14100
GCTATGCTAG	CATGTTATGA	TGCCACATTA	GGACCTGCAG	TTAATTATTA	TAATTCAGGT	14160
TTGAATATAA	CAGATGTAAT	TGGTCAACGA	GAATTGAAAA	TATTTCTTTC	AGAGGTATCA	14220
TTAGTAGGTA	AAAAATTAGG	AAATGTGACA	CAGATTCTTA	ACAGGGTAAA	AGTACTGTTC	14280
AATGGGAATC	CTAATTCAAC	ATGGATAGGA	AATATGGAAT	GTGAGAGCTT	AATATGGAGT	14340
GAATTAAATG	ATAAGTCCAT	TGGATTAGTA	CATTGTGATA	TGGAAGGAGC	TATCGGTAAA	14400
TCAGAAGAAA	CTGTTCTACA	TGAACATTAT	AGTGTTATAA	GAATTACATA	CTTGATTGGG	14460
GATGATGATG	TTGTTTTAGT	TTCCAAAATT	ATACCTACAA	TCACTCCGAA	TTGGTCTAGA	14520

155

ATACTTTATC TATATAAATT ATATTGGAAA GATGTAAGTA TAATATCACT CAAAACCTTCT 14580
 AATCCTGCAT CAACAGAATT ATATCTAATT TCGAAAGATG CATATTGTAC TATAATGGAA 14640
 CCTAGTGAAA TTGTTTTATC AAAACTTAAA AGATTGTCAC TCTTGAAGA AAATAATCTA 14700
 TTAAAATGGA TCATTTTATC AAAGAAGAGG AATAATGAAT GGTTACATCA TGAAATCAAA 14760
 GAAGGAGAAA GAGATTATGG AATCATGAGA CCATATCATA TGGCACTACA AATCTTTGGA 14820
 TTTCAAATCA ATTTAAATCA TCTGGCGAAA GAATTTTAT CAACCCAGAG TCTGACTAAT 14880
 ATCAACAATA TAATCCAAAG TTTTCAGCGA ACAATAAAGG ATGTTTTATT TGAATGGATT 14940
 AATATAACTC ATGATGATAA GAGACATAAA TTAGGCGGAA GATATAACAT ATTCCCCTG 15000
 AAAAATAAGG GAAAGTTAAG ACTGCTATCG AGAAGACTAG TATTAAGTTG GATTTCATTA 15060
 TCATTATCGA CTCGATTACT TACAGGTCGC TTTCCTGATG AAAAATTGGA ACATAGAGCA 15120
 CAGACTGGAT ATGTATCATT AGCTGATACT GATTTAGAAT CATTAAAGTT ATTGTGCGAA 15180
 AACATCATTA AGAATTACAG AGAGTGTATA GGATCAATAT CATATTGGTT TCTAACCAAA 15240
 GAAGTTAAAA TACTTATGAA ATTGATCGGT GGTGCTAAAT TATTAGGAAT TCCAGACAA 15300
 TATAAGAAC CCGAAGACCA GTTATTAGAA AACTACAATC AACATGATGA ATTTGATATC 15360
 GATTAATAACA TAAATACAAT GAAGATATAT CCTAACCTTT ATCTTTAAGC CTAGGAATAG 15420
 AAAAAAGTA AGAAAAACAT GTAATATATA TATACCAAAC AGAGTTCTTC TCTGTTTGG 15480
 TGGGTCGGCA TGGCATCTCC ACCTCCTCGC GGTCCGGACC TGGGCATCCG AAGGAGGACG 15540
 CACGTCCACT CGGATGGCTA AGGGAGAGCC TGCAGTAGCA TAACCCCTTG GGGCCTCTAA 15600
 ACGGGTCTTG AGGGGTTTTT TGCTGAAAGG AGGAACTATA TACGCGTCGA CGGGCCCCGC 15660
 GCTCAC 15666

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGATTTGCGC GCAATTAAA TCATCTGG

28

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 70 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCCAGGTCGG ACCGCGAGGA GGTGGAGATG CCATGCCAGC CCACCAAAC AAGAGAAGAA 60
CTCTGTTTGG 70

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GGCCCCGTCGA CGCGTAATAC GACTCACTAT AGGACCAAAC AAGAG 45

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGGCATCACG TGCTAC 16

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6843 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATGTTGAGCC TATTTGATAC ATTTAATGCA CGTAGGCAAG AAAACATAAC AAAATCAGCC 60
GGTGGAGCTA TCATTCCTGG ACAGAAAAAT ACTGTCTCTA TATTCGCCCT TGGACCGACA 120
ATAACTGATG ATAATGAGAA AATGACATTA GCTCTTCTAT TTCTATCTCA TTCACTAGAT 180
AATGAGAAAC AACATGCACA AAGGGCAGGG TTCTTGGTGT CTTTATTGTC AATGGCTTAT 240
GCCAATCCAG AGCTCTACCT AACAACAAAT GGAAGTAATG CAGATGTCAA GTATGTCATA 300
TACATGATTG AGAAAGATCT AAAACGGCAA AAGTATGGAG GATTGTGGT TAAGACGAGA 360
GAGATGATAT ATGAAAAGAC AACTGATTGG ATATTTGGAA GTGACCTGGA TTATGATCAG 420
GAAACTATGT TGCAGAACGG CAGGAACAAT TCAACAATTG AAGACCTTGT CCACACATTT 480
GGGTATCCAT CATGTTTtagg AGCTCTTATA ATACAGATCT GGATAGTTCT GGTCAAAGCT 540

ATCACTAGTA	TCTCAGGGTT	AAGAAAAGGC	TTTTTCACCC	GATTGGAAGC	TTTCAGACAA	600
GATGGAACAG	TGCAGGCAGG	GCTGGTATTG	AGCGGTGACA	CAGTGGATCA	GATTGGGTCA	660
ATCATGCGGT	CTCAACAGAG	CTTGGAACCT	CTTATGGTTG	AAACATTAAT	AACAATGAAT	720
ACCAGCAGAA	ATGACCTCAC	AACCATAGAA	AAGAATATAC	AAATTGTTGG	CAACTACATA	780
AGAGATGCAG	GTCTCGCTTC	ATTCTTCAAT	ACAATCAGAT	ATGGAATTGA	GACCAGAATG	840
GCAGCTTTGA	CTCTATCCAC	TCTCAGACCA	GATATCAATA	GATTAAAAGC	TTTGATGGAA	900
CTGTATTTAT	CAAAGGGACC	ACGCGCTCCT	TTCATCTGTA	TCCTCAGAGA	TCCTATACAT	960
GGTGAGTTTC	CACCAGGCAA	CTATCCTGCC	ATATGGAGCT	ATGCAATGGG	GGTGGCAGTT	1020
GTACAAAATA	GAGCCATGCA	ACAGTATGTG	ACGGGAAGAT	CATATCTAGA	CATTGATATG	1080
TTCCAGCTAG	GACAAGCAGT	AGCACGTGAT	GCCGAAGCTC	AAATGAGCTC	AACACTGGAA	1140
GATGAACTTG	GAGTGACACA	CGAATCTAAA	GAAAGCTTGA	AGAGACATAT	AAGGAACATA	1200
AACAGTTCAG	AGACATCTTT	CCACAAACCG	ACAGGTGGAT	CAGCCATAGA	GATGGCAATA	1260
GATGAAGAGC	CAGAACAATT	CGAACATAGA	GCAGATCAAG	AACAAAATGG	AGAACCTCAA	1320
TCATCCATAA	TTCAATATGC	CTGGGCAGAA	GGAAATAGAA	GCGATGATCA	GACTGAGCAA	1380
GCTACAGAAT	CTGACAATAT	CAAGACCGAA	CAACAAAACA	TCAGAGACAG	ACTAAACAAG	1440
AGACTCAACG	ACAAGAAGAA	ACAAAGCAGT	CAACCACCCA	CTAATCCCAC	AAACAGAACA	1500
AACCAGGACG	AAATAGATGA	TCTGTTTAAC	GCATTTGGAA	GCAACTAAGT	CGACGATCCG	1560
GCTGCTAACA	AAGCCCGAAA	GGAAGCTGAG	TTGGCTGCTG	CCACCGCTGA	GCAATAACTA	1620
GCATAACCCC	TTGGGGCCTC	TAAACGGGTC	TTGAGGGGTT	TTTGCTGAA	AGGAGGAACT	1680
ATATCCGGAT	CGAGATCAAT	TCTGTGAGCG	TATGGCAAAC	GAAGGAAAAA	TAGTTATAGT	1740
AGCCGCACTC	GATGGGACAT	TTCAACGTAA	ACCGTTTAAT	AATATTTTGA	ATCTTATTCC	1800
ATTATCTGAA	ATGGTGGTAA	AACTAACTGC	TGTGTGTATG	AAATGCTTTA	AGGAGGCTTC	1860
CTTTTCTAAA	CGATTGGGTG	AGGAAACCGA	GATAGAAATA	ATAGGAGGTA	ATGATATGTA	1920
TCAATCGGTG	TGTAGAAAGT	GTTACATCGA	CTCATAATAT	TATATTTTTT	ATCTAAAAAA	1980
CTAAAAATAA	ACATTGATTA	AATTTTAATA	TAATACTTAA	AAATGGATGT	TGTGTCGTTA	2040
GATAAACCGT	TTATGTATTT	TGAGGAAATT	GATAATGAGT	TAGATTACGA	ACCAGAAAGT	2100
GCAATGAGG	TCGCAAAAAA	ACTGCCGTAT	CAAGGACAGT	TAAACTATT	ACTAGGAGAA	2160
TTATTTTTTC	TTAGTAAGTT	ACAGCGACAC	GGTATATTAG	ATGGTGCCAC	CGTAGTGTAT	2220
ATAGGATCTG	CTCCCGGTAC	ACATATACGT	TATTTGAGAG	ATCATTTCTA	TAATTTAGGA	2280
GTGATCATCA	AATGGATGCT	AATTGACGGC	CGCCATCATG	ATCCTATTTT	AAATGGATTG	2340
CGTGATGTGA	CTCTAGTGAC	TCGGTTCGTT	GATGAGGAAT	ATCTACGATC	CATCAAAAAA	2400
CAACTGCATC	CTTCTAAGAT	TATTTTAATT	TCTGATGTGA	GATCCAAACG	AGGAGGAAAT	2460
GAACCTAGTA	CGGCGGATTT	ACTAAGTAAT	TACGCTCTAC	AAAATGTCAT	GATTAGTATT	2520
TTAAACCCCG	TGGCGTCTAG	TCTTAAATGG	AGATGCCCGT	TTCCAGATCA	ATGGATCAAG	2580
GACTTTTATA	TCCACACCGG	TAATAAAATG	TTACAACCTT	TTGCTCCTTC	ATATTCAGGG	2640

CCGTCGTTTT	ACAACGTCGT	GACTGGGAAA	ACCCTGGCGT	TACCCAACTT	AATCGCCTTG	2700
CAGCACATCC	CCCTTTTCGCC	AGCTGGCGTA	ATAGCGAAGA	GGCCCGCACC	GATCGCCCTT	2760
CCCAACAGTT	GCGCAGCCTG	AATGGCGAAT	GGCGCGACGC	GCCCTGTAGC	GGCGCATTAA	2820
GCGCGGCGGG	TGTGGTGGTT	ACGCGCAGCG	TGACCGCTAC	ACTTGCCAGC	GCCCTAGCGC	2880
CCGCTCCTTT	CGCTTTCTTC	CCTTCCTTTC	TCGCCACGTT	CGCCGGCTTT	CCCCGTCAAG	2940
CTCTAAATCG	GGGGCTCCCT	TTAGGGTTCC	GATTTAGTGC	TTTACGGCAC	CTCGACCCCA	3000
AAAAACTTGA	TTAGGGTGAT	GGTTCACGTA	GTGGGCCATC	GCCCTGATAG	ACGGTTTTTC	3060
GCCCTTTGAC	GTTGGAGTCC	ACGTTCTTTA	ATAGTGGACT	CTTGTTCCAA	ACTGGAACAA	3120
CACTCAACCC	TATCTCGGTC	TATTCTTTTG	ATTTATAAGG	GATTTTGCCG	ATTTCGGCCT	3180
ATTGGTTAAA	AAATGAGCTG	ATTTAACAAA	AATTTAACGC	GAATTTTAAC	AAAATATTAA	3240
CGTTTACAAT	TTCCAGGTG	GCACTTTTCG	GGGAAATGTG	CGCGGAACCC	CTATTTGTTT	3300
ATTTTCTAA	ATACATTCAA	ATATGTATCC	GCTCATGAGA	CAATAACCCT	GATAAATGCT	3360
TCAATAATAT	TGAAAAAGGA	AGAGTATGAG	TATTCAACAT	TTCCGTGTG	CCCTTATTCC	3420
CTTTTTTGCG	GCATTTTGCC	TTCCTGTTTT	TGCTCACCCA	GAAACGCTGG	TGAAAGTAAA	3480
AGATGCTGAA	GATCAGTTGG	GTGCACGAGT	GGGTTACATC	GAAGTGGATC	TCAACAGCGG	3540
TAAGATCCTT	GAGAGTTTTT	GCCCCGAAGA	ACGTTTTCCA	ATGATGAGCA	CTTTTAAAGT	3600
TCTGCTATGT	GGCGCGGTAT	TATCCCGTAT	TGACGCCGGG	CAAGAGCAAC	TCGGTCGCCG	3660
CATACACTAT	TCTCAGAATG	ACTTGTTGA	GTACTCACCA	GTCACAGAAA	AGCATCTTAC	3720
GGATGGCATG	ACAGTAAGAG	AATTATGCAG	TGCTGCCATA	ACCATGAGTG	ATAACACTGC	3780
GGCCAACTTA	CTTCTGACAA	CGATCGGAGG	ACCGAAGGAG	CTAACCCTT	TTTTGCACAA	3840
CATGGGGGAT	CATGTAACTC	GCCTTGATCG	TTGGGAACCG	GAGCTGAATG	AAGCCATACC	3900
AAACGACGAG	CGTGACACCA	CGATGCCTGT	AGCAATGGCA	ACAACGTTGC	GCAAACTATT	3960
AACTGGCGAA	CTACTTACTC	TAGCTTCCCG	GCAACAATTA	ATAGACTGGA	TGGAGGCGGA	4020
TAAAGTTGCA	GGACCACTTC	TGCGCTCGGC	CCTTCCGGCT	GGCTGGTTTA	TTGCTGATAA	4080
ATCTGGAGCC	GGTGAGCGTG	GGTCTCGCGG	TATCATTGCA	GCACTGGGGC	CAGATGGTAA	4140
GCCCTCCCGT	ATCGTAGTTA	TCTACACGAC	GGGAGTCAG	GCAACTATGG	ATGAACGAAA	4200
TAGACAGATC	GCTGAGATAG	GTGCCTCACT	GATTAAGCAT	TGGTAACTGT	CAGACCAAGT	4260
TTACTCATAT	ATACTTTAGA	TTGATTTAAA	ACTTCATTTT	TAATTTAAAA	GGATCTAGGT	4320
GAAGATCCTT	TTTGATAATC	TCATGACCAA	AATCCCTTAA	CGTGAGTTTT	CGTTCCACTG	4380
AGCGTCAGAC	CCCGTAGAAA	AGATCAAAGG	ATCTTCTTGA	GATCCTTTTT	TTCTGCGCGT	4440
AATCTGCTGC	TTGCAAACAA	AAAAACCACC	GCTACCAGCG	GTGGTTTGTT	TGCCGGATCA	4500
AGAGCTACCA	ACTCTTTTTT	CGAAGGTAAC	TGGCTTCAGC	AGAGCGCAGA	TACCAAATAC	4560
TGTCCTTCTA	GTGTAGCCGT	AGTTAGGCCA	CCACTTCAAG	AACTCTGTAG	CACCGCCTAC	4620
ATACCTCGCT	CTGCTAATCC	TGTTACCAGT	GGCTGCTGCC	AGTGGCGATA	AGTCGTGTCT	4680
TACCGGGTTG	GACTCAAGAC	GATAGTTACC	GGATAAGGCG	CAGCGGTCGG	GCTGAACGGG	4740

GGGTTCGTGC	ACACAGCCCA	GCTTGGAGCG	AACGACCTAC	ACCGAACTGA	GATACCTACA	4800
GCGTGAGCTA	TGAGAAAGCG	CCACGCTTCC	CGAAGGGAGA	AAGGCGGACA	GGTATCCGGT	4860
AAGCGGCAGG	GTCGGAACAG	GAGAGCGCAC	GAGGGAGCTT	CCAGGGGGAA	ACGCCTGGTA	4920
TCTTTATAGT	CCTGTCGGGT	TTCGCCACCT	CTGACTTGAG	CGTCGATTTT	TGTGATGCTC	4980
GTCAGGGGGG	CGGAGCCTAT	GGAAAAACGC	CAGCAACGCG	GCCTTTTTTAC	GGTTCCTGGC	5040
CTTTTGCTGG	CCTTTTGCTC	ACATGTTCTT	TCCTGCGTTA	TCCCCTGATT	CTGTGGATAA	5100
CCGTATTACC	GCCTTTGAGT	GAGCTGATAC	CGCTCGCCGC	AGCCGAACGA	CCGAGCGCAG	5160
CGAGTCAGTG	AGCGAGGAAG	CGGAAGAGCG	CCCAATACGC	AAACCGCCTC	TCCCCGCGCG	5220
TTGGCCGATT	CATTAATGCA	GCTGGCACGA	CAGGTTTCCC	GACTGGAAAG	CGGGCAGTGA	5280
GCGCAACGCA	ATTAATGTGA	GTTAGCTCAC	TCATTAGGCA	CCCCAGGCTT	TACACTTTAT	5340
GCTTCCGGCT	CGTATGTTGT	GTGGAATTGT	GAGCGGATAA	CAATTTTACA	CAGGAAACAG	5400
CTATGACCAT	GATTACGCCA	AGCTTTTGCG	ATCAATAAAT	GGATCACAAC	CAGTATCTCT	5460
TAACGATGTT	CTTCGCAGAT	GATGATTCAT	TTTTTAAGTA	TTTGGCTAGT	CAAGATGATG	5520
AATCTTCATT	ATCTGATATA	TTGCAAATCA	CTCAATATCT	AGACTTTCTG	TTATTATTAT	5580
TGATCCAATC	AAAAAATAAA	TTAGAAGCCG	TGGGTCATTG	TTATGAATCT	CTTTCAGAGG	5640
AATACAGACA	ATTGACAAAA	TTACAGACT	TTCAAGATTT	TAAAAAACTG	TTTAACAAGG	5700
TCCCTATTGT	TACAGATGGA	AGGGTCAAAC	TTAATAAAGG	ATATTTGTTC	GACTTTGTGA	5760
TTAGTTTGAT	GCGATTCAAA	AAAGAATCCT	CTCTAGCTAC	CACCGCAATA	GATCCTGTTA	5820
GATACATAGA	TCCTCGTCGC	AATATCGCAT	TTTCTAACGT	GATGGATATA	TTAAAGTCGA	5880
ATAAAGTGAA	CAATAATTAA	TTCTTTATTG	TCATCATGAA	CGGCGGACAT	ATTCAAGTGA	5940
TAATCGGCCC	CATGTTTTC	GGTAAAAGTA	CAGAATTAAT	TAGACGAGTT	AGACGTTATC	6000
AAATAGCTCA	ATATAAATGC	GTGACTATAA	AATATTCTAA	CGATAATAGA	TACGGAACGG	6060
GACTATGGAC	GCATGATAAG	AATAATTTTG	AAGCATTGGA	AGCAACTAAA	CTATGTGATG	6120
TCTTGGAATC	AATTACAGAT	TTCTCCGTGA	TAGGTATCGA	TGAAGGACAG	TTCTTTCCAG	6180
ACATTGTTGA	ATTGATCTCG	ATCCCGCGAA	ATTAATACGA	CTCACTATAG	GGAGACCACA	6240
ACGGTTTCCC	TCTAGCGGGA	TCAATTCCGC	CCCTCTCCCT	CCCCCCCCC	TAACGTTACT	6300
GGCCGAAGCC	GCTTGGAATA	AGGCCGGTGT	GCGTTTGTCT	ATATGTTATT	TTCCACCATA	6360
TTGCCGTCTT	TTGGCAATGT	GAGGGCCCGG	AAACCTGGCC	CTGTCTTCTT	GACGAGCATT	6420
CCTAGGGGTC	TTTCCCCTCT	CGCCAAAGGA	ATGCAAGGTC	TGTTGAATGT	CGTGAAGGAA	6480
GCAGTTCCTC	TGGAAGCTTC	TTGAAGACAA	ACAACGTCTG	TAGCGACCCT	TTGCAGGCAG	6540
CGGAACCCCC	CACCTGGCGA	CAGGTGCCTC	TGCGGCCAAA	AGCCACGTGT	ATAAGATACA	6600
CCTGCAAAGG	CGGCACAACC	CCAGTGCCAC	GTTGTGAGTT	GGATAGTTGT	GGAAAGAGTC	6660
AAATGGCTCT	CCTCAAGCGT	ATTCAACAAG	GGGCTGAAGG	ATGCCCAGAA	GGTACCCCAT	6720
TGTATGGGAT	CTGATCTGGG	GCCTCGGTGC	ACATGCTTTA	CATGTGTTTA	GTCGAGGTGA	6780
AAAAACGTCT	AGGCCCCCCG	AACCACGGGG	ACGTGGTTTT	CCTTTGAAAA	ACACGATAAT	6840

ACC

6843

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7107 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ATGGAAAGCG ACGCTAAAAA CTATCAAATC ATGGATTCTT GGGAAGAGGA ATCAAGAGAT	60
AAATCAACTA ATATCTCCTC GGCCCTCAAC ATCATTGAAT TCATACTCAG CACCGACCCC	120
CAAGAAGACT TATCGGAAAA CGACACAATC AACACAAGAA CCCAGCAACT CAGTGCCACC	180
ATCTGTCAAC CAGAAATCAA ACCAACAGAA ACAAGTGAGA AAGATAGTGG ATCAACTGAC	240
AAAAATAGAC AGTCCGGGTC ATCACACGAA TGTACAACAG AAGCAAAAGA TAGAAATATT	300
GATCAGGAAA CTGTACAGAG AGGACCTGGG AGAAGAAGCA GCTCAGATAG TAGAGCTGAG	360
ACTGTGGTCT CTGGAGGAAT CCCAGAAGC ATCACAGATT CTAAAAATGG AACCCAAAAC	420
ACGGAGGATA TTGATCTCAA TGAAATTAGA AAGATGGATA AGGACTCTAT TGAGGGGAAA	480
ATGCGACAAT CTGCAAATGT TCCAAGCGAG ATATCAGGAA GTGATGACAT ATTTACAACA	540
GAACAAAGTA GAAACAGTGA TCATGGAAGA AGCCTGGAAT CTATCAGTAC ACCTGATACA	600
AGATCAATAA GTGTTGTTAC TGCTGCAACA CCAGATGATG AAGAAGAAAT ACTAATGAAA	660
AATAGTAGGA CAAAGAAAAG TTCTTCAACA CATCAAGAAG ATGACAAAAG AATTAAAAAA	720
GGGGGAAAAG GGAAAGACTG GTTTAAGAAA TCAAAAGATA CCGACAACCA GATACCAACA	780
TCAGACTACA GATCCACATC AAAAGGGCAG AAGAAAATCT CAAAGACAAC AACCACCAAC	840
ACCGACACAA AGGGGCAAAC AGAAATACAG ACAGAATCAT CAGAAACACA ATCCTCATCA	900
TGGAATCTCA TCATCGACAA CAACACCGAC CGGAACGAAC AGACAAGCAC AACTCCTCCA	960
ACAACAACCT CCAGATCAAC TTATACAAAA GAATCGATCC GAACAACTC TGAATCCAAA	1020
CCCAAGACAC AAAAGACAAA TGGAAAGGAA AGGAAGGATA CAGAAGAGAG CAATCGATTT	1080
ACAGAGAGGG CAATTACTCT ATTGCAGAAT CTTGGTGTAA TTCAATCCAC ATCAAAACTA	1140
GATTTATATC AAGACAAACG AGTTGTATGT GTAGCAAATG TACTAAACAA TGTAGATACT	1200
GCATCAAAGA TAGATTTCTT GGCAGGATTA GTCATAGGGG TTTCAATGGA CAACGACACA	1260
AAATTAACAC AGATACAAAA TGAAATGCTA AACCTCAAAG CAGATCTAAA GAAAATGGAC	1320
GAATCACATA GAAGATTGAT AGAAAATCAA AGAGAACAAC TGTCATTGAT CACGTCACTA	1380
ATTTCAAATC TCAAAATTAT GACTGAGAGA GGAGGAAAGA AAGACCAAAA TGAATCCAAT	1440
GAGAGAGTAT CCATGATCAA AACAAAATTG AAAGAAGAAA AGATCAAGAA GACCAGGTTT	1500
GACCCACTTA TGGAGGCACA AGGCATTGAC AAGAATATAC CCGATCTATA TCGACATGCA	1560
GGAGATACAC TAGAGAACGA TGTACAAGTT AAATCAGAGA TATTAAGTTC ATACAATGAG	1620

TCAAATGCAA	CAAGACTAAT	ACCCAAAAAA	GTGAGCAGTA	CAATGAGATC	ACTAGTTGCA	1680
GTCATCAACA	ACAGCAATCT	CTCACAAAGC	ACAAAACAAT	CATACATAAA	CGAACTCAAA	1740
CGTTGCAAAA	ATGATGAAGA	AGTATCTGAA	TTAATGGACA	TGTTCAATGA	AGATGTCAAC	1800
AATTGCCAAT	GAGTCGACGA	TCCGGCTGCT	AACAAAGCCC	GAAAGGAAGC	TGAGTTGGCT	1860
GCTGCCACCG	CTGAGCAATA	ACTAGCATAA	CCCCTTGGGG	CCTCTAAACG	GGTCTTGAGG	1920
GGTTTTTTGC	TGAAAGGAGG	AACTATATCC	GGATCGAGAT	CAATTCTGTG	AGCGTATGGC	1980
AAACGAAGGA	AAAATAGTTA	TAGTAGCCGC	ACTCGATGGG	ACATTTCAAC	GTAAACCGTT	2040
TAATAATATT	TTGAATCTTA	TTCCATTATC	TGAAATGGTG	GTAAACTAA	CTGCTGTGTG	2100
TATGAAATGC	TTTAAGGAGG	CTTCCTTTTC	TAAACGATTG	GGTGAGGAAA	CCGAGATAGA	2160
AATAATAGGA	GGTAATGATA	TGTATCAATC	GGTGTGTAGA	AAGTGTTACA	TCGACTCATA	2220
ATATTATATT	TTTTATCTAA	AAAACAAAA	ATAAACATTG	ATTAAATTTT	AATATAATAC	2280
TTAAAAATGG	ATGTTGTGTC	GTTAGATAAA	CCGTTTATGT	ATTTTGAGGA	AATTGATAAT	2340
GAGTTAGATT	ACGAACCAGA	AAGTGCAAA	GAGGTCGCAA	AAAACTGCC	GTATCAAGGA	2400
CAGTTAAAA	TATTACTAGG	AGAATTATTT	TTTCTTAGTA	AGTTACAGCG	ACACGTATA	2460
TTAGATGGTG	CCACCGTAGT	GTATATAGGA	TCTGCTCCCG	GTACACATAT	ACGTTATTTG	2520
AGAGATCATT	TCTATAATTT	AGGAGTGATC	ATCAAATGGA	TGCTAATTGA	CGGCCGCCAT	2580
CATGATCCTA	TTTTAAATGG	ATTGCGTGAT	GTGACTCTAG	TGACTCGGTT	CGTTGATGAG	2640
GAATATCTAC	GATCCATCAA	AAAACAACCT	CATCCTTCTA	AGATTATTTT	AATTTCTGAT	2700
GTGAGATCCA	AACGAGGAGG	AAATGAACCT	AGTACGGCGG	ATTTACTAAG	TAATTACGCT	2760
CTACAAAATG	TCATGATTAG	TATTTTAAAC	CCCGTGGCGT	CTAGTCTTAA	ATGGAGATGC	2820
CCGTTTCCAG	ATCAATGGAT	CAAGGACTTT	TATATCCAC	ACGGTAATAA	AATGTTACAA	2880
CCTTTTGCTC	CTTCATATTC	AGGGCCGTCG	TTTTACAACG	TCGTGACTGG	GAAAACCTG	2940
GCGTTACCCA	ACTTAATCGC	CTTGACGAC	ATCCCCCTTT	CGCCAGCTGG	CGTAATAGCG	3000
AAGAGGCCCG	CACCGATCGC	CCTTCCCAAC	AGTTGCGCAG	CCTGAATGGC	GAATGGCGCG	3060
ACGCGCCCTG	TAGCGGCGCA	TTAAGCGCGG	CGGGTGTGGT	GGTTACGCGC	AGCGTGACCG	3120
CTACACTTGC	CAGCGCCCTA	GCGCCGCTC	CTTTCGCTTT	CTTCCCTTCC	TTTCTCGCCA	3180
CGTTCGCCCG	CTTTCCCGGT	CAAGCTCTAA	ATCGGGGGCT	CCCTTTAGGG	TTCCGATTTA	3240
GTGCTTTACG	GCACCTCGAC	CCCAAAAAAC	TTGATTAGGG	TGATGGTTCA	CGTAGTGGGC	3300
CATCGCCCTG	ATAGACGGTT	TTTCGCCCTT	TGACGTTGGA	GTCCACGTTC	TTTAATAGTG	3360
GACTCTTGTT	CCAACTGGA	ACAACACTCA	ACCCTATCTC	GGTCTATTCT	TTTGATTTAT	3420
AAGGGATTTT	GCCGATTTCG	GCCTATTGGT	TAAAAATGA	GCTGATTTAA	CAAAAATTTA	3480
ACGCGAATTT	TAACAAAATA	TTAACGTTTA	CAATTTCCCA	GGTGGCACTT	TTCGGGGAAA	3540
TGTGCGCGGA	ACCCCTATTT	GTTTATTTTT	CTAAATACAT	TCAAATATGT	ATCCGCTCAT	3600
GAGACAATAA	CCCTGATAAA	TGCTTCAATA	ATATTGAAAA	AGGAAGAGTA	TGAGTATTCA	3660
ACATTTCCGT	GTCGCCCTTA	TTCCCTTTTT	TGCGGCATTT	TGCCTTCCTG	TTTTTGCTCA	3720

CCCAGAAACG	CTGGTGAAAG	TAAAAGATGC	TGAAGATCAG	TTGGGTGCAC	GAGTGGGTTA	3780
CATCGAACTG	GATCTCAACA	GCGGTAAGAT	CCTTGAGAGT	TTTCGCCCCG	AAGAACGTTT	3840
TCCAATGATG	AGCACTTTTA	AAGTTCTGCT	ATGTGGCGCG	GTATTATCCC	GTATTGACGC	3900
CGGGCAAGAG	CAACTCGGTC	GCCGCATACA	CTATTCTCAG	AATGACTTGG	TTGAGTACTC	3960
ACCAGTCACA	GAAAAGCATC	TTACGGATGG	CATGACAGTA	AGAGAATTAT	GCAGTGCTGC	4020
CATAACCATG	AGTGATAACA	CTGCGGCCAA	CTTACTTCTG	ACAACGATCG	GAGGACCGAA	4080
GGAGCTAACC	GCTTTTTTGC	ACAACATGGG	GGATCATGTA	ACTCGCCTTG	ATCGTTGGGA	4140
ACCGGAGCTG	AATGAAGCCA	TACCAAACGA	CGAGCGTGAC	ACCACGATGC	CTGTAGCAAT	4200
GGCAACAACG	TTGCGCAAAC	TATTAAC TGG	CGAACTACTT	ACTCTAGCTT	CCCGGCAACA	4260
ATTAATAGAC	TGGATGGAGG	CGGATAAAGT	TGCAGGACCA	CTTCTGCGCT	CGGCCCTTCC	4320
GGCTGGCTGG	TTTATTGCTG	ATAAATCTGG	AGCCGGTGAG	CGTGGGTCTC	GCGGTATCAT	4380
TGCAGCACTG	GGGCCAGATG	GTAAGCCCTC	CCGTATCGTA	GTTATCTACA	CGACGGGGAG	4440
TCAGGCAACT	ATGGATGAAC	GAAATAGACA	GATCGCTGAG	ATAGGTGCCT	CACTGATTAA	4500
GCATTGGTAA	CTGTCAGACC	AAGTTTACTC	ATATATACTT	TAGATTGATT	TAAAACTTCA	4560
TTTTTAATTT	AAAAGGATCT	AGGTGAAGAT	CCTTTTTGAT	AATCTCATGA	CCAAAATCCC	4620
TTAACGTGAG	TTTTCGTTCC	ACTGAGCGTC	AGACCCCGTA	GAAAAGATCA	AAGGATCTTC	4680
TTGAGATCCT	TTTTTTCTGC	GCGTAATCTG	CTGCTTGCAA	ACAAAAAAC	CACCGCTACC	4740
AGCGGTGGTT	TGTTTGCCGG	ATCAAGAGCT	ACCAACTCTT	TTTCCGAAGG	TAAGTGGCTT	4800
CAGCAGAGCG	CAGATACCAA	ATACTGTCCT	TCTAGTGTAG	CCGTAGTTAG	GCCACCACTT	4860
CAAGAACTCT	GTAGCACCGC	CTACATACCT	CGCTCTGCTA	ATCCTGTTAC	CAGTGGCTGC	4920
TGCCAGTGGC	GATAAGTCGT	GTCTTACCGG	GTTGGACTCA	AGACGATAGT	TACCGGATAA	4980
GGCGCAGCGG	TCGGGCTGAA	CGGGGGGTTC	GTGCACACAG	CCCAGCTTGG	AGCGAACGAC	5040
CTACACCGAA	CTGAGATACC	TACAGCGTGA	GCTATGAGAA	AGCGCCACGC	TTCCCGAAGG	5100
GAGAAAGGCG	GACAGGTATC	CGGTAAGCGG	CAGGGTCGGA	ACAGGAGAGC	GCACGAGGGA	5160
GCTTCCAGGG	GGAAACGCCT	GGTATCTTTA	TAGTCCTGTC	GGGTTTCGCC	ACCTCTGACT	5220
TGAGCGTCGA	TTTTTGTGAT	GCTCGTCAGG	GGGGCGGAGC	CTATGGAAAA	ACGCCAGCAA	5280
CGCGGCCTTT	TTACGGTTCC	TGGCCTTTTG	CTGGCCTTTT	GCTCACATGT	TCTTTCTGTC	5340
GTTATCCCCT	GATTCTGTGG	ATAACCGTAT	TACCGCCTTT	GAGTGAGCTG	ATACCGCTCG	5400
CCGCAGCCGA	ACGACCGAGC	GCAGCGAGTC	AGTGAGCGAG	GAAGCGGAAG	AGCGCCCAAT	5460
ACGCAAACCG	CCTCTCCCCG	CGCGTTGGCC	GATTCATTAA	TGCAGCTGGC	ACGACAGGTT	5520
TCCCGACTGG	AAAGCGGGCA	GTGAGCGCAA	CGCAATTAAT	GTGAGTTAGC	TCACTCATTA	5580
GGCACCCAG	GCTTTTAACT	TTATGCTTCC	GGCTCGTATG	TTGTGTGGAA	TTGTGAGCGG	5640
ATAACAATTT	CACACAGGAA	ACAGCTATGA	CCATGATTAC	GCCAAGCTTT	TGCGATCAAT	5700
AAATGGATCA	CAACCAGTAT	CTCTTAACGA	TGTTCTTCGC	AGATGATGAT	TCATTTTTTA	5760
AGTATTTGGC	TAGTCAAGAT	GATGAATCTT	CATTATCTGA	TATATTGCAA	ATCACTCAAT	5820

ATCTAGACTT TCTGTTATTA TTATTGATCC AATCAAAAAA TAAATTAGAA GCCGTGGGTC	5880
ATTGTTATGA ATCTCTTTCA GAGGAATACA GACAATTGAC AAAATTCACA GACTTTCAAG	5940
ATTTTAAAAA ACTGTTTAAAC AAGGTCCCTA TTGTTACAGA TGGAAGGGTC AAACCTAATA	6000
AAGGATATTT GTTCGACTTT GTGATTAGTT TGATGCGATT CAAAAAGAA TCCTCTCTAG	6060
CTACCACCGC AATAGATCCT GTTAGATACA TAGATCCTCG TCGCAATATC GCATTTTCTA	6120
ACGTGATGGA TATATTAAAG TCGAATAAAG TGAACAATAA TTAATTCTTT ATTGTCATCA	6180
TGAACGGCGG ACATATTCAG TTGATAATCG GCCCATGTT TTCAGGTAA AGTACAGAAT	6240
TAATTAGACG AGTTAGACGT TATCAAATAG CTCAATATAA ATGCGTGACT ATAAAATATT	6300
CTAACGATAA TAGATACGGA ACGGGACTAT GGACGCATGA TAAGAATAAT TTTGAAGCAT	6360
TGGAAGCAAC TAAACTATGT GATGTCTTGG AATCAATTAC AGATTCTCC GTGATAGGTA	6420
TCGATGAAGG ACAGTTCTTT CCAGACATTG TTGAATTGAT CTCGATCCCG CGAAATTAAT	6480
ACGACTCACT ATAGGGAGAC CACAACGGTT TCCCTCTAGC GGGATCAATT CCGCCCCCTCT	6540
CCCTCCCCC CCCCTAACGT TACTGGCCGA AGCCGCTTGG AATAAGGCCG GTGTGCGTTT	6600
GTCTATATGT TATTTTCCAC CATATTGCCG TCTTTTGGCA ATGTGAGGGC CCGGAAACCT	6660
GGCCCTGTCT TCTTGACGAG CATTCTTAGG GGTCTTTCCC CTCTCGCAA AGGAATGCAA	6720
GGTCTGTTGA ATGTCGTGAA GGAAGCAGTT CCTCTGGAAG CTTCTTGAAG ACAAACAACG	6780
TCTGTAGCGA CCCTTTGCAG GCAGCGGAAC CCCCCACCTG GCGACAGGTG CCTCTGCGGC	6840
CAAAGCCAC GTGTATAAGA TACACCTGCA AAGGCGGCAC AACCCAGTG CCACGTTGTG	6900
AGTTGGATAG TTGTGGAAG AGTCAAATGG CTCTCCTCAA GCGTATTCAA CAAGGGGCTG	6960
AAGGATGCCC AGAAGGTACC CCATTGTATG GGATCTGATC TGGGGCCTCG GTGCACATGC	7020
TTTACATGTG TTAGTCGAG GTTAAAAAAC GTCTAGGCCC CCCGAACCAC GGGGACGTGG	7080
TTTTCCTTTG AAAAACACGA TAATACC	7107

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 12011 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATGGACACTG AATCTAACAA TGGCACTGTA TCTGACATAC TCTATCCTGA GTGTCACCTT	60
AACTCTCCTA TCGTTAAAGG TAAAATAGCA CAATTACACA CTATTATGAG TCTACCTCAG	120
CCTTATGATA TGGATGACGA CTCAATACTA GTTATCACTA GACAGAAAAT AAAACTTAAT	180
AAATTGGATA AAAGACAACG ATCTATTAGA AGATTAAAAT TAATATTAAC TGAAAAAGTG	240
AATGACTTAG GAAAATACAC ATTTATCAGA TATCCAGAAA TGTCAAAGA AATGTTCAAA	300
TTATATATAC CTGGTATTAA CAGTAAAGTG ACTGAATTAT TACTTAAAGC AGATAGAACA	360

TATAGTCAAA	TGACTGATGG	ATTAAGAGAT	CTATGGATTA	ATGTGCTATC	AAAATTAGCC	420
TCAAAAAATG	ATGGAAGCAA	TTATGATCTT	AATGAAGAAA	TTAATAATAT	ATCGAAAGTT	480
CACACAACCT	ATAAATCAGA	TAAATGGTAT	AATCCATTCA	AAACATGGTT	TACTATCAAG	540
TATGATATGA	GAAGATTACA	AAAAGCTCGA	AATGAGATCA	CTTTTAATGT	TGGGAAGGAT	600
TATAACTTGT	TAGAAGACCA	GAAGAATTTT	TTATTGATAC	ATCCAGAATT	GGTTTTGATA	660
TTAGATAAAC	AAAACATATA	TGGTTATCTA	ATTACTCCTG	AATTAGTATT	GATGTATTGT	720
GACGTAGTCG	AAGGCCGATG	GAATATAAGT	GCATGTGCTA	AGTTAGATCC	AAAATTACAA	780
TCTATGTATC	AGAAAGGTAA	TAACCTGTGG	GAAGTGATAG	ATAAATTGTT	TCCAATTATG	840
GGAGAAAAGA	CATTTGATGT	GATATCGTTA	TTAGAACCAC	TTGCATTATC	CTTAATTCAA	900
ACTCATGATC	CTGTTAAACA	ACTAAGAGGA	GCTTTTTTAA	ATCATGTGTT	ATCCGAGATG	960
GAATTAATAT	TTGAATCTAG	AGAATCGATT	AAGGAATTTT	TGAGTGTAGA	TTACATTGAT	1020
AAAATTTTAG	ATATATTTAA	TAAGTCTACA	ATAGATGAAA	TAGCAGAGAT	TTTCTCTTTT	1080
TTTAGAACAT	TTGGGCATCC	TCCATTAGAA	GCTAGTATTG	CAGCAGAAAA	GGTTAGAAAA	1140
TATATGTATA	TTGGAACA	ATTAAAATTT	GACACTATTA	ATAAATGTCA	TGCTATCTTC	1200
TGTACAATAA	TAATTAACGG	ATATAGAGAG	AGGCATGGTG	GACAGTGGCC	TCCTGTGACA	1260
TTACCTGATC	ATGCACACGA	ATTCATCATA	AATGCTTACG	GTTCAAACCTC	TGCGATATCA	1320
TATGAAAATG	CTGTTGATTA	TTACCAGAGC	TTTATAGGAA	TAAAATTCAA	TAAATTCATA	1380
GAGCCTCAGT	TAGATGAGGA	TTTGACAATT	TATATGAAAG	ATAAAGCATT	ATCTCCAAAA	1440
AAATCAAATT	GGGACACAGT	TTATCCTGCA	TCTAATTTAC	TGTACCGTAC	TAACGCATCC	1500
AACGAATCAC	GAAGATTAGT	TGAAGTATTT	ATAGCAGATA	GTAAATTTGA	TCCTCATCAG	1560
ATATTGGATT	ATGTAGAATC	TGGGGACTGG	TTAGATGATC	CAGAATTTAA	TATTTCTTAT	1620
AGTCTTAAAG	AAAAAGAGAT	CAAACAGGAA	GGTAGACTCT	TTGCAAAAAT	GACATACAAA	1680
ATGAGAGCTA	CACAAGTTTT	ATCAGAGACA	CTACTTGCAA	ATAACATAGG	AAAATTCTTT	1740
CAAGAAAATG	GGATGGTGAA	GGGAGAGATT	GAATTACTTA	AGAGATTAAC	AACCATATCA	1800
ATATCAGGAG	TTCCACGGTA	TAATGAAGTG	TACAATAATT	CTAAAAGCCA	TACAGATGAC	1860
CTTAAAACCT	ACAATAAAAT	AAGTAATCTT	AATTTGTCTT	CTAATCAGAA	ATCAAAGAAA	1920
TTTGAATTCA	AGTCAACGGA	TATCTACAAT	GATGGATACG	AGACTGTGAG	CTGTTTCCTA	1980
ACAACAGATC	TCAAAAATA	CTGTCTTAAT	TGGAGATATG	AATCAACAGC	TCTATTTGGA	2040
GAAACTTGCA	ACCAAATATT	TGGATTAAAT	AAATTGTTTA	ATTGGTTACA	CCCTCGTCTT	2100
GAAGGAAGTA	CAATCTATGT	AGGTGATCCT	TACTGTCCTC	CATCAGATAA	AGAACATATA	2160
TCATTAGAGG	ATCACCCTGA	TTCTGGTTTT	TACGTTTATA	ACCCAAGAGG	GGGTATAGAA	2220
GGATTTTGTC	AAAAATTATG	GACACTCATA	TCTATAAGTG	CAATACATCT	AGCAGCTGTT	2280
AGAATAGGCG	TGAGGGTGAC	TGCAATGGTT	CAAGGAGACA	ATCAAGCTAT	AGCTGTAACC	2340
ACAAGAGTAC	CCAACAATTA	TGACTACAGA	GTTAAGAAGG	AGATAGTTTA	TAAAGATGTA	2400
GTGAGATTTT	TTGATTTCATT	AAGAGAAGTG	ATGGATGATC	TAGGTCATGA	ACTTAAATTA	2460

AATGAAACGA	TTATAAGTAG	CAAGATGTTC	ATATATAGCA	AAAGAATCTA	TTATGATGGG	2520
AGAATTCCTC	CTCAAGCTCT	AAAAGCATT	TCTAGATGTG	TCTTCTGGTC	AGAGACAGTA	2580
ATAGACGAAA	CAAGATCAGC	ATCTTCAAAT	TTGGCAACAT	CATTTGCAAA	AGCAATTGAG	2640
AATGGTTATT	CACCTGTTCT	AGGATATGCA	TGCTCAATTT	TTAAGAACAT	TCAACAACATA	2700
TATATTGCCC	TTGGGATGAA	TATCAATCCA	ACTATAACAC	AGAATATCAG	AGATCAGTAT	2760
TTTAGGAATC	CAAATTGGAT	GCAATATGCC	TCTTTAATAC	CTGCTAGTGT	TGGGGGATTC	2820
AATTACATGG	CCATGTCAAG	ATGTTTTGTA	AGGAATATTG	GTGATCCATC	AGTTGCCGCA	2880
TTGGCTGATA	TTAAAAGATT	TATTAAGGCG	AATCTATTAG	ACCGAAGTGT	TCTTTATAGG	2940
ATTATGAATC	AAGAACCAGG	TGAGTCATCT	TTTTTGGACT	GGGCTTCAGA	TCCATATTCA	3000
TGCAATTTAC	CACAATCTCA	AAATATAACC	ACCATGATAA	AAAATATAAC	AGCAAGGAAT	3060
GTATTACAAG	ATTACCAAAA	TCCATTATTA	TCTGGATTAT	TCACAAATAC	AATGATAGAA	3120
GAAGATGAAG	AATTAGCTGA	GTCCTGATG	GACAGGAAGG	TAATTCTCCC	TAGAGTTGCA	3180
CATGATATTC	TAGATAATTC	TCTCACAGGA	ATTAGAAATG	CCATAGCTGG	AATGTTAGAT	3240
ACGACAAAAT	CACTAATTCTG	GGTTGGCATA	AATAGAGGAG	GA CTGACATA	TAGTTTGTTG	3300
AGGAAAATCA	GTAATTACGA	TCTAGTACAA	TATGAAACAC	TAAGTAGGAC	TTTGCGACTA	3360
ATTGTAAGTG	ATAAAATCAA	GTATGAAGAT	ATGTGTTCCG	TAGACCTTGC	CATAGCATTG	3420
CGACAAAAGA	TGTGGATTCA	TTTATCAGGA	GGAAGGATGA	TAAGTGGACT	TGAAACGCCT	3480
GACCCATTAG	AATTACTATC	TGGGGTAGTA	ATAACAGGAT	CAGAACATTG	TAAAATATGT	3540
TATTCTTCAG	ATGGCACAAA	CCCATATACT	TGGATGTATT	TACCCGGTAA	TATCAAAATA	3600
GGATCAGCAG	AAACAGGTAT	ATCGTCATTA	AGAGTTCCTT	ATTTTGGATC	AGTCACTGAT	3660
GAAAGATCTG	AAGCACAATT	AGGATATATC	AAGAATCTTA	GTAAACCTGC	AAAAGCCGCA	3720
ATAAGAATAG	CAATGATATA	TACATGGGCA	TTTGGTAATG	ATGAGATATC	TTGGATGGAA	3780
GCCTCACAGA	TAGCACAAAC	ACGTGCAAAT	TTTACACTAG	ATAGTCTCAA	AATTTTAACA	3840
CCGGTAGCTA	CATCAACAAA	TTTATCACAC	AGATTAAAGG	ATACTGCAAC	TCAGATGAAA	3900
TTCTCCAGTA	CATCATTGAT	CAGAGTCAGC	AGATTCATAA	CAATGTCCAA	TGATAACATG	3960
TCTATCAAAG	AAGCTAATGA	AACCAAAGAT	ACTAATCTTA	TTTATCAACA	AATAATGTTA	4020
ACAGGATTAA	GTGTTTTCGA	ATATTTATTT	AGATTAAAAG	AAACCACAGG	ACACAACCCT	4080
ATAGTTATGC	ATCTGCACAT	AGAAGATGAG	TGTTGTATTA	AAGAAAGTTT	TAATGATGAA	4140
CATATTAATC	CAGAGTCTAC	ATTAGAATTA	ATTGATATC	CTGAAAGTAA	TGAATTTATT	4200
TATGATAAAG	ACCCACTCAA	AGATGTGGAC	TTATCAAAAC	TTATGGTTAT	TAAAGACCAT	4260
TCTTACACAA	TTGATATGAA	TTATTGGGAT	GATACTGACA	TCATACATGC	AATTTCAATA	4320
TGTACTGCAA	TTACAATAGC	AGATACTATG	TCACAATTAG	ATCGAGATAA	TTTAAAAGAG	4380
ATAATAGTTA	TTGCAAATGA	TGATGATATT	AATAGCTTAA	TCACTGAATT	TTTGACTCTT	4440
GACATACTTG	TATTTCTCAA	GACATTTGGT	GGATTATTAG	TAAATCAATT	TGCATACACT	4500
CTTTATAGTC	TAAAAATAGA	AGGTAGGGAT	CTCATTTGGG	ATTATATAAT	GAGAACACTG	4560

AGAGATACTT	CCCATTCAAT	ATTAAAAAGTA	TTATCTAATG	CATTATCTCA	TCCTAAAGTA	4620
TTCAAGAGGT	TCTGGGATTG	TGGAGTTTTA	AACCCTATTT	ATGGTCCTAA	TACTGCTAGT	4680
CAAGACCAGA	TAAAACTTGC	CCTATCTATA	TGTGAATATT	CACTAGATCT	ATTTATGAGA	4740
GAATGGTTGA	ATGGTGTATC	ACTTGAAATA	TACATTTGTG	ACAGCGATAT	GGAAGTTGCA	4800
AATGATAGGA	AACAAGCCTT	TATTTCTAGA	CACCTTTCAT	TTGTTTGTTG	TTTAGCAGAA	4860
ATTGCATCTT	TCGGACCTAA	CCTGTTA AAC	TTAACATACT	TGGAGAGACT	TGATCTATTG	4920
AAACAATATC	TTGAATTAAA	TATTAAAGAA	GACCCTACTC	TTAAATATGT	ACAAATATCT	4980
GGATTATTAA	TTAAATCGTT	CCCATCAACT	GTAACATACG	TAAGAAAGAC	TGCAATCAAA	5040
TATCTAAGGA	TTCGCGGTAT	TAGTCCACCT	GAGGTAATTG	ATGATTGGGA	TCCGGTAGAA	5100
GATGAAAATA	TGCTGGATAA	CATTGTCAAA	ACTATAAATG	ATAACTGTAA	TAAAGATAAT	5160
AAAGGGAATA	AAATTAACAA	TTTCTGGGGA	CTAGCACTTA	AGAACTATCA	AGTCCTTAAA	5220
ATCAGATCTA	TAACAAGTGA	TTCTGATGAT	AATGATAGAC	TAGATGCTAA	TACAAGTGGT	5280
TTGACACTTC	CTCAAGGAGG	GAATTATCTA	TCGCATCAAT	TGAGATTATT	CGGAATCAAC	5340
AGCACTAGTT	GTCTGAAAGC	TCTTGAGTTA	TCACAAATTT	TAATGAAGGA	AGTCAATAAA	5400
GACAAGGACA	GGCTCTTCCT	GGGAGAAGGA	GCAGGAGCTA	TGCTAGCATG	TTATGATGCC	5460
ACATTAGGAC	CTGCAGTTAA	TTATTATAAT	TCAGGTTTGA	ATATAACAGA	TGTAATTGGT	5520
CAACGAGAAT	TGAAAATATT	TCCTTCAGAG	GTATCATTAG	TAGGTAAAAA	ATTAGGAAAT	5580
GTGACACAGA	TTCTTAACAG	GGTAAAAGTA	CTGTTCAATG	GGAATCCTAA	TTCAACATGG	5640
ATAGGAAATA	TGGAATGTGA	GAGCTTAATA	TGGAGTGAAT	TAAATGATAA	GTCCATTGGA	5700
TTAGTACATT	GTGATATGGA	AGGAGCTATC	GGTAAATCAG	AAGAAACTGT	TCTACATGAA	5760
CATTATAGTG	TTATAAGAAT	TACATACTTG	ATTGGGGATG	ATGATGTTGT	TTTAGTTTCC	5820
AAAATTATAC	CTACAATCAC	TCCGAATTGG	TCTAGAATAC	TTTATCTATA	TAAATTATAT	5880
TGGAAAGATG	TAAGTATAAT	ATCACTCAAA	ACTTCTAATC	CTGCATCAAC	AGAATTATAT	5940
CTAATTTTCGA	AAGATGCATA	TTGTACTATA	ATGGAACCTA	GTGAAATTGT	TTTATCAAAA	6000
CTTAAAAGAT	TGTCACTCTT	GGAAGAAAAT	AATCTATTAA	AATGGATCAT	TTTATCAAAG	6060
AAGAGGAATA	ATGAATGGTT	ACATCATGAA	ATCAAAGAAG	GAGAAAGAGA	TTATGGAATC	6120
ATGAGACCAT	ATCATATGGC	ACTACAAATC	TTTGGATTTT	AAATCAATTT	AAATCATCTG	6180
GCGAAAGAAT	TTTTATCAAC	CCCAGATCTG	ACTAATATCA	ACAATATAAT	CCAAAGTTTT	6240
CAGCGAACAA	TAAAGGATGT	TTTATTTGAA	TGGATTAATA	TAACTCATGA	TGATAAGAGA	6300
CATAAATTAG	GCGGAAGATA	TAACATATTC	CCACTGAAAA	ATAAGGGAAA	GTTAAGACTG	6360
CTATCGAGAA	GACTAGTATT	AAGTTGGATT	TCATTATCAT	TATCGACTCG	ATTACTTACA	6420
GGTCGCTTTC	CTGATGAAAA	ATTTGAACAT	AGAGCACAGA	CTGGATATGT	ATCATTAGCT	6480
GATACTGATT	TAGAATCATT	AAAGTTATTG	TCGAAAAACA	TCATTAAGAA	TTACAGAGAG	6540
TGTATAGGAT	CAATATCATA	TTGGTTTCTA	ACCAAAGAAG	TTAAAATACT	TATGAAATTG	6600
ATTGGTGGTG	CTAAATTATT	AGGAATTCCC	AGACAATATA	AAGAACCCGA	AGACCAGTTA	6660

TTAGAAACT	ACAATCAACA	TGATGAATTT	GATATCGATT	AAAACATAAA	TACAATGTCG	6720
ACGATCCGGC	TGCTAACAAA	GCCCGAAAGG	AAGCTGAGTT	GGCTGCTGCC	ACCGCTGAGC	6780
AATAACTAGC	ATAACCCCTT	GGGGCCTCTA	AACGGGTCTT	GAGGGGTTTT	TTGCTGAAAG	6840
GAGGAACTAT	ATCCGGATCG	AGATCAATTC	TGTGAGCGTA	TGGCAAACGA	AGGAAAAATA	6900
GTTATAGTAG	CCGCACTCGA	TGGGACATTT	CAACGTAAAC	CGTTTAATAA	TATTTTGAAT	6960
CTTATTCCAT	TATCTGAAAT	GGTGGTAAAA	CTAACTGCTG	TGTGTATGAA	ATGCTTTAAG	7020
GAGGCTTCCT	TTTCTAAACG	ATTGGGTGAG	GAAACCGAGA	TAGAAATAAT	AGGAGGTAAT	7080
GATATGTATC	AATCGGTGTG	TAGAAAGTGT	TACATCGACT	CATAATATTA	TATTTTTTAT	7140
CTAAAAAAT	AAAAATAAAC	ATTGATTAAA	TTTAAATATA	ATACTTAAAA	ATGGATGTTG	7200
TGTCGTTAGA	TAAACCGTTT	ATGTATTTTG	AGGAAATTGA	TAATGAGTTA	GATTACGAAC	7260
CAGAAAGTGC	AAATGAGGTC	GCAAAAAAAC	TGCCGTATCA	AGGACAGTTA	AACTATTAC	7320
TAGGAGAATT	ATTTTTTCTT	AGTAAGTTAC	AGCGACACGG	TATATTAGAT	GGTGCCACCG	7380
TAGTGTATAT	AGGATCTGCT	CCCGGTACAC	ATATACGTTA	TTTGAGAGAT	CATTTCTATA	7440
ATTTAGGAGT	GATCATCAAA	TGGATGCTAA	TTGACGGCCG	CCATCATGAT	CCTATTTTAA	7500
ATGGATTGCG	TGATGTGACT	CTAGTGACTC	GTTTCGTTGA	TGAGGAATAT	CTACGATCCA	7560
TCAAAAAACA	ACTGCATCCT	TCTAAGATTA	TTTAAATTTC	TGATGTGAGA	TCCAAACGAG	7620
GAGGAAATGA	ACCTAGTACG	GCGGATTTAC	TAAGTAATTA	CGCTCTACAA	AATGTCATGA	7680
TTAGTATTTT	AAACCCCGTG	GCGTCTAGTC	TTAAATGGAG	ATGCCCCTTT	CCAGATCAAT	7740
GGATCAAGGA	CTTTTATATC	CCACACGGTA	ATAAAATGTT	ACAACCTTTT	GCTCCTTCAT	7800
ATTCAGGGCC	GTCGTTTTAC	AACGTCGTGA	CTGGGAAAAC	CCTGGCGTTA	CCCAACTTAA	7860
TCGCCTTGCA	GCACATCCCC	CTTTCGCCAG	CTGGCGTAAT	AGCGAAGAGG	CCCGCACCGA	7920
TCGCCCTTCC	CAACAGTTGC	GCAGCCTGAA	TGGCGAATGG	CGCGACGCGC	CCTGTAGCGG	7980
CGCATTAAGC	GCGGCGGGTG	TGGTGGTTAC	GCGCAGCGTG	ACCGCTACAC	TTGCCAGCGC	8040
CCTAGCGCCC	GCTCCTTTTC	CTTCTTCCCC	TTCCTTTCTC	GCCACGTTTC	CCGGCTTTCC	8100
CCGTCAAGCT	CTAAATCGGG	GGCTCCCTTT	AGGGTTCCGA	TTAGTGCTT	TACGGCACCT	8160
CGACCCCAAA	AACTTGATT	AGGGTGATGG	TTCACGTAGT	GGGCCATCGC	CCTGATAGAC	8220
GGTTTTTCGC	CCTTTGACGT	TGGAGTCCAC	GTTCTTTAAT	AGTGGACTCT	TGTTCCAAAC	8280
TGGAACAACA	CTCAACCCTA	TCTCGGTCTA	TTCTTTTGAT	TTATAAGGGA	TTTTGCCGAT	8340
TTCGGCCTAT	TGGTTAAAAA	ATGAGCTGAT	TTAACAAAAA	TTTAACGCGA	ATTTTAACAA	8400
AATATTAACG	TTTACAATTT	CCCAGGTGGC	ACTTTTCGGG	GAAATGTGCG	CGGAACCCCT	8460
ATTTGTTTAT	TTTTCTAAAT	ACATTCAAAT	ATGTATCCGC	TCATGAGACA	ATAACCCTGA	8520
TAAATGCTTC	AATAATATTG	AAAAAGGAAG	AGTATGAGTA	TTCAACATTT	CCGTGTCGCC	8580
CTTATTCCTT	TTTTTGCGGC	ATTTTGCTTT	CCTGTTTTTG	CTCACCAGCA	AACGCTGGTG	8640
AAAGTAAAAG	ATGCTGAAGA	TCAGTTGGGT	GCACGAGTGG	GTTACATCGA	ACTGGATCTC	8700
AACAGCGGTA	AGATCCTTGA	GAGTTTTTCGC	CCCGAAGAAC	GTTTTCCAAT	GATGAGCACT	8760

TTTAAAGTTC	TGCTATGTGG	CGCGGTATTA	TCCCGTATTG	ACGCCGGGCA	AGAGCAACTC	8820
GGTCGCCGCA	TACACTATTC	TCAGAAATGAC	TTGGTTGAGT	ACTCACCAGT	CACAGAAAAG	8880
CATCTTACGG	ATGGCATGAC	AGTAAGAGAA	TTATGCAGTG	CTGCCATAAC	CATGAGTGAT	8940
AACACTGCGG	CCAACTTACT	TCTGACAACG	ATCGGAGGAC	CGAAGGAGCT	AACCGCTTTT	9000
TTGCACAACA	TGGGGGATCA	TGTAACTCGC	CTTGATCGTT	GGGAACCGGA	GCTGAATGAA	9060
GCCATACCAA	ACGACGAGCG	TGACACCACG	ATGCCTGTAG	CAATGGCAAC	AACGTTGCGC	9120
AAACTATTAA	CTGGCGAACT	ACTTACTCTA	GCTTCCCGGC	AACAATTAAT	AGACTGGATG	9180
GAGGCGGATA	AAGTTGCAGG	ACCACTTCTG	CGCTCGGCCC	TTCCGGCTGG	CTGTTTATT	9240
GCTGATAAAT	CTGGAGCCGG	TGAGCGTGGG	TCTCGCGGTA	TCATTGCAGC	ACTGGGGCCA	9300
GATGGTAAGC	CCTCCCGTAT	CGTAGTTATC	TACACGACGG	GGAGTCAGGC	AACTATGGAT	9360
GAACGAAATA	GACAGATCGC	TGAGATAGGT	GCCTCACTGA	TTAAGCATTG	GTAAGTGTCA	9420
GACCAAGTTT	ACTCATATAT	ACTTTAGATT	GATTTAAAAC	TTCATTTTTA	ATTTAAAAGG	9480
ATCTAGGTGA	AGATCCTTTT	TGATAATCTC	ATGACCAAAA	TCCCTTAACG	TGAGTTTTCG	9540
TTCCACTGAG	CGTCAGACCC	CGTAGAAAAG	ATCAAAGGAT	CTTCTTGAGA	TCCTTTTTTT	9600
CTGCGCGTAA	TCTGCTGCTT	GCAAACAAAA	AAACCACCGC	TACCAGCGGT	GGTTTGTTTG	9660
CCGGATCAAG	AGCTACCAAC	TCTTTTTTCCG	AAGGTAAGTG	GCTTCAGCAG	AGCGCAGATA	9720
CCAAATACTG	TCCTTCTAGT	GTAGCCGTAG	TTAGGCCACC	ACTTCAAGAA	CTCTGTAGCA	9780
CCGCCTACAT	ACCTCGCTCT	GCTAATCCTG	TTACCAGTGG	CTGCTGCCAG	TGGCGATAAG	9840
TCGTGTCTTA	CCGGGTGGA	CTCAAGACGA	TAGTTACCGG	ATAAGGCGCA	GCGGTCGGGC	9900
TGAACGGGGG	GTTCTGTCAC	ACAGCCCAGC	TTGGAGCGAA	CGACCTACAC	CGAACTGAGA	9960
TACCTACAGC	GTGAGCTATG	AGAAAGCGCC	ACGCTTCCCG	AAGGGAGAAA	GGCGGACAGG	10020
TATCCGGTAA	GCGGCAGGGT	CGGAACAGGA	GAGCGCACGA	GGGAGCTTCC	AGGGGGAAAC	10080
GCCTGGTATC	TTTATAGTCC	TGTCGGGTTT	CGCCACCTCT	GACTTGAGCG	TCGATTTTTG	10140
TGATGCTCGT	CAGGGGGGCG	GAGCCTATGG	AAAAACGCCA	GCAACGCGGC	CTTTTTACGG	10200
TTCCTGGCCT	TTTGCTGGCC	TTTTGCTCAC	ATGTTCTTTC	CTGCGTTATC	CCCTGATTCT	10260
GTGGATAACC	GTATTACCGC	CTTTGAGTGA	GCTGATACCG	CTCGCCGCAG	CCGAACGACC	10320
GAGCGCAGCG	AGTCAGTGAG	CGAGGAAGCG	GAAGAGCGCC	CAATACGCAA	ACCGCCTCTC	10380
CCCGCGCGTT	GGCCGATTCA	TTAATGCAGC	TGGCACGACA	GGTTTCCCGA	CTGGAAAGCG	10440
GGCAGTGAGC	GCAACGCAAT	TAATGTGAGT	TAGCTCACTC	ATTAGGCACC	CCAGGCTTTA	10500
CACTTTATGC	TTCCGGCTCG	TATGTTGTGT	GGAATTGTGA	GCGGATAACA	ATTTACACA	10560
GGAAACAGCT	ATGACCATGA	TTACGCCAAG	CTTTTGCGAT	CAATAAATGG	ATCACAACCA	10620
GTATCTCTTA	ACGATGTTCT	TCGCAGATGA	TGATTCAATT	TTAAGTATT	TGGCTAGTCA	10680
AGATGATGAA	TCTTCATTAT	CTGATATATT	GCAAATCACT	CAATATCTAG	ACTTTCTGTT	10740
ATTATTATTG	ATCCAATCAA	AAAATAAATT	AGAAGCCGTG	GGTCATTGTT	ATGAATCTCT	10800
TTCAGAGGAA	TACAGACAAT	TGACAAAATT	CACAGACTTT	CAAGATTTTA	AAAACTGTT	10860

169

```
TAACAAGGTC CCTATTGTTA CAGATGGAAG GGTCAAACCTT AATAAAGGAT ATTTGTTCGA 10920
CTTTGTGATT AGTTTGATGC GATTCAAAAA AGAATCCTCT CTAGCTACCA CCGCAATAGA 10980
TCCTGTTAGA TACATAGATC CTCGTCGCAA TATCGCATT TCTAACGTGA TGGATATATT 11040
AAAGTCGAAT AAAGTGAACA ATAATTAATT CTTTATTGTC ATCATGAACG GCGGACATAT 11100
TCAGTTGATA ATCGGCCCCA TGTTTTTCAGG TAAAAGTACA GAATTAATTA GACGAGTTAG 11160
ACGTTATCAA ATAGCTCAAT ATAAATGCGT GACTATAAAA TATTCTAACG ATAATAGATA 11220
CGGAACGGGA CTATGGACGC ATGATAAGAA TAATTTTGAA GCATTGGAAG CAACTAAACT 11280
ATGTGATGTC TTGGAATCAA TTACAGATTT CTCCGTGATA GGTATCGATG AAGGACAGTT 11340
CTTTCCAGAC ATTGTTGAAT TGATCTCGAT CCCGCGAAAT TAATACGACT CACTATAGGG 11400
AGACCACAAC GGTTCCTCTC TAGCGGGATC AATTCCGCCC CTCTCCCTCC CCCCCCCTA 11460
ACGTTACTGG CCGAAGCCGC TTGGAATAAG GCCGGTGTGC GTTTGTCTAT ATGTTATTTT 11520
CCACCATATT GCCGTCTTTT GGCAATGTGA GGGCCCGGAA ACCTGGCCCT GTCTTCTTGA 11580
CGAGCATTC TAGGGGTCTT TCCCTCTCG CCAAAGGAAT GCAAGGCTG TTGAATGTCG 11640
TGAAGGAAGC AGTTCCTCTG GAAGCTTCTT GAAGACAAAC AACGTCTGTA GCGACCCTTT 11700
GCAGGCAGCG GAACCCCCCA CCTGGCGACA GGTGCCTCTG CGGCCAAAAG CCACGTGTAT 11760
AAGATACACC TGCAAAGGCG GCACAACCCC AGTGCCACGT TGTGAGTTGG ATAGTTGTGG 11820
AAAGAGTCAA ATGGCTCTCC TCAAGCGTAT TCAACAAGGG GCTGAAGGAT GCCCAGAAGG 11880
TACCCCATTG TATGGGATCT GATCTGGGGC CTCGGTGCAC ATGCTTTACA TGTGTTTAGT 11940
CGAGGTAAA AAACGTCTAG GCCCCCGAA CCACGGGGAC GTGGTTTTCC TTTGAAAAAC 12000
ACGATAATAC C 12011
```

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GATCGATGCT AGCCC

15

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

170

GATCGGGCTA GCATC

15

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TTACATGGCC AT

12

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TCACATGGCG AT

12

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TTTGGACTGG GC

12

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

TTTTGATTGG GC

12

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:

171

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

TGGTCCTAAT ACTG

14

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TGGGCCTAAT ATCG

14

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GCATTATCTA GATGTGTCTT CTGGTCAGAG

30

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CCTGAATTAT AATAATTAAC TGCAGGTCCT

30

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GGGAAAGAAT CCAGAGACAA GAACGG

26

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGTGAAGTTG TGGATCCATT TGATTG

26

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

CAACCTGTAA GGTACCAGCA TCCG

24

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GATATGGTGT TAGGCCTTGA TCTGTTC

27

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

CGCCATGGAA AAATCAGAGA TCCTCTTCT

29

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CTGGATCCTA ATTGGAGTTG TTACCCATGT A

31

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

AACCATGGCT GAAAAAGGGA AAA

23

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GGTGAAGCTT AAGATGTGAT TTTACATATT TTA

33

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

AAATAGGATC CCTACAGATC ATTAGATATT AAAAT

35

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:

174

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

CGCCATGGTG TTCAGTGCTT GTTG

24

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CCACAAGCTT AATTAACCAT AATATGCATC A

31

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TTCCATGGAT TTGGATTGT CTATTGGGT

29

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15462 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ACCAAACAAG AGAAGAACT TGCTTGTA TATAAATTTA ACTTAAAT AACTTAGGAT	60
TTAAGACATT GACTAGAAGG TCAAGAAAAG GGAAGCTCTAT AATTTCAAAA ATGTTGAGCC	120
TATTTGATAC ATTTAATGCA CGTAGGCAAG AAAACATAAC AAAATCAGCC GGTGGAGCTA	180
TCATTCCTGG ACAGAAAAAT ACTGTCTCTA TATTCGCCCT TGGACCGACA ATAAGTATG	240
ATAATGAGAA AATGACATTA GCTCTTCTAT TTCTATCTCA TTCACTAGAT AATGAGAAAC	300

AACATGCACA AAGGGCAGGG TTCTTGGTGT CTTTATTGTC AATGGCTTAT GCCAATCCAG	360
AGCTCTACCT AACACAAAT GGAAGTAATG CAGATGCCAA GTATGTCATA TACATGATTG	420
AGAAAGATCT AAAACGGCAA AAGTATGGAG GATTTGTGGT TAAGACGAGA GAGATGATAT	480
ATGAAAAGAC AACTGATTGG ATATTTGGAA GTGACCTGGA TTATGATCAG GAAACTATGT	540
TGCAGAACGG CAGGAACAAT TCAACAATTG AAGACCTTGT CCACACATTT GGGTATCCAT	600
CATGTTTAGG AGCTCTTATA ATACAGATCT GGATAGTTCT GGTCAAAGCT ATCACTAGTA	660
TCTCAGGGTT AAGAAAAGGC TTTTTCACCC GATTGGAAGC TTTCAGACAA GATGGAACAG	720
TGCAGGCAGG GCTGGTATTG AGCGGTGACA CAGTGGATCA GATTGGGTCA ATCATGCGGT	780
CTCAACAGAG CTTGGTAACT CTTATGGTTG AAACATTAAT AACAATGAAT ACCAGCAGAA	840
ATGACCTCAC AACCATAGAA AAGAATATAC AAATTGTTGG CAACTACATA AGAGATGCAG	900
GTCTCGCTTC ATTCTTCAAT ACAATCAGAT ATGGAATTGA GACCAGAATG GCAGCTTTGA	960
CTCTATCCAC TCTCAGACCA GATATCAATA GATTAAAAGC TTTGATGGAA CTGTATTTAT	1020
CAAAGGGACC ACGCGCTCCT TTCATCTGTA TCCTCAGAGA TCCTATACAT GGTGAGTTCTG	1080
CACCAGGCAA CTATCCTGCC ATATGGAGCT ATGCAATGGG GGTGGCAGTT GTACAAAATA	1140
GAGCCATGCA ACAGTATGTG ACGGGAAGAT CATATCTAGA CATTGATATG TTCCAGCTAG	1200
GACAAGCAGT AGCACGTGAT GCCGAAGCTC AAATGAGCTC AACACTGGAA GATGAACTTG	1260
GAGTGACACA CGAAGCTAAA GAAAGCTTGA AGAGACATAT AAGGAACATA AACAGTTCAG	1320
AGACATCTTT CCACAAACCG ACAGGTGGAT CAGCCATAGA GATGGCAATA GATGAAGAGC	1380
CAGAACAATT CGAACATAGA GCAGATCAAG AACAAAATGG AGAACCTCAA TCATCCATAA	1440
TTCAATATGC CTGGGCAGAA GGAAATAGAA GCGATGATCA GACTGAGCAA GCTACAGAAT	1500
CTGACAATAT CAAGACCGAA CAACAAAACA TCAGAGACAG ACTAAACAAG AGACTCAACG	1560
ACAAGAAGAA ACAAAGCAGT CAACCACCCA CTAATCCCAC AAACAGAACA AACCAGGACG	1620
AAATAGATGA TCTGTTTAAC GCATTTGGAA GCAACTAATC GAATCAACAT TTTAATCTAA	1680
ATCAATAATA AATAAGAAAA ACTTAGGATT AAAGAATCCT ATCATACCGG AATATAGGGT	1740
GGTAAATTTA GAGTCTGCTT GAAACTCAAT CAATAGAGAG TTGATGGAAA GCGATGCTAA	1800
AAACTATCAA ATCATGGATT CTTGGGAAGA GGAATCAAGA GATAAATCAA CTAATATCTC	1860
CTCGGCCCTC AACATCATTG AATTCATACT CAGCACCGAC CCCCAAGAAG ACTTATCGGA	1920
AAACGACACA ATCAACACAA GAACCCAGCA ACTCAGTGCC ACCATCTGTC AACCAGAAAT	1980
CAAACCAACA GAAACAAGTG AGAAAGATAG TGGATCAACT GACAAAATA GACAGTCTGG	2040
GTCATCACAC GAATGTACAA CAGAAGCAAA AGATAGAAAC ATTGATCAGG AAAGTGTACA	2100
GAGAGGACCT GGGAGAAGAA GCAGCTCAGA TAGTAGAGCT GAGACTGTGG TCTCTGGAGG	2160
AATCCCAGAG AGCATCACAG ATTCTAAAAA TGAACCCAA AACACGGAGG ATATTGATCT	2220
CAATGAAATT AGAAAGATGG ATAAGGACTC TATTGAGGGG AAAATGCGAC AATCTGCAAA	2280
TGTTCCAAGC GAGATATCAG GAAGTGATGA CATATTTACA ACAGAACAAA GTAGAAACAG	2340
TGATCATGGA AGAAGCCTGG AATCTATCAG TACACCTGAT ACAAGATCAA TAAGTGTGTG	2400

TACTGCTGCA	ACACCAGATG	ATGAAGAAGA	AATACTAATG	AAAAATAGTA	GGACAAAGAA	2460
AAGTTCTTCA	ACACATCAAG	AAGATGACAA	AAGAATTAA	AAAGGGGGAA	AAGGGAAAGA	2520
CTGGTTTAAG	AAATCAAAAAG	ATACCGACAA	CCAGATACCA	ACATCAGACT	ACAGATCCAC	2580
ATCAAAAGGG	CAGAAGAAAA	TCTCAAAGAC	AACAACCACC	AACACCGACA	CAAAGGGGCA	2640
AACAGAAATA	CAGACAGAAT	CATCAGAAAC	ACAATCCTCA	TCATGGAATC	TCATCATCGA	2700
CAACAACACC	GACCGGAACG	AACAGACAAG	CACAACCTCT	CCAACAACAA	CTTCCAGATC	2760
AACTTATACA	AAAGAATCGA	TCCGAACAAA	CTCTGAATCC	AAACCCAAGA	CACAAAAGAC	2820
AAATGGAAAG	GAAAGGAAGG	ATACAGAAGA	GAGCAATCGA	TTTACAGAGA	GGGCAATTAC	2880
TCTATTGCAG	AATCTTGGTG	TAATTCAATC	CACATCAAAA	CTAGATTTAT	ATCAAGACAA	2940
ACGAGTTGTA	TGTGTAGCAA	ATGTACTAAA	CAATGTAGAT	ACTGCATCAA	AGATAGATTT	3000
CCTGGCAGGA	TTAGTCATAG	GGGTTTCAAT	GGACAACGAC	ACAAAATTAA	CACAGATACA	3060
AAATGAAATG	CTAAACCTCA	AAGCAGATCT	AAAGAAAATG	GACGAATCAC	ATAGAAGATT	3120
GATAGAAAAT	CAAAGAGAAC	AACTGTCTAT	GATCACGTCA	CTAATTTCAA	ATCTCAAAAT	3180
TATGACTGAG	AGAGGAGGAA	AGAAAGACCA	AAATGAATCC	AATGAGAGAG	TATCCATGAT	3240
CAAAACAAAA	TTGAAAGAAG	AAAAGATCAA	GAAGACCAGG	TTTGACCCAC	TTATGGAGGC	3300
ACAAGGCATT	GACAAGAATA	TACCCGATCT	ATATCGACAT	GCAGGAGATA	CCTAGAGAA	3360
CGATGTACAA	GTAAATCAG	AGATATTAAG	TTCATACAAT	GAGTCAAATG	CAACAAGACT	3420
AATACCCAAA	AAAGTGAGCA	GTACAATGAG	ATCACTAGTT	GCAGTCATCA	ACAACAGCAA	3480
TCTCTCACAA	AGCACAAAAC	AATCATACAT	AAACGAACTC	AAACGTTGCA	AAAATGATGA	3540
AGAAGTATCT	GAATTAATGG	ACATGTTCAA	TGAAGATGTC	AACAATTGCC	AATGATCCAA	3600
CAAAGAAACG	ACACCGAACA	AACAGACAAG	AAACAACAGT	AGATCAAAAAC	CTGTCAACAC	3660
ACACAAAATC	AAGCAGAATG	AAACAACAGA	TATCAATCAA	TATACAAATA	AGAAAACTT	3720
AGGATTAAAAG	AATAAATTAA	TCCTTGTTCA	AAATGAGTAT	AACTAACTCT	GCAATATACA	3780
CATTCCCAGA	ATCATCATTC	TCTGAAAATG	GTCATATAGA	ACCATTACCA	CTCAAAGTCA	3840
ATGAACAGAG	GAAAGCAGTA	CCCCACATTA	GAGTTGCCAA	GATCGGAAAT	CCACCAAAAC	3900
ACGGATCCCG	GTATTTAGAT	GTCTTCTTAC	TCGGCTTCTT	CGAGATGGAA	CGAATCAAAG	3960
ACAAATACGG	GAGTGTGAAT	GATCTCGACA	GTGACCCGAG	TTACAAAGTT	TGTGGCTCTG	4020
GATCATTACC	AATCGGATTG	GCTAAGTACA	CTGGGAATGA	CCAGGAATTG	TTACAAGCCG	4080
CAACCAAAC	GGATATAGAA	GTGAGAAGAA	CAGTCAAAGC	GAAAGAGATG	GTTGTTTACA	4140
CGGTACAAAA	TATAAAACCA	GAAGTGTACC	CATGGTCCAA	TAGACTAAGA	AAAGGAATGC	4200
TGTTTCGATGC	CAACAAAGTT	GCTCTTGCTC	CTCAATGTCT	TCCACTAGAT	AGGAGCATAA	4260
AATTTAGAGT	AATCTTCGTG	AATTGTACGG	CAATTGGATC	AATAACCTTG	TTCAAAATTC	4320
CTAAGTCAAT	GGCATCACTA	TCGTTAACCA	ACACAATATC	AATCAATCTG	CAGGTACACA	4380
TAAAAACAGG	GGTTCAGACT	GATTCTAAAG	GGATAGTTCA	AATTTTGGAT	GAGAAAGGCG	4440
AAAAATCACT	GAATTTTCATG	GTCCATCTCG	GATTGATCAA	AAGAAAAGTA	GGCAGAATGT	4500

ACTCTGTTGA ATACTGTAAA CAGAAAATCG AGAAAATGAG ATTGATATTT TCTTTAGGAC 4560
TAGTTGGAGG AATCAGTCTT CATGTCAATG CAACTGGGTC CATATCAAAA AACTAGCAA 4620
GTCAGCTGGT ATTCAAAAGA GAGATTGTGTT ATCCTTTAAT GGATCTAAAT CCGCATCTCA 4680
ATCTAGTTAT CTGGGCTTCA TCAGTAGAGA TTACAAGAGT GGATGCAATT TTCCAACCTT 4740
CTTTACCTGG CGAGTTCAGA TACTATCCTA ATATTATTGC AAAAGGAGTT GGGAAAATCA 4800
AACAAATGGAA CTAGTAATCT CTATTTTAGT CCGGACGTAT CTATTAAGCC GAAGCAAATA 4860
AAGGATAATC AAAAAGCTAG GACAAAAGAG GTCAATACCA ACAACTATTA GCAGTCACAC 4920
TCGCAAGAAT AAGAGAGAAG GGACCAAAAA AGTCAAATAG GAGAAATCAA AACAAAAGGT 4980
ACAGAACACC AGAACAACAA AATCAAAACA TCCAATCAC TCAAAACAAA AATTCCAAAA 5040
GAGACCGGCA ACACAACAAG CACTGAACAC AATGCCAACT TCAATACTGC TAATTATTAC 5100
AACCATGATC ATGGCATCTT TCTGCCAAAT AGATATCACA AACTACAGC ACGTAGGTGT 5160
ATTGGTCAAC AGTCCCAAAG GGATGAAGAT ATCACAAAAC TTTGAAACAA GATATCTAAT 5220
TTTGAGCCTC ATACCAAAAA TAGAAGACTC TAACTCTTGT GGTGACCAAC AGATCAAGCA 5280
ATACAAGAAG TTATTGGATA GACTGATCAT CCCTTTTATAT GATGGATTAA GATTACAGAA 5340
AGATGTGATA GTAACCAATC AAGAATCCAA TGAAAACACT GATCCCAGAA CAAAACGATT 5400
CTTTGGAGGG GTAATTGGAA CCATTGCTCT GGGAGTAGCA ACCTCAGCAC AAATTACAGC 5460
GGCAGTTGCT CTGTTGAAG CCAAGCAGGC AAGATCAGAC ATCGAAAAAC TCAAAGAAGC 5520
AATTAGGGAC ACAACAAAG CAGTGCAGTC AGTTCAGAGC TCCATAGGAA ATTTAATAGT 5580
AGCAATTAA TCAGTCCAGG ATTATGTTAA CAAAGAAATC GTGCCATCGA TTGCGAGGCT 5640
AGGTTGTGAA GCAGCAGGAC TTCAATTAGG AATTGCATTA ACACAGCATT ACTCAGAATT 5700
AACAAACATA TTTGGTGATA ACATAGGATC GTTACAAGAA AAAGGAATAA AATTACAAGG 5760
TATAGCATCA TTATACCGCA CAAATATCAC AGAAATATTC ACAACATCAA CAGTTGATAA 5820
ATATGATATC TATGATCTGT TATTTACAGA ATCAATAAAG GTGAGAGTTA TAGATGTTGA 5880
CTTGAATGAT TACTCAATCA CCCTCCAAGT CAGACTCCCT TTATTAATA GGCTGCTGAA 5940
CACTCAGATC TACAAAGTAG ATTCCATATC ATATAACATC CAAAACAGAG AATGGTATAT 6000
CCCTCTTCCC AGCCATATCA TGACGAAAGG GGCATTTCTA GGTGGAGCAG ACGTCAAAGA 6060
ATGTATAGAA GCATTACAGCA GCTATATATG CCCTTCTGAT CCAGGATTTG TATTAAACCA 6120
TGAAATAGAG AGCTGCTTAT CAGGAAACAT ATCCCAATGT CCAAGAACAA CGGTCACATC 6180
AGACATTGTT CCAAGATATG CATTTGTCAA TGGAGGAGTG GTTGCAAACT GTATAACAAC 6240
CACCTGTACA TGCAACGGAA TTGGTAATAG AATCAATCAA CCACCTGATC AAGGAGTAAA 6300
AATTATAACA CATAAAGAAT GTAGTACTGT GGGTATCAAC GGAATGCTGT TCAATACAAA 6360
TAAAGAAGGA ACTCTTGCA TCTATACACC AAATGATATA AACTAAACA ATTCTGTTAC 6420
ACTGGATCCA ATTGACATAT CAATCGAGCT CAACAAGGCC AAATCAGATC TAGAAGAATC 6480
AAAAGAATGG ATAAGAAGGT CAAATCAAAA ACTAGATTCT ATTGGAAATT GGCATCAATC 6540
TAGCACTACA ATCATAATTA TTTTGATAAT GATCATTATA TTGTTTATAA TTAATATAAC 6600

GATAATTACA	ATTGCAATTA	AGTATTACAG	AATTCAAAAG	AGAAATCGAG	TGGATCAAAA	6660
TGACAAGCCA	TATGTACTAA	CAAACAAATA	ACATATCTAC	AGATCATTAG	ATATTAAAAT	6720
TATAAAAAAC	TTAGGAGTAA	AGTTACGCAA	TCCAACCTCTA	CTCATATAAT	TGAGGAAGGA	6780
CCCAATAGAC	AAATCCAAAT	TCGAGATGGA	ATACTGGAAG	CATACCAATC	ACGGAAAGGA	6840
TGCTGGTAAT	GAGCTGGAGA	CGTCTATGGC	TACTCATGGC	AACAAGCTCA	CTAATAAGAT	6900
AATATACATA	TTATGGACAA	TAATCCTGGT	GTTATTATCA	ATAGTCTTCA	TCATAGTGCT	6960
AATTAATTCC	ATCAAAAGTG	AAAAGGCCCA	CGAATCATTG	CTGCAAGACA	TAAATAATGA	7020
GTTTATGGAA	ATTACAGAAA	AGATCCAAAT	GGCATCGGAT	AATACCAATG	ATCTAATACA	7080
GTCAGGAGTG	AATACAAGGC	TTCTTACAAT	TCAGAGTCAT	GTCCAGAATT	ACATACCAAT	7140
ATCATTGACA	CAACAGATGT	CAGATCTTAG	GAAATTCATT	AGTGAAATTA	CAATTAGAAA	7200
TGATAATCAA	GAAGTGCTGC	CACAAAGAAT	AACACATGAT	GTAGGTATAA	AACCTTTAAA	7260
TCCAGATGAT	TTTTGGAGAT	GCACGTCTGG	TCTTCCATCT	TTAATGAAAA	CTCCAAAAAT	7320
AAGGTTAATG	CCAGGGCCGG	GATTATTAGC	TATGCCAACG	ACTGTTGATG	GCTGTGTTAG	7380
AACTCCGTCT	TTAGTTATAA	ATGATCTGAT	TTATGCTTAT	ACCTCAAATC	TAATTACTCG	7440
AGGTTGTCAG	GATATAGGAA	AATCATATCA	AGTCTTACAG	ATAGGGATAA	TAACTGTAAA	7500
CTCAGACTTG	GTACCTGACT	TAAATCCTAG	GATCTCTCAT	ACCTTTAACA	TAAATGACAA	7560
TAGGAAGTCA	TGTTCTCTAG	CACTCCTAAA	TATAGATGTA	TATCAACTGT	GTTCAACTCC	7620
CAAAGTTGAT	GAAAGATCAG	ATTATGCATC	ATCAGGCATA	GAAGATATTG	TACTTGATAT	7680
TGTCAATTAT	GATGGTTCAA	TCTCAACAAC	AAGATTTAAG	AATAATAACA	TAAGCTTTGA	7740
TCAACCATAT	GCTGCACTAT	ACCCATCTGT	TGGACCAGGG	ATATACTACA	AAGGCAAAAT	7800
AATATTTCTC	GGGTATGGAG	GTCTTGAACA	TCCAATAAAT	GAGAATGTAA	TCTGCAACAC	7860
AACTGGGTGC	CCCGGGAAAA	CACAGAGAGA	CTGTAATCAA	GCATCTCATA	GTACTTGGTT	7920
TTCAGATAGG	AGGATGGTCA	ACTCCATCAT	TGTGGCTGAC	AAAGGCTTAA	ACTCAATTCC	7980
AAAATTGAAA	GTATGGACGA	TATCTATGCG	ACAAAATTAC	TGGGGGTCAG	AAGGAAGGTT	8040
ACTTCTACTA	GGTAACAAGA	TCTATATATA	TACAAGATCT	ACAAGTTGGC	ATAGCAAGTT	8100
ACAATTAGGA	ATAATTGATA	TTACTGATTA	CAGTGATATA	AGGATAAAAT	GGACATGGCA	8160
TAATGTGCTA	TCAAGACCAG	GAAACAATGA	ATGTCCATGG	GGACATTCAT	GTCCAGATGG	8220
ATGTATAACA	GGAGTATATA	CTGATGCATA	TCCACTCAAT	CCCACAGGGA	GCATTGTGTC	8280
ATCTGTCATA	TTAGACTCAC	AAAAATCGAG	AGTGAACCCA	GTCATAACTT	ACTCAACAGC	8340
AACCGAAAGA	GTAAACGAGC	TGGCCATCCT	AAACAGAACA	CTCTCAGCTG	GATATACAAC	8400
AACAAGCTGC	ATTACACACT	ATAACAAAGG	ATATTGTTTT	CATATAGTAG	AAATAAATCA	8460
TAAAAGCTTA	AACACATTTT	AACCCATGTT	GTTCAAAACA	GAGATTCCAA	AAAGCTGCAG	8520
TTAATCATAA	TTAACCATAA	TATGCATCAA	TCTATCTATA	ATACAAGTAT	ATGATAAGTA	8580
ATCAGCAATC	AGACAATAGA	CAAAAGGGAA	ATATAAAAAA	CTTAGGAGCA	AAGCGTGCTC	8640
GGGAAATGGA	CACTGAATCT	AACAATGGCA	CTGTATCTGA	CATACTCTAT	CCTGAGTGTC	8700

ACCTTAACTC	TCCTATCGT	AAAGGTAAAA	TAGCACAATT	ACACACTATT	ATGAGTCTAC	8760
CTCAGCCTTA	TGATATGGAT	GACGACTCAA	TACTAGTTAT	CACTAGACAG	AAAATAAAAC	8820
TTAATAAATT	GGATAAAAGA	CAACGATCTA	TTAGAAGATT	AAAATTAATA	TTAACTGAAA	8880
AAGTGAATGA	CTTAGGAAAA	TACACATTTA	TCAGATATCC	AGAAATGTCA	AAAGAAATGT	8940
TCAAATTATA	TATACCTGGT	ATTAACAGTA	AAGTGAAGTA	ATTATTACTT	AAAGCAGATA	9000
GAACATATAG	TCAAATGACT	GATGGATTAA	GAGATCTATG	GATTAATGTG	CTATCAAAAT	9060
TAGCCTCAAA	AAATGATGGA	AGCAATTATG	ATCTTAATGA	AGAAATTAAT	AATATATCGA	9120
AAGTTCACAC	AACCTATAAA	TCAGATAAAT	GGTATAATCC	ATTCAAAACA	TGGTTTACTA	9180
TCAAGTATGA	TATGAGAAGA	TTACAAAAAG	CTCGAAATGA	GATCACTTTT	AATGTTGGGA	9240
AGGATTATAA	CTTGTTAGAA	GACCAGAAGA	ATTTCTTATT	GATACATCCA	GAATTGGTTT	9300
TGATATTAGA	TAAACAAAAC	TATAATGGTT	ATCTAATTAC	TCCTGAATTA	GTATTGATGT	9360
ATTGTGACGT	AGTCGAAGGC	CGATGGAATA	TAAGTGCATG	TGCTAAGTTA	GATCCAAAAT	9420
TACAATCTAT	GTATCAGAAA	GGTAATAACC	TGTGGGAAGT	GATAGATAAA	TTGTTTCCAA	9480
TTATGGGAGA	AAAGACATTT	GATGTGATAT	CGTTATTAGA	ACCACTTGCA	TTATCCTTAA	9540
TTCAAACCTCA	TGATCCTGTT	AAACAACATA	GAGGAGCTTT	TTTAAATCAT	GTGTTATCCG	9600
AGATGGAATT	AATATTTGAA	TCTAGAGAAT	CGATTAAAGGA	ATTTCTGAGT	GATAGATTACA	9660
TTGATAAAAT	TTTAGATATA	TTTAATAAGT	CTACAATAGA	TGAAATAGCA	GAGATTTTCT	9720
CTTTTTTTTAG	AACATTTGGG	CATCCTCCAT	TAGAAGCTAG	TATTGCAGCA	GAAAAGGTTA	9780
GAAAATATAT	GTATATTGGA	AAACAATTAA	AATTTGACAC	TATTAATAAA	TGTCATGCTA	9840
TCTTCTGTAC	AATAATAATT	AACGGATATA	GAGAGAGGCA	TGGTGGACAG	TGGCCTCCTG	9900
TGACATTACC	TGATCATGCA	CACGAATTCA	TCATAAATGC	TTACGGTTCA	AACTCTGCGA	9960
TATCATATGA	AAATGCTGTT	GATTATTACC	AGAGCTTTAT	AGGAATAAAA	TTCAATAAAT	10020
TCATAGAGCC	TCAGTTAGAT	GAGGATTGGA	CAATTTATAT	GAAAGATAAA	GCATTATCTC	10080
CAAAAAAATC	AAATTGGGAC	ACAGTTTATC	CTGCATCTAA	TTTACTGTAC	CGTACTAACG	10140
CATCCAACGA	ATCACGAAGA	TTAGTTGAAG	TATTTATAGC	AGATAGTAAA	TTTGATCCTC	10200
ATCAGATATT	GGATTATGTA	GAATCTGGGG	ACTGGTTAGA	TGATCCAGAA	TTTAATATTT	10260
CTTATAGTCT	TAAAGAAAAA	GAGATCAAAC	AGGAAGGTAG	ACTCTTTGCA	AAAATGACAT	10320
ACAAAATGAG	AGCTACACAA	GTTTTATCAG	AGACCCTACT	TGCAAATAAC	ATAGGAAAAT	10380
TCTTTCAAGA	AAATGGGATG	GTGAAGGGAG	AGATTGAATT	ACTTAAGAGA	TTAACAACCA	10440
TATCAATATC	AGGAGTTCCA	CGGTATAATG	AAGTGACAA	TAATCTAAA	AGCCATACAG	10500
ATGACCTTAA	AACCTACAAT	AAAATAAGTA	ATCTTAATTT	GTCTTCTAAT	CAGAAATCAA	10560
AGAAATTTGA	ATTCAAGTCA	ACGGATATCT	ACAATGATGG	ATACGAGACT	GTGAGCTGTT	10620
TCCTAACAAAC	AGATCTCAAA	AAATACTGTC	TTAATTGGAG	ATATGAATCA	ACAGCTCTAT	10680
TTGGAGAAAC	TTGCAACCAA	ATATTTGGAT	TAAATAAATT	GTTTAATTGG	TTACACCCCTC	10740
GTCTTGAAGG	AAGTACAATC	TATGTAGGTG	ATCCTTACTG	TCCTCCATCA	GATAAAGAAC	10800

ATATATCATT	AGAGGATCAC	CCTGATTCTG	GTTTTTACGT	TCATAACCCA	AGAGGGGGTA	10860
TAGAAGGATT	TTGTCAAAAA	TTATGGACAC	TCATATCTAT	AAGTGCAATA	CATCTAGCAG	10920
CTGTTAGAAT	AGGCGTGAGG	GTGACTGCAA	TGGTTCAAGG	AGACAATCAA	GCTATAGCTG	10980
TAACCACAAG	AGTACCCAAC	AATTATGACT	ACAGAGTTAA	GAAGGAGATA	GTTTATAAAG	11040
ATGTAGTGAG	ATTTTTTGAT	TCATTAAGAG	AAGTGATGGA	TGATCTAGGT	CATGAACTTA	11100
AATTAAATGA	AACGATTATA	AGTAGCAAGA	TGTTTCATATA	TAGCAAAAGA	ATCTATTATG	11160
ATGGGAGAAT	TCTTCCTCAA	GCTCTAAAAG	CATTATCTAG	ATGTGTCTTC	TGGTCAGAGA	11220
CAGTAATAGA	CGAAACAAGA	TCAGCATCTT	CAAATTTGGC	AACATCATTT	GCAAAAAGCAA	11280
TTGAGAATGG	TTATTCACCT	GTTCTAGGAT	ATGCATGCTC	AATTTTTTAAG	AATATTCAAC	11340
AACTATATAT	TGCCCTTGGG	ATGAATATCA	ATCCAACAT	AACACAGAAT	ATCAGAGATC	11400
AGTATTTTAG	GAATCCAAAT	TGGATGCAAT	ATGCCTCTTT	AATACCTGCT	AGTGTGGGG	11460
GATTCAATCA	CATGGCGATG	TCAAGATGTT	TTGTAAGGAA	TATTGGTGAT	CCATCAGTTG	11520
CCGCATTGGC	TGATATTAAA	AGATTTATTA	AGGCGAATCT	ATTAGACCGA	AGTGTCTCTT	11580
ATAGGATTAT	GAATCAAGAA	CCAGGTGAGT	CATCTTTTTT	TGATTGGGCT	TCAGATCCAT	11640
ATTCATGCAA	TTTACCACAA	TCTCAAAATA	TAACCACCAT	GATAAAAAAT	ATAACAGCAA	11700
GGAATGTATT	ACAAGATTCA	CCAAATCCAT	TATTATCTGG	ATTATTCACA	AATACAATGA	11760
TAGAAGAAGA	TGAAGAATTA	GCTGAGTTCC	TGATGGACAG	GAAGGTAATT	CTCCCTAGAG	11820
TTGCACATGA	TATTCTAGAT	AATTCTCTCA	CAGGAATTAG	AAATGCCATA	GCTGGAATGT	11880
TAGATACGAC	AAAATCACTA	ATTCGGGTTG	GCATAAATAG	AGGAGGACTG	ACATATAGTT	11940
TGTTGAGGAA	AATCAGTAAT	TACGATCTAG	TACAATATGA	AACACTAAGT	AGGACTTTGC	12000
GACTAATTGT	AAGTGATAAA	ATCAAGTATG	AAGATATGTG	TTCGGTAGAC	CTTGCCATAG	12060
CATTGCGACA	AAAGATGTGG	ATTCATTTAT	CAGGAGGAAG	GATGATAAGT	GGACTTGAAA	12120
CGCCTGACCC	ATTAGAATTA	CTATCTGGGG	TAGTAATAAC	AGGATCAGAA	CATTGTAAAA	12180
TATGTTATTC	TTCAGATGGC	ACAAACCCAT	ATACTTGGAT	GTATTTACCC	GGTAATATCA	12240
AAATAGGATC	AGCAGAAACA	GGTATATCGT	CATTAAGAGT	TCCTTATTTT	GGATCAGTCA	12300
CTGATGAAAG	ATCTGAAGCA	CAATTAGGAT	ATATCAAGAA	TCTTAGTAAA	CCTGCAAAAG	12360
CCGCAATAAG	AATAGCAATG	ATATATACAT	GGGCATTTGG	TAATGATGAG	ATATCTTGGA	12420
TGGAAGCCTC	ACAGATAGCA	CAAACACGTG	CAAATTTTAC	ACTAGATAGT	CTCAAAATTT	12480
TAACACCGGT	AGCTACATCA	ACAAATTTAT	CACACAGATT	AAAGGATACT	GCAACTCAGA	12540
TGAAATTCTC	CAGTACATCA	TTGATCAGAG	TCAGCAGATT	CATAACAATG	TCCAATGATA	12600
ACATGTCTAT	CAAAGAAGCT	AATGAAACCA	AAGATACTAA	TCTTATTTAT	CAACAAATAA	12660
TGTTAACAGG	ATTAAGTGTT	TTCGAATATT	TATTTAGATT	AAAAGAAACC	ACAGGACACA	12720
ACCCTATAGT	TATGCATCTG	CACATAGAAG	ATGAGTGTTG	TATTAAAGAA	AGTTTTAATG	12780
ATGAACATAT	TAATCCAGAG	TCTACATTAG	AATTAATTCG	ATATCCTGAA	AGTAATGAAT	12840
TTATTTATGA	TAAAGACCCA	CTCAAAGATG	TGGACTTATC	AAAACCTTATG	GTTATTAAAG	12900

ACCATTCTTA CACAATTGAT ATGAATTATT GGGATGATAC TGACATCATA CATGCAATTT 12960
CAATATGTAC TGCAATTACA ATAGCAGATA CTATGTCACA ATTAGATCGA GATAATTTAA 13020
AAGAGATAAT AGTTATTGCA AATGATGATG ATATTAATAG CTTAATCACT GAATTTTGA 13080
CTCTTGACAT ACTTGTATTT CTCAAGACAT TTGGTGGATT ATTAGTAAAT CAATTTGCAT 13140
ACACTCTTTA TAGTCTAAAA ATAGAAGGTA GGGATCTCAT TTGGGATTAT ATAATGAGAA 13200
CACTGAGAGA TACTTCCCAT TCAATATTAA AAGTATTATC TAATGCATTA TCTCATCCTA 13260
AAGTATTCAA GAGGTTCTGG GATTGTGGAG TTTTAAACCC TATTTATGGG CCTAATATCG 13320
CTAGTCAAGA CCAGATAAAA CTTGCCCTAT CTATATGTGA ATATTCATA GATCTATTTA 13380
TGAGAGAATG GTTGAATGGT GTATCACTTG AAATATACAT TTGTGACAGC GATATGGAAG 13440
TTGCAAATGA TAGGAAACAA GCCTTTATTT CTAGACACCT TTCATTTGTT TGTGTTTAG 13500
CAGAAATTGC ATCTTTCGGA CCTAACCTGT TAAACTTAAC ATACTGGAG AGACTTGATC 13560
TATTGAAACA ATATCTTGAA TTAAATATTA AAGAAGACCC TACTCTTAAA TATGTACAAA 13620
TATCTGGATT ATTAATTAATA TCGTTCCCAT CAACTGTAAC ATACGTAAGA AAGACTGCAA 13680
TCAAATATCT AAGGATTTCG GGTATTAGTC CACCTGAGGT AATTGATGAT TGGGATCCGG 13740
TAGAAGATGA AAATATGCTG GATAACATTG TCAAACTAT AAATGATAAC TGTAATAAAG 13800
ATAATAAAGG GAATAAAATT AACAATTTCT GGGGACTAGC ACTTAAGAAC TATCAAGTCC 13860
TTAAAATCAG ATCTATAACA AGTGATTCTG ATGATAATGA TAGACTAGAT GCTAATACAA 13920
GTGGTTTGAC ACTTCCTCAA GGAGGGAATT ATCTATCGCA TCAATTGAGA TTATTCGGAA 13980
TCAACAGCAC TAGTTGTCTG AAAGCTCTTG AGTTATCACA AATTTTAATG AAGGAAGTCA 14040
ATAAAGACAA GGACAGGCTC TTCCTGGGAG AAGGAGCAGG AGCTATGCTA GCATGTTATG 14100
ATGCCACATT AGGACCTGCA GTTAATTATT ATAATTCAGG TTTGAATATA ACAGATGTAA 14160
TTGGTCAACG AGAATTGAAA ATATTTCTTT CAGAGGTATC ATTAGTAGGT AAAAAATTAG 14220
GAAATGTGAC ACAGATTCTT AACAGGGTAA AAGTACTGTT CAATGGGAAT CCTAATTCAA 14280
CATGGATAGG AAATATGGAA TGTGAGAGCT TAATATGGAG TGAATTAAAT GATAAGTCCA 14340
TTGGATTAGT ACATTGTGAT ATGGAAGGAG CTATCGGTAA ATCAGAAGAA ACTGTTCTAC 14400
ATGAACATTA TAGTGTTATA AGAATTACAT ACTTGATTGG GGATGATGAT GTTGTTTTAG 14460
TTTCCAAAAT TATACCTACA ATCACTCCGA ATTGGTCTAG AATACTTTAT CTATATAAAT 14520
TATATTGGAA AGATGTAAGT ATAATATCAC TCAAACTTC TAATCCTGCA TCAACAGAAT 14580
TATATCTAAT TTCGAAAGAT GCATATTGTA CTATAATGGA ACCTAGTGAA ATTGTTTTAT 14640
CAAACTTAA AAGATTGTCA CTCTTGGAAG AAAATAATCT ATTAAAATGG ATCATTTTAT 14700
CAAAGAAGAG GAATAATGAA TGGTTACATC ATGAAATCAA AGAAGGAGAA AGAGATTATG 14760
GAATCATGAG ACCATATCAT ATGGCACTAC AAATCTTTGG ATTTCAAATC AATTTAAATC 14820
ATCTGGCGAA AGAATTTTAA TCAACCCAG ATCTGACTAA TATCAACAAT ATAATCCAAA 14880
GTTTTCAGCG AACAATAAAG GATGTTTTAT TTGAATGGAT TAATATAACT CATGATGATA 14940
AGAGACATAA ATTAGGCGGA AGATATAACA TATTCCCACT GAAAAATAAG GGAAAGTTAA 15000

GACTGCTATC GAGAAGACTA GTATTAAGTT GGATTTTCATT ATCATTATCG ACTCGATTAC 15060
TTACAGGTCG CTTTCCTGAT GAAAAATTTG AACATAGAGC ACAGACTGGA TATGTATCAT 15120
TAGCTGATAC TGATTTAGAA TCATTAAAGT TATTGTCGAA AAACATCATT AAGAATTACA 15180
GAGAGTGTAT AGGATCAATA TCATATTGGT TTCTAACCAA AGAAGTTAAA ATACTTATGA 15240
AATTGATCGG TGGTGCTAAA TTATTAGGAA TTCCCAGACA ATATAAAGAA CCCGAAGACC 15300
AGTTATTAGA AAATAACAAT CAACATGATG AATTTGATAT CGATTAAAC ATAAATACAA 15360
TGAAGATATA TCCTAACCTT TATCTTTAAG CCTAGGAATA GACAAAAAGT AAGAAAAACA 15420
TGTAATATAT ATATACCAA CAGAGTTCTT CTCTTGTTTG GT 15462

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

TTGTCTGGGA AT 12

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TTGCCTGGGA AT 12

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

TTGTTTGGGA AT 12

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid

183

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

TTGTCTGGTA AT

12

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

AACTTTAAAT TA

12

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

AACTTAAAAT TA

12

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

TTAAAGACAT TG

12

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

TTTAAGACAT TG

12

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GCAGATGTCA AG

12

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GCAGATGCCA AG

12

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CGAATCTAAA GA

12

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CGAAGCTAAA GA

12

185

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

GAAATATTGA TC

12

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

GAAACATTGA TC

12

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

TCTCTACCCA AC

12

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

TCGTTAACCA AC

12

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid

186

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

AGTACAATAG GT

12

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

AGTACTGTGG GT

12

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

GCACTTGATC CA

12

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

AACTTGGATC CA

12

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

CCATCATTGT TGTGACAA

19

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

CCATCATTGT GGCTGACAA

19

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

TTACATGGCC A

11

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

TCACATGGCG A

11

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

TTTGACTGG GC

12

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

TTTTGATTGG GC

12

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

GGTCCTAATA CT

12

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

GGGCCTAATA TC

12

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74

CCATAGAGAG TCCATGGAAA GCGACGCTAA AAACATATC

38

WHAT IS CLAIMED IS:

1. An isolated polynucleotide molecule comprising an operably linked transcriptional promoter, a polynucleotide sequence encoding a PIV genome or antigenome, and a
5 transcriptional terminator, wherein said polynucleotide sequence encoding said PIV genome or antigenome is modified by introduction of a heterologous PIV sequence selected from a HPIV1 sequence, a HPIV2 sequence, a HPIV3 sequence, a BPIV sequence or a MPIV sequence to form a chimeric PIV genome or
10 antigenome.
2. The isolated polynucleotide molecule of claim 1, wherein a gene or gene segment of human PIV3 is replaced with a counterpart gene or gene segment from a heterologous PIV.
- 15 3. The isolated polynucleotide molecule of claim 2, wherein the counterpart gene or gene segment is a HN or F glycoprotein gene or gene segment of HPIV1 or HPIV2.
4. The isolated polynucleotide molecule of claim 2, wherein an HN or F glycoprotein gene of PIV1 or PIV2 is
20 substituted for the counterpart HN or F glycoprotein gene of HPIV3.
5. The isolated polynucleotide molecule of claim 1, wherein the polynucleotide sequence encoding the genome or antigenome incorporates a BPIV gene or gene segment.
- 25 6. The isolated polynucleotide molecule of claim 1, which incorporates a heterologous sequence from RSV.
7. The isolated polynucleotide molecule of claim 6, wherein the heterologous sequence from RSV is a G or F gene or gene segment.

8. The isolated polynucleotide molecule of claim 1, which incorporates a heterologous sequence from measles virus.

5 9. The isolated polynucleotide molecule of claim 8, wherein the heterologous sequence from measles virus is a HA or F gene or gene segment.

10 10. An isolated polynucleotide molecule comprising an operably linked transcriptional promoter, a polynucleotide sequence encoding a PIV genome or antigenome, and a transcriptional terminator, wherein said polynucleotide sequence encoding said PIV genome or antigenome is selected from the group consisting of:

15 i) p218(131) (SEQ ID NO: 1);
ii) p3/7(131) (SEQ ID NO: 14);
15 iii) p3/7(131)2G (SEQ ID NO: 15); or
iv) the isolated polynucleotide of i), ii) or iii) modified by introduction of a heterologous PIV sequence selected from a HPiV1 sequence, a HPiV2 sequence, a BPiV sequence or a MPiV sequence or by a nucleotide insertion,
20 rearrangement, deletion or substitution specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

25 11. An isolated polynucleotide molecule comprising an operably linked transcriptional promoter, a polynucleotide sequence encoding a PIV genome or antigenome, and a transcriptional terminator, wherein said polynucleotide sequence encoding said PIV genome or antigenome is modified by a nucleotide insertion, rearrangement, deletion or
30 substitution.

12. The isolated polynucleotide molecule of claim 11, wherein said nucleotide insertion, rearrangement, deletion or substitution specifies a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation,

small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

13. The isolated polynucleotide molecule of claim 12, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates multiple *ts* mutations.

14. The isolated polynucleotide molecule of claim 12, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates multiple non-*ts* attenuating mutations.

15. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates one or more mutations of JS cp45.

16. The isolated polynucleotide molecule of claim 15, wherein said polynucleotide sequence encoding said PIV genome or antigenome encodes at least one amino acid substitution in the polymerase L protein.

17. The isolated polynucleotide molecule of claim 16, wherein the amino acid substitution in the polymerase L protein occurs at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS cp45.

18. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome encodes at least one amino acid substitution in the N protein.

19. The isolated polynucleotide molecule of claim 28, wherein the amino acid substitution in the N protein occurs at a position corresponding to residues Val₉₆ or Ser₃₈₉ of JS cp45.

20. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome encodes an amino acid substitution in the C protein.

5 21. The isolated polynucleotide molecule of claim 20, wherein the amino acid substitution in the C protein occurs at a position corresponding to Ile₉₆ of JS cp45.

10 22. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome encodes at least one amino acid substitution in the F protein.

15 23. The isolated polynucleotide molecule of claim 22, wherein the amino acid substitution in the F protein occurs at a position corresponding to Ile₄₂₀ or Ala₄₅₀ of JS cp45.

24. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome encodes an amino acid substitution in the HN protein.

20 25. The isolated polynucleotide molecule of claim 24, wherein the amino acid substitution in the HN protein occurs at a position corresponding to residue Val₃₈₄ of JS cp45.

25 26. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates at least one mutation in a 3' leader sequence.

30 27. The isolated polynucleotide molecule of claim 26, wherein the mutation in the 3' leader occurs at a position corresponding to nucleotide 23, 24, 28, or 45 of JS cp45.

28. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates a mutation in a N gene start sequence.

5 29. The isolated polynucleotide molecule of claim 28, wherein the mutation in the N gene start sequence occurs at a position corresponding to nucleotide 62 of JS *cp45*.

10 30. The isolated polynucleotide molecule of claim 12, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates a plurality and up to a full complement of mutations present in *rcp45*, *rcp45* 3'NCMFHN, *rcp45* 3'NL, *rcp45* 3'N, or *rcp45* F.

15 31. The isolated polynucleotide molecule of claim 12, which is an antigenomic cDNA selected from *rcp45*, *rcp45* 3'NCMFHN, *rcp45* 3'NL, *rcp45* 3'N, *rcp45* L, *rcp45* F, *rcp45* M, *rcp45* HN, or *rcp45* C.

20 32. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates a mutation stabilized by multiple nucleotide substitutions in a codon specifying the mutation.

25 33. The isolated polynucleotide molecule of claim 11, wherein said polynucleotide sequence encoding said PIV genome or antigenome incorporates a heterologous sequence from HPIV1, HPIV2, HPIV3, BPIV or MPIV to form a chimeric genome or antigenome.

34. The isolated polynucleotide molecule of claim 33, wherein said chimeric genome or antigenome incorporates one or more *ts* mutations.

35. The isolated polynucleotide molecule of claim 33, wherein said chimeric genome or antigenome incorporates one or more non-ts attenuating mutations.

5 36. The isolated polynucleotide molecule of claim 33, wherein said chimeric genome or antigenome incorporates one or more mutations of JS cp45.

10 37. The isolated polynucleotide molecule of claim 36, wherein said one or more mutations of JS cp45 occur in one or more PIV proteins selected from L, M, N, C, F, or HN or in a PIV extragenic sequence selected from a 3' leader or N-gene start sequence.

15 38. The isolated polynucleotide molecule of claim 33, wherein said chimeric genome or antigenome incorporates multiple mutations each specifying a phenotype selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, or host range restriction.

20 39. The isolated polynucleotide molecule of claim 33, wherein said chimeric genome or antigenome incorporates at least one and up to a full complement of mutations present in rcp45, rcp45 3'NCMFHN, rcp45 3'NL, rcp45 3'N, or rcp45 F.

25 40. The isolated polynucleotide molecule of claim 33, wherein a mutation specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV is incorporated in a chimeric PIV background comprising a genome or antigenome having one or more PIV3 HN or F glycoprotein genes substituted by one or more counterpart PIV1 or PIV2 HN and F glycoprotein genes.

30 41. The isolated polynucleotide molecule of claim 33, wherein the heterologous sequence specifies a phenotypic alteration selected from attenuation, temperature-sensitivity,

cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of a chimeric PIV.

42. The isolated polynucleotide molecule of claim 11, which incorporates a cis-acting regulatory sequence of HPIV1, HPIV2, BPIV or MPIV.

43. The isolated polynucleotide molecule of claim 11, which incorporates a heterologous sequence from RSV.

44. The isolated polynucleotide molecule of claim 43, wherein the heterologous sequence from RSV is a G or F gene or gene segment.

45. The isolated polynucleotide molecule of claim 11, which incorporates a heterologous sequence from measles virus.

46. The isolated polynucleotide molecule of claim 45, wherein the heterologous sequence from measles virus is a HA or F gene or gene segment.

47. The isolated polynucleotide molecule of claim 11, which incorporates a polynucleotide sequence encoding a non-PIV molecule selected from a cytokine, a T-helper epitope, a restriction site marker, or a protein of a microbial pathogen capable of eliciting a protective immune response in a mammalian host.

48. A cell or cell-free composition including an expression vector which comprises an isolated polynucleotide molecule encoding a PIV genome or antigenome and an expression vector which comprises one or more isolated polynucleotide molecules that encode(s) N, P and L proteins of PIV, whereby expression of said PIV genome or antigenome and N, P, and L proteins yields an infectious PIV particle.

49. The cell or cell-free composition of claim 48, wherein the infectious PIV particle is a virus.

50. The cell or cell-free composition of claim 48, wherein the infectious PIV particle is a subviral particle.

5 51. The cell or cell-free composition of claim 48, wherein the polynucleotide encoding the PIV genome or antigenome and the one or more polynucleotides encoding N, P and L proteins of PIV are incorporated within a single vector.

10 52. A method for producing an infectious PIV particle from one or more isolated polynucleotide molecules encoding said PIV, comprising:

coexpressing in a cell or cell-free system an expression vector which comprises a polynucleotide molecule encoding a PIV genome or antigenome and an expression vector
15 which comprises one or more polynucleotide molecules encoding N, P and L proteins, thereby producing an infectious PIV particle.

20 53. The method of claim 52, wherein the PIV genome or antigenome and the N, P, and L proteins are expressed by the same expression vector.

54. The method of claim 52, wherein the N, P, and L proteins are encoded on two or three different expression vectors.

25 55. The method of claim 52, wherein at least one of the N, P and L proteins is supplied by coinfection with PIV.

56. The method of claim 52, wherein the polynucleotide molecule that encodes the PIV genome or antigenome is cDNA.

30 57. The method of claim 52, wherein the infectious PIV particle is a virus.

58. The method of claim 52, wherein the infectious PIV particle is a subviral particle.

59. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome
5 is a human, bovine or murine PIV sequence.

60. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome encodes the sequence of a wild-type PIV strain.

61. The method of claim 52, wherein the
10 polynucleotide molecule encoding the PIV genome or antigenome encodes HPIV3.

62. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome incorporates an attenuating mutation from a biologically
15 derived PIV strain.

63. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome incorporates one or more *ts* mutations.

64. The method of claim 52, wherein the
20 polynucleotide molecule encoding the PIV genome or antigenome incorporates one or more non-*ts* attenuating mutations.

65. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome incorporates at least one mutation of JS *cp45*.

25 66. The method of claim 65, wherein the polynucleotide molecule encoding the PIV genome or antigenome incorporates multiple mutations of JS *cp45*.

67. The method of claim 65, wherein the mutation of JS cp45 specifies at least one amino acid substitution in the polymerase L protein.

5 68. The method of claim 67, wherein the amino acid substitution in the polymerase L occurs at a position corresponding to Tyr₉₄₂, Leu₉₉₂, or Thr₁₅₅₈ of JS cp45.

69. The method of claim 65, wherein said mutation of JS cp45 specifies a change in a PIV protein selected from L, M, N, C, F, or HN or in a PIV extragenic sequence selected from a 3' leader or N gene start sequence.

10 70. The method of claim 52, wherein said polynucleotide molecule encoding the PIV genome or antigenome incorporates a mutation that is stabilized by multiple nucleotide substitutions in a codon which specifies the mutation.

15 71. The method of claim 52, wherein said polynucleotide molecule encoding said PIV genome or antigenome incorporates a plurality and up to a full complement of mutations present in rcp45, rcp45 3'NCMFHN, rcp45 3'NL, rcp45 3'N, or rcp45 F.

20 72. The method of claim 69, wherein said polynucleotide molecule encoding said PIV genome or antigenome is an antigenomic cDNA selected from rcp45, rcp45 3'NCMFHN, rcp45 3'NL, rcp45 3'N, rcp45 L, rcp45 F, rcp45 M, rcp45 HN, or rcp45 C.

25 73. The method of claim 52, wherein said polynucleotide molecule encoding said PIV genome or antigenome incorporates a heterologous sequence from HPIV1, HPIV2, HPIV3, BPIV or MPIV to form a chimeric genome or antigenome.

30 74. The method of claim 73, wherein the polynucleotide molecule encoding the PIV genome or antigenome

is a chimera of a HPIV3 sequence and a HPIV1, HPIV2, BPIV or MPIV sequence.

75. The method of claim 74, wherein a heterologous sequence from HPIV1 or HPIV2 encoding a gene or gene segment
5 of an HN or F glycoprotein is substituted for a corresponding gene or gene segment of HPIV3.

76. The method of claim 73, wherein said chimeric genome or antigenome incorporates one or more *ts* mutations.

77. The method of claim 73, wherein said chimeric
10 genome or antigenome incorporates one or more non-*ts* attenuating mutations.

78. The method of claim 73, wherein said chimeric genome or antigenome incorporates one or more mutations of JS *cp45*.

79. The method of claim 73, wherein said chimeric
15 genome or antigenome incorporates multiple mutations each specifying a phenotype selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, or host range restriction.

80. The method of claim 73, wherein a mutation
20 specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV is incorporated in a chimeric PIV background comprising
25 a genome or antigenome having one or more PIV3 HN or F glycoprotein genes substituted by one or more counterpart PIV1 or PIV2 HN and F glycoprotein genes.

81. The method of claim 80, wherein one or more
30 mutations of JS *cp45* are incorporated in a chimeric background comprising a genome or antigenome having both PIV3 HN and F

glycoprotein genes substituted by counterpart PIV1 or PIV2 HN and F glycoprotein genes.

82. The method of claim 81, wherein said one or more mutations of JS cp45 occur in one or more PIV proteins selected from L, M, N, C, F, or HN or in a PIV extragenic sequence selected from a 3' leader or N gene start sequence.

83. The method of claim 73, wherein the heterologous sequence specifies a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of a chimeric PIV.

84. The method of claim 52, wherein said polynucleotide molecule encoding said PIV genome or antigenome incorporates a heterologous sequence from RSV.

85. The method of claim 84, wherein the heterologous sequence from RSV is a G or F gene or gene segment.

86. The method of claim 52, wherein said polynucleotide molecule encoding said PIV genome or antigenome incorporates a heterologous sequence from measles virus.

87. The method of claim 86, wherein the heterologous sequence from measles virus is a HA or F gene or gene segment.

88. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome is selected from:

- i) p218(131) (SEQ ID NO: 1);
- ii) p3/7(131) (SEQ ID NO: 14);
- iii) p3/7(131)2G (SEQ ID NO: 15); or
- iv) the polynucleotide molecule of i), ii) or iii) modified by introduction of a heterologous PIV sequence

selected from a HPIV1 sequence, a HPIV2 sequence, a BPIV sequence or a MPIV sequence or by a nucleotide insertion, rearrangement, deletion or substitution specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

89. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome is selected from:

- i) p218(131) (SEQ ID NO: 1);
 - ii) p3/7(131) (SEQ ID NO: 14);
 - iii) p3/7(131)2G (SEQ ID NO: 15); or
 - iv) the polynucleotide molecule of i), ii) or iii)
- modified by introduction of a heterologous PIV sequence selected from a HPIV1 sequence, a HPIV2 sequence, a BPIV sequence or a MPIV sequence and by a nucleotide insertion, rearrangement, deletion or substitution different from said introduction of said heterologous PIV sequence specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

90. The method of claim 52, wherein the polynucleotide molecule encoding the PIV genome or antigenome is modified to encode a non-PIV molecule selected from a cytokine, a T-helper epitope, a restriction site marker, or a protein of a microbial pathogen capable of eliciting a protective immune response in a mammalian host.

91. An isolated infectious PIV particle which comprises a recombinant PIV genome or antigenome, a N protein, a P protein, and a L protein.

92. The isolated infectious PIV particle of claim 91, which is a subviral particle.

93. The isolated infectious PIV particle of claim 91, which is a virus.

94. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome
5 incorporates a heterologous sequence from RSV or measles virus.

95. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome is a cDNA.

10 96. The isolated infectious PIV particle of claim 91, which is a human PIV.

97. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome is a chimera of heterologous PIV sequences selected from HPIV1,
15 HPIV2, HPIV3, BPIV, or MPIV sequences.

98. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome is selected from:

20 i) p218(131) (SEQ ID NO: 1);
ii) p3/7(131) (SEQ ID NO: 14);
iii) p3/7(131)2G (SEQ ID NO: 15); or
iv) the genome or antigenome of i), ii) or iii)
modified by introduction of a heterologous PIV sequence
selected from a HPIV1 sequence, a HPIV2 sequence, a BPIV
25 sequence or a MPIV sequence or by a nucleotide insertion,
rearrangement, deletion or substitution specifying a
phenotypic alteration selected from attenuation, temperature-
sensitivity, cold-adaptation, small plaque size, host range
restriction, or a change in an immunogenic epitope of PIV.

30 99. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome is selected from:

- i) p218(131) (SEQ ID NO: 1);
 - ii) p3/7(131) (SEQ ID NO: 14);
 - iii) p3/7(131)2G (SEQ ID NO: 15); or
 - iv) the genome or antigenome of i), ii) or iii)
- 5 modified by introduction of a heterologous PIV sequence selected from a HPIV1 sequence, a HPIV2 sequence, a BPIV sequence or a MPIV sequence and by a nucleotide insertion, rearrangement, deletion or substitution different from said
- 10 introduction of said heterologous PIV sequence specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

15 100. The isolated infectious PIV particle of claim 91, wherein the counterpart gene or gene segment is a gene or gene segment of the HN or F glycoprotein gene of HPIV1 or HPIV2.

20 101. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome incorporates a heterologous sequence from RSV or measles virus.

25 102. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome is modified by a nucleotide insertion, rearrangement, deletion or substitution encoding a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV.

30 103. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome incorporates multiple ts mutations.

104. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome incorporates multiple non-ts attenuating mutations.

105. The isolated infectious PIV particle of claim 91, wherein the recombinant PIV genome or antigenome incorporates at least one mutation of JS cp45.

106. The isolated infectious PIV particle of claim 105, wherein the mutation of JS cp45 specifies an amino acid substitution in the polymerase L protein.

107. The isolated infectious PIV particle of claim 97, wherein said chimeric genome or antigenome incorporates one or more *ts* mutations.

108. The isolated infectious PIV particle of claim 97, wherein said chimeric genome or antigenome incorporates one or more non-*ts* attenuating mutations.

109. The isolated infectious PIV particle of claim 97, wherein said chimeric genome or antigenome incorporates one or more mutations of JS cp45.

110. The isolated infectious PIV particle of claim 129, wherein said chimeric genome or antigenome incorporates multiple mutations each specifying a phenotype selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, or host range restriction.

111. The isolated infectious PIV particle of claim 109, wherein said chimeric genome or antigenome incorporates at least one and up to a full complement of mutations present in *rcp45*, *rcp45* 3'NCMFHN, *rcp45* 3'NL, *rcp45* 3'N, or *rcp45* F.

112. The isolated infectious PIV particle of claim 91, wherein a mutation specifying a phenotypic alteration selected from attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of PIV is incorporated in a chimeric PIV background comprising a genome or antigenome having one or more PIV3 HN or F glycoprotein genes or gene

segments substituted by one or more counterpart PIV1 or PIV2
HN and F glycoprotein genes or gene segments.

5 113. The isolated infectious PIV particle of claim
112, wherein one or more mutations of JS cp45 are incorporated
in said chimeric background.

10 114. The isolated infectious PIV particle of claim
of claim 113, wherein said one or more mutations of JS cp45
occur in one or more PIV proteins selected from L, M, N, C, F,
or HN or in a PIV extragenic sequence selected from a 3'
leader or N gene start sequence.

15 115. The isolated infectious PIV particle of claim
91, wherein the recombinant PIV genome or antigenome is
modified to encode a non-PIV molecule selected from a
cytokine, a T-helper epitope, a restriction site marker, or a
protein of a microbial pathogen capable of eliciting a
protective immune response in a mammalian host.

116. The isolated infectious PIV particle of claim
91, further comprising an RSV antigen or epitope which elicits
protective immunity to RSV in an immunized host.

20 117. The isolated infectious PIV particle of claim
91, which is selected from r942, r992, r1558, r942/992,
r992/1558, r942/1558, or r942/992/1558, rcp45 3'N, rcp45 C,
rcp45 M, rcp45 F, rcp45 HN, rcp45L, rcp45 3'NL, rcp45
3'NCMFHN, and rcp45.

25 118. An immunogenic composition comprising an
immunogenically effective amount of an infectious PIV particle
in a pharmaceutically acceptable carrier, said PIV particle
comprising a recombinant PIV genome or antigenome, a N
protein, a P protein, and a L protein.

30 119. The immunogenic composition of claim 118,
wherein said infectious PIV particle is a subviral particle.

120. The immunogenic composition of claim 118,
wherein said infectious PIV particle is a virus.

121. The immunogenic composition of claim 118,
wherein the recombinant PIV genome or antigenome incorporates
5 a heterologous sequence from RSV or measles virus.

122. The immunogenic composition of claim 118,
wherein the recombinant PIV genome or antigenome is a chimera
of heterologous PIV sequences selected from HPIV1, HPIV2,
HPIV3, BPIV, or MPIV sequences to form an infectious, chimeric
10 PIV particle.

123. The immunogenic composition of claim 118,
wherein the recombinant PIV genome or antigenome encodes a
human PIV in which a gene or gene segment is replaced with a
counterpart gene or gene segment from a heterologous PIV.

124. The immunogenic composition of claim 123,
wherein one or both HN and F glycoprotein genes of HPIV1 are
substituted for HN and F glycoprotein genes of HPIV3 to form
said infectious, chimeric PIV particle.

125. The immunogenic composition of claim 123,
20 wherein the recombinant PIV genome or antigenome of said
infectious, chimeric PIV particle is modified by a nucleotide
insertion, rearrangement, deletion or substitution encoding a
phenotypic alteration selected from attenuation, temperature-
sensitivity, cold-adaptation, small plaque size, host range
25 restriction, or a change in an immunogenic epitope of PIV.

126. The immunogenic composition of claim 125,
wherein said recombinant PIV genome or antigenome incorporates
multiple mutations selected from *ts* and non-*ts* attenuating
mutations to form an attenuated, infectious, chimeric PIV
30 particle.

127. The immunogenic composition of claim 118,
wherein the recombinant PIV genome or antigenome incorporates
a mutation of JS cp45.

128. The immunogenic composition of claim 118,
5 wherein the recombinant PIV genome or antigenome incorporates
multiple mutations of JS cp45.

1/18

FIG. 1




FIG. 14

FIG. 14A

FIG. 14B

FIG. 2





FIG. 15A



3/18

FIG. 3





FIG. 4



4/18

FIG. 5A





FIG. 5B



SUBSTITUTE SHEET (RULE 26)

5/18

FIG. 6A





FIG. 6B



6/18

FIG. 7A





FIG. 7B



7/18

FIG. 8






FIG. 12




8/18

FIG. 9



9/18

FIG. 10



10/18




FIG. 11

FIG. 13

cp 45 MUTATIONS INTRODUCED INTO rPIV3





Figure 1: Schematic representation of recombinant PIV3 viruses bearing mutations (*) in the 3' leader, N gene start, N, C, M, F, HN and/or L proteins.

12/18

FIG. 14A



SUBSTITUTE SHEET (RULE 26)





FIG. 14B

14/18

FIG. 15B



15/18

FIG. 16

REGION I		F ORF	
rPIV3	5039	AAGAGACCGGCAACACAAAGCAGCTGAACACAAATGCCAACTTCAATACCTCTAATTAATACAACCAATG	5107
rPIV3-1	5039	AAGAGACCGGCAACACAAAGCAGCTGAACACCAATGGAATAATCAGAGATCCCTCTCTTAGTATACTCA	5107
PIV1		AAATTCAATAAACAACCAAGATAGAGACAAAATGCCAAAATCAGAGATCCCTCTCTTAGTATACTCA	
REGION II		F ORF	
		Gln to Glu	
rPIV3	6659	AATGACAAGCCATATGTACTAACAAACAATAACATATCTACAGATCATTAGATTAATAAATTAATAAAA	6727
rPIV3-1	6707	AGAAATCCCTACATGGGTAACAACTCCAAATAGCATCCCTACAGATCATTAGATTAATAAATTAATAAAA	6775
PIV1		AGAAATCCCTACATGGGTAACAACTCCAAATTAATAAATCAGATCAAGTACATTGTAGCATACATACAAC	
REGION III		HN ORF	
rPIV3	6773	ACGAAGGACCCCAATAGACAAATCCAAATTCGAGATGCAATACIGGAAGCAIACCAATCAGGAAGCAT	6841
rPIV3-1	6821	ACGAAGGACCCCAATAGACAAATCCAAATCCATGGCTGAAAAAGGGAAACAAATAGTTCATAATTGG	6877
PIV1		GAGCTCTGGTACCAGCATCCGATTATACAAACCAATGGCTGAAAAAGGGAAACAAATAGTTCATAATTGG	
REGION IV		HN ORF	
rPIV3	8486	ATGTTGTTCAAAACAGAGATTCCAAAAAGCTCCAGTTAATCATAATTAACCATAAATGCAATCAATCTA	8554
rPIV3-1	8540	AAGACAAGTATCCCTAAAATATGTAATAATCACAATCTAAGCTTAATTAACCATAAATGCAATCAATCTA	8608
PIV1		AAGACAAGTATCCCTAAAATATGTAATAATCACAATCTAAGCTTAAGGCGCTTAAGCTAGTCTGTAAT	

-16/18-




FIG. 17B




FIG. 17A

17/18

FIG. 18





FIG. 19



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/10551

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/45 C07K14/115 C12N5/10 C12N7/01 A61K39/155

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>WO 97 20468 A (UNIV ST LOUIS ; BELSHE ROBERT B (US); RAY RANJIT (US)) 12 June 1997</p> <p>see abstract see page 25, line 24 - page 27, line 27 see claims 1-20</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1-17, 26, 27, 30, 31, 33-69, 71-116, 118-128</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

30 September 1998

Date of mailing of the international search report

13/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

International Application No
PCT 98/10551

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		Relevant to claim No.
Category	Citation of document, with indication where appropriate, of the relevant passages	
X	DIMOCK K. & COLLINS P.L.: "Rescue of synthetic analogs of genomic RNA and replicative intermediate RNA of human parainfluenza virus type 3" J. VIROL., vol. 67, no. 5, May 1993, pages 2772-2778, XP002078295 cited in the application see figure 1	11,47, 52,53, 55-57, 59-61, 90,91, 93,95, 96,115
Y	EP 0 702 085 A (AKZO NOBEL NV) 20 March 1996	10-17, 22-27, 30,31, 48-72, 88, 91-98, 102-106, 117-120, 127,128
Y	see the whole document --- STOKES A ET AL: "THE COMPLETE NUCLEOTIDE SEQUENCE OF TWO COLD-ADAPTED, TEMPERATURE-SENSITIVE ATTENUATED MUTANT VACCINE CIRUSES (CP12 AND CP45) DERIVED FROM THE JS STRAIN OF HUMAN PARAINFLUENZA VIRUS TYPE 3 (PIV3)" VIRUS RESEARCH, vol. 30, 1993, pages 43-52, XP002051711 cited in the application see the whole document	10-17, 22-27, 30,31, 48-72, 88, 91-98, 102-106, 117-120, 127,128
A	& DATABASE GENBANK Accession No. U51116, 6 March 1997 STOKES A.: "Human parainfluenza virus 3" see the whole document	18-21, 28,29,32
A	--- STOKES A. ET AL.: "The complete nucleotide sequence of the JS strain of human parainfluenza virus type 3: comparison with the Wash/47885/57 prototype strain" VIRUS RESEARCH, vol. 25, 1992, pages 91-103, XP002078296 see the whole document & DATABASE GENBANK Accession No. Z11575, 23 August 1993 STOKES A.: "Human parainfluenza virus type 3 (wt)" see page P	1-128
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/10551

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>COLLINS P L ET AL: "PRODUCTION OF INFECTIOUS HUMAN RESPIRATORY SYNCYTIAL VIRUS FROM CLONED CDNA CONFIRMS AN ESSENTIAL ROLE FOR THE TRANSCRIPTION ELONGATION FACTOR FROM THE 5' PROXIMAL OPEN READING FRAME OF THE M2 MRNA IN GENE EXPRESSION AND PROVIDES A CAPABILITY FOR VACCINE DEVELOPMENT"</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 92, December 1995, pages 11563-11567, XP002066592 see figure 1</p> <p>---</p>	<p>1,6,7, 11,43, 44,52, 84,85, 91,94, 101,116, 118,121</p>
A	<p>EP 0 440 219 A (SCHWEIZ SERUM & IMPFINST) 7 August 1991</p> <p>see abstract</p> <p>---</p>	<p>8,9,11, 45,46, 52,86, 87,91, 94,101, 118,121</p>
A	<p>WO 97 11093 A (CONNAUGHT LAB ;CATES GEORGE A (CA); EWASYSHYN MARY E (CA); FAHIM R) 27 March 1997</p> <p>see abstract</p> <p>---</p>	<p>1-11,30, 33-62, 73-101, 107-114, 118-126</p>
P,X	<p>TAO T. ET AL.: "Recovery of a fully viable chimeric human parainfluenza virus (PIV) type 3 in which the hemagglutinin-Neuraminidase and Fusion glycoproteins have been replaced by those of PIV type 1"</p> <p>J. VIROL., vol. 72, no. 4, April 1998, pages 2955-2961, XP002078294 see the whole document</p> <p>---</p>	<p>1-17, 22-27, 30-128</p>
P,X	<p>SKIADOPOULOS M.H. ET AL.: "Three amino acid substitutions in the L protein of the human parainfluenzavirus type 3 cp45 live attenuated vaccine candidate contribute to its temperature-sensitive and attenuation phenotypes."</p> <p>J. VIROLOGY, vol. 72, no. 3, March 1998, pages 1762-1768, XP002078298 see the whole document</p> <p>---</p> <p style="text-align: center;">-/--</p>	<p>10-17, 48-69, 71,72, 88, 91-93, 95-98, 102-106, 117</p>

INTERNATIONAL SEARCH REPORT

International Application No
PCT/98/10551

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P;X	<p>DURBIN A.P. ET AL.: "Recovery of infectious human parainfluenza virus type 3 from cDNA" VIROLOGY, vol. 235, 1 September 1997, pages 323-332, XP002078299 see the whole document</p>	<p>10, 52-61, 88,89, 91-93, 96,98,99</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

CT/US 98/10551

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 9720468	A	12-06-1997	AU	1149597 A		27-06-1997
EP 0702085	A	20-03-1996	CA	2154023 A		19-01-1996
			CN	1121959 A		08-05-1996
			JP	8168381 A		02-07-1996
EP 0440219	A	07-08-1991	AU	7007491 A		08-08-1991
			CA	2035386 A		03-08-1991
			JP	4211377 A		03-08-1992
WO 9711093	A	27-03-1997	AU	6982096 A		09-04-1997
			EP	0851873 A		08-07-1998